

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 October 2001 (18.10.2001)

PCT

(10) International Publication Number
WO 01/77330 A2

- (51) International Patent Classification⁷: C12N 15/12, C07K 14/705, C12N 15/86, 5/10, A61K 48/00, 38/16, C07K 16/28, G01N 33/68, C12Q 1/68, C12N 15/11 // A61P 25/28
- (74) Agents: DELUCA, Mark et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).
- (21) International Application Number: PCT/US01/11330
- (22) International Filing Date: 6 April 2001 (06.04.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|-------------------------------|----|
| 60/195,150 | 6 April 2000 (06.04.2000) | US |
| 60/195,099 | 6 April 2000 (06.04.2000) | US |
| 60/195,151 | 6 April 2000 (06.04.2000) | US |
| 60/195,148 | 6 April 2000 (06.04.2000) | US |
| 60/195,093 | 6 April 2000 (06.04.2000) | US |
| 60/195,098 | 6 April 2000 (06.04.2000) | US |
| 60/230,149 | 5 September 2000 (05.09.2000) | US |
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:
— of inventorship (Rule 4.17(iv)) for US only
- (71) Applicant (*for all designated States except US*): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).
- Published:
— without international search report and to be republished upon receipt of that report
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): VOGELI, Gabriel [US/US]; 3005 First Avenue, Seattle, WA 98121 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/77330 A2

(54) Title: NOVEL G PROTEIN-COUPLED RECEPTORS

(57) Abstract: The present invention provides a gene encoding a G protein-coupled receptor termed nGPCR-x; constructs and recombinant host cells incorporating the genes; the nGPCR-x polypeptides encoded by the gene; antibodies to the nGPCR-x polypeptides; and methods of making and using all of the foregoing.

NOVEL G PROTEIN-COUPLED RECEPTORS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application claims priority of Application Serial No. 60/195,150, filed 2000 April 6; Application Serial No. 60/195,099 filed 2000 April 6; Application Serial No. 60/195,151 filed 2000 April 6; Application Serial No. 60/195,148 filed 2000 April 6; Application Serial No. 60/195,093 filed 2000 April 6; Application Serial No. 60/195,098 filed 2000 April 6; and Application Serial No. 60/230,149 filed 2000 September 5; each of which is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0002] The present invention relates generally to the fields of genetics and cellular and molecular biology. More particularly, the invention relates to novel G protein coupled receptors, to polynucleotides that encode such novel receptors, to reagents such as antibodies, probes, primers and kits comprising such antibodies, probes, primers related to the same, and to methods which use the novel G protein coupled receptors, polynucleotides or reagents.

BACKGROUND OF THE INVENTION

[0003] The G protein-coupled receptors (GPCRs) form a vast superfamily of cell surface receptors which are characterized by an amino-terminal extracellular domain, a carboxyl-terminal intracellular domain, and a serpentine structure that passes through the cell membrane seven times. Hence, such receptors are sometimes also referred to as seven transmembrane (7TM) receptors. These seven transmembrane domains define three extracellular loops and three intracellular loops, in addition to the amino- and carboxy-terminal domains. The extracellular portions of the receptor have a role in recognizing and binding one or more extracellular binding partners (*e.g.*, ligands), whereas the intracellular portions have a role in recognizing and communicating with downstream molecules in the signal transduction cascade.

[0004] The G protein-coupled receptors bind a variety of ligands including calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and even photons, and are important in the normal (and sometimes the aberrant) function of many cell types. [See generally Strosberg, *Eur. J. Biochem.* 196:1-10 (1991) and Bohmet *et al.*, *Biochem J.* 322:1-18 (1997).] When a specific ligand binds to its corresponding receptor, the ligand typically stimulates the receptor to activate a specific heterotrimeric guanine-nucleotide-binding

regulatory protein (G-protein) that is coupled to the intracellular portion of the receptor. The G protein in turn transmits a signal to an effector molecule within the cell, by either stimulating or inhibiting the activity of that effector molecule. These effector molecules include adenylate cyclase, phospholipases and ion channels. Adenylate cyclase and phospholipases are enzymes that are involved in the production of the second messenger molecules cAMP, inositol triphosphate and diacylglycerol. It is through this sequence of events that an extracellular ligand stimuli exerts intracellular changes through a G protein-coupled receptor. Each such receptor has its own characteristic primary structure, expression pattern, ligand-binding profile, and intracellular effector system.

[0005] Because of the vital role of G protein-coupled receptors in the communication between cells and their environment, such receptors are attractive targets for therapeutic intervention, for example by activating or antagonizing such receptors. For receptors having a known ligand, the identification of agonists or antagonists may be sought specifically to enhance or inhibit the action of the ligand. Some G protein-coupled receptors have roles in disease pathogenesis (*e.g.*, certain chemokine receptors that act as HIV co-receptors may have a role in AIDS pathogenesis), and are attractive targets for therapeutic intervention even in the absence of knowledge of the natural ligand of the receptor. Other receptors are attractive targets for therapeutic intervention by virtue of their expression pattern in tissues or cell types that are themselves attractive targets for therapeutic intervention. Examples of this latter category of receptors include receptors expressed in immune cells, which can be targeted to either inhibit autoimmune responses or to enhance immune responses to fight pathogens or cancer; and receptors expressed in the brain or other neural organs and tissues, which are likely targets in the treatment of mental disorder, depression, bipolar disease, or other neurological disorders. This latter category of receptor is also useful as a marker for identifying and/or purifying (*e.g.*, via fluorescence-activated cell sorting) cellular subtypes that express the receptor. Unfortunately, only a limited number of G protein receptors from the central nervous system (CNS) are known. Thus, a need exists for G protein-coupled receptors that have been identified and show promise as targets for therapeutic intervention in a variety of animals, including humans.

SUMMARY OF THE INVENTION

[0006] The present invention relates to an isolated nucleic acid molecule that comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, or a fragment thereof. The nucleic acid molecule encodes at least a portion of nGPCR α . In some embodiments, the nucleic acid molecule comprises a sequence that encodes a polypeptide

comprising a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, or a fragment thereof. In some embodiments, the nucleic acid molecule comprises a sequence homologous to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or a fragment thereof. In some embodiments, the nucleic acid molecule comprises a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, and fragments thereof.

[0007] According to some embodiments, the present invention provides vectors which comprise the nucleic acid molecule of the invention. In some embodiments, the vector is an expression vector.

[0008] According to some embodiments, the present invention provides host cells which comprise the vectors of the invention. In some embodiments, the host cells comprise expression vectors.

[0009] The present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence complementary to at least a portion of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, said portion comprising at least 10 nucleotides.

[00010] The present invention provides a method of producing a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, or a homolog or fragment thereof. The method comprising the steps of introducing a recombinant expression vector that includes a nucleotide sequence that encodes the polypeptide into a compatible host cell, growing the host cell under conditions for expression of the polypeptide and recovering the polypeptide.

[00011] The present invention provides an isolated antibody which binds to an epitope on a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, or a homolog or fragment thereof.

[00012] The present invention provides a method of inducing an immune response in a mammal against a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, or a homolog or fragment thereof. The method comprises administering to a mammal an amount of the polypeptide sufficient to induce said immune response.

[00013] The present invention provides a method for identifying a compound which binds nGPCR-x. The method comprises the steps of contacting nGPCR-x with a compound and determining whether the compound binds nGPCR-x.

[00014] The present invention provides a method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-x. The method comprises the steps of contacting said

nucleic acid molecule encoding nGPCR-x with a compound and determining whether said compound binds said nucleic acid molecule.

- [00015] The present invention provides a method for identifying a compound which modulates the activity of nGPCR-x. The method comprises the steps of contacting nGPCR-x with a compound and determining whether nGPCR-x activity has been modulated.
- [00016] The present invention provides a method of identifying an animal homolog of nGPCR-x. The method comprises the steps screening a nucleic acid database of the animal with a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or a portion thereof and determining whether a portion of said library or database is homologous to said sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or portion thereof.
- [00017] The present invention provides a method of identifying an animal homolog of nGPCR-x. The methods comprises the steps screening a nucleic acid library of the animal with a nucleic acid molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or a portion thereof, and determining whether a portion of said library or database is homologous to said sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or a portion thereof.
- [00018] Another aspect of the present invention relates to methods of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor. The methods comprise the steps of assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR-x that is expressed in the brain. The nGPCR-x comprise an amino acid sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, and allelic variants thereof. A diagnosis of the disorder or predisposition is made from the presence or absence of the mutation. The presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR-x in the nucleic acid correlates with an increased risk of developing the disorder.
- [00019] The present invention further relates to methods of screening for a nGPCR-x hereditary mental disorder genotype in a human patient. The methods comprise the steps of providing a biological sample comprising nucleic acid from the patient, in which the nucleic acid includes sequences corresponding to alleles of nGPCR-x. The presence of one or more mutations in the nGPCR-x allele is indicative of a hereditary mental disorder genotype.
- [00020] The present invention provides kits for screening a human subject to diagnose mental disorder or a genetic predisposition therefor. The kits include an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-x gene. The oligonucleotide comprises 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a

wild type human nGPCR-x gene sequence or nGPCR-x coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution. The kit also includes a media packaged with the oligonucleotide. The media contains information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

[00021] The present invention further relates to methods of identifying nGPCR-x allelic variants that correlates with mental disorders. The methods comprise the steps of providing biological samples that comprise nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny, and detecting in the nucleic acid the presence of one or more mutations in an nGPCR-x that is expressed in the brain. The nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, and allelic variants thereof. The nucleic acid includes sequences corresponding to the gene or genes encoding nGPCR-x. The one or more mutations detected indicate an allelic variant that correlates with a mental disorder.

[00022] The present invention further relates to purified polynucleotides comprising nucleotide sequences encoding alleles of nGPCR-x from a human with mental disorder. The polynucleotide hybridizes to the complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 under the following hybridization conditions: (a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS. The polynucleotide that encodes nGPCR-x amino acid sequence of the human differs from a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116 by at least one residue.

[00023] The present invention also provides methods for identifying a modulator of biological activity of nGPCR-x comprising the steps of contacting a cell that expresses nGPCR-x in the presence and in the absence of a putative modulator compound and measuring nGPCR-x biological activity in the cell. The decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

[00024] The present invention further provides methods to identify compounds useful for the treatment of mental disorders. The methods comprise the steps of contacting a composition comprising nGPCR-x with a compound suspected of binding nGPCR-x. The binding between nGPCR-x and the compound suspected of binding nGPCR-x is detected. Compounds identified as binding nGPCR-x are candidate compounds useful for the treatment of mental disorder.

Compounds identified as binding nGPCR-x may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity.

[00025] The present invention further provides methods for identifying a compound useful as a modulator of binding between nGPCR-x and a binding partner of nGPCR-x. The methods comprise the steps of contacting the binding partner and a composition comprising nGPCR-x in the presence and in the absence of a putative modulator compound and detecting binding between the binding partner and nGPCR-x. Decreased or increased binding between the binding partner and nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a related disease or disorder. Compounds identified as modulating binding between nGPCR-x and a nGPCR-x binding partner may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity as modulators.

[00026] Another aspect of the present invention relates to methods of purifying a G protein from a sample containing a G protein. The methods comprise the steps of contacting the sample with an nGPCR-x for a time sufficient to allow the G protein to form a complex with the nGPCR-x; isolating the complex from remaining components of the sample; maintaining the complex under conditions which result in dissociation of the G protein from the nGPCR-x; and isolating said G protein from the nGPCR-x.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Definitions

[00027] Various definitions are made throughout this document. Most words have the meaning that would be attributed to those words by one skilled in the art. Words specifically defined either below or elsewhere in this document have the meaning provided in the context of the present invention as a whole and as are typically understood by those skilled in the art.

[00028] "Synthesized" as used herein and understood in the art, refers to polynucleotides produced by purely chemical, as opposed to enzymatic, methods. "Wholly" synthesized DNA sequences are therefore produced entirely by chemical means, and "partially" synthesized DNAs embrace those wherein only portions of the resulting DNA were produced by chemical means.

[00029] By the term "region" is meant a physically contiguous portion of the primary structure of a biomolecule. In the case of proteins, a region is defined by a contiguous portion of the amino acid sequence of that protein.

[00030] The term "domain" is herein defined as referring to a structural part of a biomolecule that contributes to a known or suspected function of the biomolecule. Domains may be co-extensive with regions or portions thereof; domains may also incorporate a portion of a

biomolecule that is distinct from a particular region, in addition to all or part of that region . Examples of GPCR protein domains include, but are not limited to, the extracellular (*i.e.*, N-terminal), transmembrane and cytoplasmic (*i.e.*, C-terminal) domains, which are co-extensive with like-named regions of GPCRs; each of the seven transmembrane segments of a GPCR; and each of the loop segments (both extracellular and intracellular loops) connecting adjacent transmembrane segments.

[00031] As used herein, the term "activity" refers to a variety of measurable indicia suggesting or revealing binding, either direct or indirect; affecting a response, *i.e.* having a measurable affect in response to some exposure or stimulus, including, for example, the affinity of a compound for directly binding a polypeptide or polynucleotide of the invention, or, for example, measurement of amounts of upstream or downstream proteins or other similar functions after some stimulus or event.

[00032] Unless indicated otherwise, as used herein, the abbreviation in lower case (*gpcr*) refers to a gene, cDNA, RNA or nucleic acid sequence, while the upper case version (*GPCR*) refers to a protein, polypeptide, peptide, oligopeptide, or amino acid sequence. The term "nGPCR-x" refers to any of the nGPCRs taught herein, while specific reference to a nGPCR (for example nGPCR-2073) refers only to that specific nGPCR.

[00033] As used herein, the term "antibody" is meant to refer to complete, intact antibodies, and Fab, Fab', F(ab)2, and other fragments thereof. Complete, intact antibodies include monoclonal antibodies such as murine monoclonal antibodies, chimeric antibodies and humanized antibodies.

[00034] As used herein, the term "binding" means the physical or chemical interaction between two proteins or compounds or associated proteins or compounds or combinations thereof. Binding includes ionic, non-ionic, Hydrogen bonds, Van der Waals, hydrophobic interactions, etc. The physical interaction, the binding, can be either direct or indirect, indirect being through or due to the effects of another protein or compound. Direct binding refers to interactions that do not take place through or due to the effect of another protein or compound but instead are without other substantial chemical intermediates. Binding may be detected in many different manners. As a non-limiting example, the physical binding interaction between a nGPCR-x of the invention and a compound can be detected using a labeled compound. Alternatively, functional evidence of binding can be detected using, for example, a cell transfected with and expressing a nGPCR-x of the invention. Binding of the transfected cell to a ligand of the nGPCR-x that was transfected into the cell provides functional evidence of binding. Other methods of detecting binding are well known to those of skill in the art.

- [00035] As used herein, the term "compound" means any identifiable chemical or molecule, including, but not limited to, small molecule, peptide, protein, sugar, nucleotide, or nucleic acid, and such compound can be natural or synthetic.
- [00036] As used herein, the term "complementary" refers to Watson-Crick basepairing between nucleotide units of a nucleic acid molecule.
- [00037] As used herein, the term "contacting" means bringing together, either directly or indirectly, a compound into physical proximity to a polypeptide or polynucleotide of the invention. The polypeptide or polynucleotide can be in any number of buffers, salts, solutions *etc.* Contacting includes, for example, placing the compound into a beaker, microtiter plate, cell culture flask, or a microarray, such as a gene chip, or the like, which contains the nucleic acid molecule, or polypeptide encoding the nGPCR or fragment thereof.
- [00038] As used herein, the phrase "homologous nucleotide sequence," or "homologous amino acid sequence," or variations thereof, refers to sequences characterized by a homology, at the nucleotide level or amino acid level, of at least the specified percentage. Homologous nucleotide sequences include those sequences coding for isoforms of proteins. Such isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. Homologous nucleotide sequences include nucleotide sequences encoding for a protein of a species other than humans, including, but not limited to, mammals. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding other known GPCRs. Homologous amino acid sequences include those amino acid sequences which contain conservative amino acid substitutions and which polypeptides have the same binding and/or activity. A homologous amino acid sequence does not, however, include the amino acid sequence encoding other known GPCRs. Percent homology can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison WI), using the default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482-489, which is incorporated herein by reference in its entirety).
- [00039] As used herein, the term "isolated" nucleic acid molecule refers to a nucleic acid molecule (DNA or RNA) that has been removed from its native environment. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules.

- [00040] As used herein, the terms "modulates" or "modifies" means an increase or decrease in the amount, quality, or effect of a particular activity or protein.
- [00041] As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues which has a sufficient number of bases to be used in a polymerase chain reaction (PCR). This short sequence is based on (or designed from) a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 50 nucleotides, preferably about 15 to 30 nucleotides. They are chemically synthesized and may be used as probes.
- [00042] As used herein, the term "probe" refers to nucleic acid sequences of variable length, preferably between at least about 10 and as many as about 6,000 nucleotides, depending on use. They are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligomers. They may be single or double-stranded and carefully designed to have specificity in PCR, hybridization membrane-based, or ELISA-like technologies.
- [00043] The term "preventing" refers to decreasing the probability that an organism contracts or develops an abnormal condition.
- [00044] The term "treating" refers to having a therapeutic effect and at least partially alleviating or abrogating an abnormal condition in the organism.
- [00045] The term "therapeutic effect" refers to the inhibition or activation factors causing or contributing to the abnormal condition. A therapeutic effect relieves to some extent one or more of the symptoms of the abnormal condition. In reference to the treatment of abnormal conditions, a therapeutic effect can refer to one or more of the following: (a) an increase in the proliferation, growth, and/or differentiation of cells; (b) inhibition (*i.e.*, slowing or stopping) of cell death; (c) inhibition of degeneration; (d) relieving to some extent one or more of the symptoms associated with the abnormal condition; and (e) enhancing the function of the affected population of cells. Compounds demonstrating efficacy against abnormal conditions can be identified as described herein.
- [00046] The term "abnormal condition" refers to a function in the cells or tissues of an organism that deviates from their normal functions in that organism. An abnormal condition can relate to cell proliferation, cell differentiation, cell signaling, or cell survival. An abnormal condition may also include obesity, diabetic complications such as retinal degeneration, and irregularities in glucose uptake and metabolism, and fatty acid uptake and metabolism.

- [00047] Abnormal cell proliferative conditions include cancers such as fibrotic and mesangial disorders, abnormal angiogenesis and vasculogenesis, wound healing, psoriasis, diabetes mellitus, and inflammation.
- [00048] Abnormal differentiation conditions include, but are not limited to, neurodegenerative disorders, slow wound healing rates, and slow tissue grafting healing rates. Abnormal cell signaling conditions include, but are not limited to, psychiatric disorders involving excess neurotransmitter activity.
- [00049] Abnormal cell survival conditions may also relate to conditions in which programmed cell death (apoptosis) pathways are activated or abrogated. A number of protein kinases are associated with the apoptosis pathways. Aberrations in the function of any one of the protein kinases could lead to cell immortality or premature cell death.
- [00050] The term "administering" relates to a method of incorporating a compound into cells or tissues of an organism. The abnormal condition can be prevented or treated when the cells or tissues of the organism exist within the organism or outside of the organism. Cells existing outside the organism can be maintained or grown in cell culture dishes. For cells harbored within the organism, many techniques exist in the art to administer compounds, including (but not limited to) oral, parenteral, dermal, injection, and aerosol applications. For cells outside of the organism, multiple techniques exist in the art to administer the compounds, including (but not limited to) cell microinjection techniques, transformation techniques and carrier techniques.
- [00051] The abnormal condition can also be prevented or treated by administering a compound to a group of cells having an aberration in a signal transduction pathway to an organism. The effect of administering a compound on organism function can then be monitored. The organism is preferably a mouse, rat, rabbit, guinea pig or goat, more preferably a monkey or ape, and most preferably a human.
- [00052] By "amplification" it is meant increased numbers of DNA or RNA in a cell compared with normal cells. "Amplification" as it refers to RNA can be the detectable presence of RNA in cells, since in some normal cells there is no basal expression of RNA. In other normal cells, a basal level of expression exists, therefore in these cases amplification is the detection of at least 1 to 2-fold, and preferably more, compared to the basal level.
- [00053] As used herein, the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which a probe, primer, or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m

is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present in excess, at T_m , 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (e.g. 10 to 50 nucleotides) and at least about 60°C for longer probes, primers or oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

[00054] The amino acid sequences are presented in the amino to carboxy direction, from left to right. The amino and carboxy groups are not presented in the sequence. The nucleotide sequences are presented by single strand only, in the 5' to 3' direction, from left to right. Nucleotides and amino acids are represented in the manner recommended by the IUPAC-IUB Biochemical Nomenclature Commission or (for amino acids) by three letters code.

Polynucleotides

[00055] The present invention provides purified and isolated polynucleotides (e.g., DNA sequences and RNA transcripts, both sense and complementary antisense strands, both single- and double-stranded, including splice variants thereof) that encode unknown G protein-coupled receptors heretofore termed novel GPCRs, or nGPCRs. These genes are described herein and designated herein collectively as nGPCR-x (where x is 86-93, 2588, 2589, 2591, 2592, 2593, 2594, 2595, 2596, 2598, 2600, 2601, 2602, 2603, 2604, 2606, 2607, 2608, 2609, 2610, 2611, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2621, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2639, 2640, 2641, 2642, 2643, 2644, and 2645). Table 1 below identifies the novel gene sequence nGPCR-x designation, the SEQ ID NO: of the gene sequence, the SEQ ID NO: of the polypeptide encoded thereby, and the U.S. Provisional Application in which the gene sequence has been disclosed.

Table 1

nGPCR	Nucleotide Sequence (SEQ ID NO:)	Amino acid Sequence (SEQ ID NO:)	Originally filed in:	nGPCR	Nucleotide Sequence (SEQ ID NO:)	Amino acid Sequence (SEQ ID NO:)	Originally filed in:
86	1	59	A	2613	30	88	D
87	2	60	A	2614	31	89	D
88	3	61	A	2615	32	90	D
89	4	62	A	2616	33	91	D
90	5	63	A	2617	34	92	D
91	6	64	A	2618	35	93	D
92	7	65	A	2619	36	94	D
93	8	66	A	2621	37	95	D
93	9	67	G	2624	38	96	D
2588	10	68	B	2625	39	97	D
2589	11	69	B	2626	40	98	E

2591	12	70	B	2627	41	99	E
2592	13	71	B	2628	42	100	E
2593	14	72	B	2629	43	101	E
2594	15	73	B	2630	44	102	E
2595	16	74	B	2631	45	103	E
2596	17	75	B	2632	46	104	E
2598	18	76	B	2633	47	105	E
2600	19	77	B	2634	48	106	E
2601	20	78	C	2635	49	107	E
2602	21	79	C	2636	50	108	F
2603	22	80	C	2637	51	109	F
2604	23	81	C	2639	52	110	F
2606	24	82	C	2640	53	111	F
2607	25	83	C	2641	54	112	F
2608	26	84	C	2642	55	113	F
2609	27	85	C	2643	56	114	F
2610	28	86	C	2644	57	115	F
2611	29	87	C	2645	58	116	F

Legend

A= Ser. No. 60/195,150

C= Ser. No. 60/195,151

E= Ser. No. 60/195,093

G= Ser. No. 60/230,149

B= Ser. No. 60/195,099

D= Ser. No. 60/195,148

F= Ser. No. 60/195,098

[00056] When a specific nGPCR is identified (for example nGPCR-2611), it is understood that only that specific nGPCR is being referred to.

[00057] It is well known that GPCRs are expressed in many different tissues, including the brain. Accordingly, the nGPCR-x of the present invention may be useful, *inter alia*, for treating and/or diagnosing mental disorders. Following the techniques described in Example 5, below, those skilled in the art could readily ascertain if nGPCR-x is expressed in a particular tissue or region.

[00058] The invention provides purified and isolated polynucleotides (*e.g.*, cDNA, genomic DNA, synthetic DNA, RNA, or combinations thereof, whether single- or double-stranded) that comprise a nucleotide sequence encoding the amino acid sequence of the polypeptides of the invention. Such polynucleotides are useful for recombinantly expressing the receptor and also for detecting expression of the receptor in cells (*e.g.*, using Northern hybridization and *in situ* hybridization assays). Such polynucleotides also are useful in the design of antisense and other molecules for the suppression of the expression of nGPCR-x in a cultured cell, a tissue, or an animal; for therapeutic purposes; or to provide a model for diseases or conditions characterized by aberrant nGPCR-x expression. Specifically excluded from the definition of polynucleotides of the invention are entire isolated, non-recombinant native chromosomes of host cells. A preferred polynucleotide has a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, which correspond to naturally occurring nGPCR-x sequences. It will be appreciated that numerous other polynucleotide sequences exist that also encode nGPCR-x

having the sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, due to the well-known degeneracy of the universal genetic code.

[00059] The invention also provides a purified and isolated polynucleotide comprising a nucleotide sequence that encodes a mammalian polypeptide, wherein the polynucleotide hybridizes to a polynucleotide having the sequence set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or the non-coding strand complementary thereto, under the following hybridization conditions:

(a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate; and

(b) washing 2 times for 30 minutes each at 60°C in a wash solution comprising 0.1% SSC, 1% SDS. Polynucleotides that encode a human allelic variant are highly preferred.

[00060] The present invention relates to molecules which comprise the gene sequences that encode the nGPCRs; constructs and recombinant host cells incorporating the gene sequences; the novel GPCR polypeptides encoded by the gene sequences; antibodies to the polypeptides and homologs; kits employing the polynucleotides and polypeptides, and methods of making and using all of the foregoing. In addition, the present invention relates to homologs of the gene sequences and of the polypeptides and methods of making and using the same.

[00061] Genomic DNA of the invention comprises the protein-coding region for a polypeptide of the invention and is also intended to include allelic variants thereof. It is widely understood that, for many genes, genomic DNA is transcribed into RNA transcripts that undergo one or more splicing events wherein intron (*i.e.*, non-coding regions) of the transcripts are removed, or "spliced out." RNA transcripts that can be spliced by alternative mechanisms, and therefore be subject to removal of different RNA sequences but still encode a nGPCR-x polypeptide, are referred to in the art as splice variants which are embraced by the invention. Splice variants comprehended by the invention therefore are encoded by the same original genomic DNA sequences but arise from distinct mRNA transcripts. Allelic variants are modified forms of a wild-type gene sequence, the modification resulting from recombination during chromosomal segregation or exposure to conditions which give rise to genetic mutation. Allelic variants, like wild type genes, are naturally occurring sequences (as opposed to non-naturally occurring variants that arise from *in vitro* manipulation).

[00062] The invention also comprehends cDNA that is obtained through reverse transcription of an RNA polynucleotide encoding nGPCR-x (conventionally followed by second strand synthesis of a complementary strand to provide a double-stranded DNA).

[00063] Preferred DNA sequences encoding human nGPCR-x polypeptides are selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58. A preferred DNA of the invention

comprises a double stranded molecule along with the complementary molecule (the "non-coding strand" or "complement") having a sequence unambiguously deducible from the coding strand according to Watson-Crick base-pairing rules for DNA. Also preferred are other polynucleotides encoding the nGPCR-x polypeptide selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, which differ in sequence from the polynucleotides selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, by virtue of the well-known degeneracy of the universal nuclear genetic code.

[00064] The invention further embraces other species, preferably mammalian, homologs of the human nGPCR-x DNA. Species homologs, sometimes referred to as "orthologs," in general, share at least 35%, at least 40%, at least 45%, at least 50%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% homology with human DNA of the invention. Generally, percent sequence "homology" with respect to polynucleotides of the invention may be calculated as the percentage of nucleotide bases in the candidate sequence that are identical to nucleotides in the nGPCR-x sequence set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity.

[00065] Polynucleotides of the invention permit identification and isolation of polynucleotides encoding related nGPCR-x polypeptides, such as human allelic variants and species homologs, by well-known techniques including Southern and/or Northern hybridization, and polymerase chain reaction (PCR). Examples of related polynucleotides include human and non-human genomic sequences, including allelic variants, as well as polynucleotides encoding polypeptides homologous to nGPCR-x and structurally related polypeptides sharing one or more biological, immunological, and/or physical properties of nGPCR-x. Non-human species genes encoding proteins homologous to nGPCR-x can also be identified by Southern and/or PCR analysis and are useful in animal models for nGPCR-x disorders. Knowledge of the sequence of a human nGPCR-x DNA also makes possible through use of Southern hybridization or polymerase chain reaction (PCR) the identification of genomic DNA sequences encoding nGPCR-x expression control regulatory sequences such as promoters, operators, enhancers, repressors, and the like. Polynucleotides of the invention are also useful in hybridization assays to detect the capacity of cells to express nGPCR-x. Polynucleotides of the invention may also provide a basis for diagnostic methods useful for identifying a genetic alteration(s) in a nGPCR-x locus that underlies a disease state or states, which information is useful both for diagnosis and for selection of therapeutic strategies.

[00066] According to the present invention, the nGPCR-x nucleotide sequences disclosed herein may be used to identify homologs of the nGPCR-x, in other animals, including but not limited to humans and other mammals, and invertebrates. Any of the nucleotide sequences disclosed herein, or any portion thereof, can be used, for example, as probes to screen databases or nucleic acid libraries, such as, for example, genomic or cDNA libraries, to identify homologs, using screening procedures well known to those skilled in the art. Accordingly, homologs having at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, more preferably at least 90%, more preferably at least 95%, and most preferably at least 100% homology with nGPCR-x sequences can be identified.

[00067] The disclosure herein of full-length polynucleotides encoding nGPCR-x polypeptides makes readily available to the worker of ordinary skill in the art every possible fragment of the full-length polynucleotide.

[00068] One preferred embodiment of the present invention provides an isolated nucleic acid molecule comprising a sequence homologous sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, and fragments thereof. Another preferred embodiment provides an isolated nucleic acid molecule comprising a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, and fragments thereof.

[00069] As used in the present invention, fragments of nGPCR-x-encoding polynucleotides comprise at least 10, and preferably at least 12, 14, 16, 18, 20, 25, 50, or 75 consecutive nucleotides of a polynucleotide encoding nGPCR-x. Preferably, fragment polynucleotides of the invention comprise sequences unique to the nGPCR-x-encoding polynucleotide sequence, and therefore hybridize under highly stringent or moderately stringent conditions only (*i.e.*, "specifically") to polynucleotides encoding nGPCR-x (or fragments thereof). Polynucleotide fragments of genomic sequences of the invention comprise not only sequences unique to the coding region, but also include fragments of the full-length sequence derived from introns, regulatory regions, and/or other non-translated sequences. Sequences unique to polynucleotides of the invention are recognizable through sequence comparison to other known polynucleotides, and can be identified through use of alignment programs routinely utilized in the art, *e.g.*, those made available in public sequence databases. Such sequences also are recognizable from Southern hybridization analyses to determine the number of fragments of genomic DNA to which a polynucleotide will hybridize. Polynucleotides of the invention can be labeled in a manner that permits their detection, including radioactive, fluorescent, and enzymatic labeling.

[00070] Fragment polynucleotides are particularly useful as probes for detection of full-length or fragments of nGPCR-x polynucleotides. One or more polynucleotides can be included in kits

that are used to detect the presence of a polynucleotide encoding nGPCR-x, or used to detect variations in a polynucleotide sequence encoding nGPCR-x.

[00071] The invention also embraces DNAs encoding nGPCR-x polypeptides that hybridize under moderately stringent or high stringency conditions to the non-coding strand, or complement, of the polynucleotides set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58.

[00072] Exemplary highly stringent hybridization conditions are as follows: hybridization at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% Dextran sulfate, and washing twice for 30 minutes at 60°C in a wash solution comprising 0.1X SSC and 1% SDS. It is understood in the art that conditions of equivalent stringency can be achieved through variation of temperature and buffer, or salt concentration as described Ausubel *et al.* (Eds.), Protocols in Molecular Biology, John Wiley & Sons (1994), pp. 6.0.3 to 6.4.10. Modifications in hybridization conditions can be empirically determined or precisely calculated based on the length and the percentage of guanosine/cytosine (GC) base pairing of the probe. The hybridization conditions can be calculated as described in Sambrook, *et al.*, (Eds.), Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press: Cold Spring Harbor, New York (1989), pp. 9.47 to 9.51.

[00073] With the knowledge of the nucleotide sequence information disclosed in the present invention, one skilled in the art can identify and obtain nucleotide sequences which encode nGPCR-x from different sources (*i.e.*, different tissues or different organisms) through a variety of means well known to the skilled artisan and as disclosed by, for example, Sambrook *et al.*, "Molecular cloning: a laboratory manual", Second Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), which is incorporated herein by reference in its entirety.

[00074] For example, DNA that encodes nGPCR-x may be obtained by screening of mRNA, cDNA, or genomic DNA with oligonucleotide probes generated from the nGPCR-x gene sequence information provided herein. Probes may be labeled with a detectable group, such as a fluorescent group, a radioactive atom or a chemiluminescent group in accordance with procedures known to the skilled artisan and used in conventional hybridization assays, as described by, for example, Sambrook *et al.*

[00075] A nucleic acid molecule comprising any of the nGPCR-x nucleotide sequences described above can alternatively be synthesized by use of the polymerase chain reaction (PCR) procedure, with the PCR oligonucleotide primers produced from the nucleotide sequences provided herein. See U.S. Patent Numbers 4,683,195 to Mullis *et al.* and 4,683,202 to Mullis. The PCR reaction provides a method for selectively increasing the concentration of a particular nucleic acid sequence even when that sequence has not been previously purified and is present.

only in a single copy in a particular sample. The method can be used to amplify either single or double-stranded DNA. The essence of the method involves the use of two oligonucleotide probes to serve as primers for the template-dependent, polymerase mediated replication of a desired nucleic acid molecule.

[00076] A wide variety of alternative cloning and *in vitro* amplification methodologies are well known to those skilled in the art. Examples of these techniques are found in, for example, Berger *et al.*, *Guide to Molecular Cloning Techniques*, Methods in Enzymology 152, Academic Press, Inc., San Diego, CA (Berger), which is incorporated herein by reference in its entirety.

[00077] Automated sequencing methods can be used to obtain or verify the nucleotide sequence of nGPCR-x. The nGPCR-x nucleotide sequences of the present invention are believed to be 100% accurate. However, as is known in the art, nucleotide sequence obtained by automated methods may contain some errors. Nucleotide sequences determined by automation are typically at least about 90%, more typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of a given nucleic acid molecule. The actual sequence may be more precisely determined using manual sequencing methods, which are well known in the art. An error in a sequence which results in an insertion or deletion of one or more nucleotides may result in a frame shift in translation such that the predicted amino acid sequence will differ from that which would be predicted from the actual nucleotide sequence of the nucleic acid molecule, starting at the point of the mutation.

[00078] The nucleic acid molecules of the present invention, and fragments derived therefrom, are useful for screening for restriction fragment length polymorphism (RFLP) associated with certain disorders, as well as for genetic mapping.

[00079] The polynucleotide sequence information provided by the invention makes possible large-scale expression of the encoded polypeptide by techniques well known and routinely practiced in the art.

Vectors

[00080] Another aspect of the present invention is directed to vectors, or recombinant expression vectors, comprising any of the nucleic acid molecules described above. Vectors are used herein either to amplify DNA or RNA encoding nGPCR-x and/or to express DNA which encodes nGPCR-x. Preferred vectors include, but are not limited to, plasmids, phages, cosmids, episomes, viral particles or viruses, and integratable DNA fragments (*i.e.*, fragments integratable into the host genome by homologous recombination). Preferred viral particles include, but are not limited to, adenoviruses, baculoviruses, parvoviruses, herpesviruses, poxviruses, adeno-associated viruses, Semliki Forest viruses, vaccinia viruses, and retroviruses. Preferred expression vectors include, but are not limited to, pcDNA3 (Invitrogen) and pSVL (Pharmacia

Biotech). Other expression vectors include, but are not limited to, pSPORT™ vectors, pGEM™ vectors (Promega), pPROEXvectors™ (LTI, Bethesda, MD), Bluescript™ vectors (Stratagene), pQE™ vectors (Qiagen), pSE420™ (Invitrogen), and pYES2™(Invitrogen).

[00081] Expression constructs preferably comprise GPCR-x-encoding polynucleotides operatively linked to an endogenous or exogenous expression control DNA sequence and a transcription terminator. Expression control DNA sequences include promoters, enhancers, operators, and regulatory element binding sites generally, and are typically selected based on the expression systems in which the expression construct is to be utilized. Preferred promoter and enhancer sequences are generally selected for the ability to increase gene expression, while operator sequences are generally selected for the ability to regulate gene expression. Expression constructs of the invention may also include sequences encoding one or more selectable markers that permit identification of host cells bearing the construct. Expression constructs may also include sequences that facilitate, and preferably promote, homologous recombination in a host cell. Preferred constructs of the invention also include sequences necessary for replication in a host cell.

[00082] Expression constructs are preferably utilized for production of an encoded protein, but may also be utilized simply to amplify a nGPCR-x-encoding polynucleotide sequence. In preferred embodiments, the vector is an expression vector wherein the polynucleotide of the invention is operatively linked to a polynucleotide comprising an expression control sequence. Autonomously replicating recombinant expression constructs such as plasmid and viral DNA vectors incorporating polynucleotides of the invention are also provided. Preferred expression vectors are replicable DNA constructs in which a DNA sequence encoding nGPCR-x is operably linked or connected to suitable control sequences capable of effecting the expression of the nGPCR-x in a suitable host. DNA regions are operably linked or connected when they are functionally related to each other. For example, a promoter is operably linked or connected to a coding sequence if it controls the transcription of the sequence. Amplification vectors do not require expression control domains, but rather need only the ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants. The need for control sequences in the expression vector will vary depending upon the host selected and the transformation method chosen. Generally, control sequences include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding and sequences which control the termination of transcription and translation.

[00083] Preferred vectors preferably contain a promoter that is recognized by the host organism. The promoter sequences of the present invention may be prokaryotic, eukaryotic or viral.

Examples of suitable prokaryotic sequences include the P_R and P_L promoters of bacteriophage lambda (The bacteriophage Lambda, Hershey, A. D., Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY (1973), which is incorporated herein by reference in its entirety; Lambda II, Hendrix, R. W., Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY (1980), which is incorporated herein by reference in its entirety); the *trp*, *recA*, heat shock, and *lacZ* promoters of *E. coli* and the SV40 early promoter (Benoist *et al. Nature*, 1981, 290, 304-310, which is incorporated herein by reference in its entirety). Additional promoters include, but are not limited to, mouse mammary tumor virus, long terminal repeat of human immunodeficiency virus, maloney virus, cytomegalovirus immediate early promoter, Epstein Barr virus, Rous sarcoma virus, human actin, human myosin, human hemoglobin, human muscle creatine, and human metallothionein.

- [00084] Additional regulatory sequences can also be included in preferred vectors. Preferred examples of suitable regulatory sequences are represented by the Shine-Dalgarno of the replicase gene of the phage MS-2 and of the gene *cII* of bacteriophage lambda. The Shine-Dalgarno sequence may be directly followed by DNA encoding nGPCR-x and result in the expression of the mature nGPCR-x protein.
- [00085] Moreover, suitable expression vectors can include an appropriate marker that allows the screening of the transformed host cells. The transformation of the selected host is carried out using any one of the various techniques well known to the expert in the art and described in Sambrook *et al.*, *supra*.
- [00086] An origin of replication can also be provided either by construction of the vector to include an exogenous origin or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter may be sufficient. Alternatively, rather than using vectors which contain viral origins of replication, one skilled in the art can transform mammalian cells by the method of co-transformation with a selectable marker and nGPCR-x DNA. An example of a suitable marker is dihydrofolate reductase (DHFR) or thymidine kinase (*see*, U.S. Patent No. 4,399,216).
- [00087] Nucleotide sequences encoding GPCR-x may be recombined with vector DNA in accordance with conventional techniques, including blunt-ended or staggered-ended termini for ligation, restriction enzyme digestion to provide appropriate termini, filling in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulation are disclosed by Sambrook *et al.*, *supra* and are well known in the art. Methods for construction of mammalian expression vectors are disclosed in, for example, Okayama *et al.*, *Mol. Cell. Biol.*, 1983, 3, 280, Cosman *et al.*, *Mol.*

Immunol., 1986, 23, 935, Cosman *et al.*, *Nature*, 1984, 312, 768, EP-A-0367566, and WO 91/18982, each of which is incorporated herein by reference in its entirety.

Host cells

- [00088] According to another aspect of the invention, host cells are provided, including prokaryotic and eukaryotic cells, comprising a polynucleotide of the invention (or vector of the invention) in a manner that permits expression of the encoded nGPCR-x polypeptide. Polynucleotides of the invention may be introduced into the host cell as part of a circular plasmid, or as linear DNA comprising an isolated protein coding region or a viral vector. Methods for introducing DNA into the host cell that are well known and routinely practiced in the art include transformation, transfection, electroporation, nuclear injection, or fusion with carriers such as liposomes, micelles, ghost cells, and protoplasts. Expression systems of the invention include bacterial, yeast, fungal, plant, insect, invertebrate, vertebrate, and mammalian cells systems.
- [00089] The invention provides host cells that are transformed or transfected (stably or transiently) with polynucleotides of the invention or vectors of the invention. As stated above, such host cells are useful for amplifying the polynucleotides and also for expressing the nGPCR-x polypeptide or fragment thereof encoded by the polynucleotide.
- [00090] In still another related embodiment, the invention provides a method for producing a nGPCR-x polypeptide (or fragment thereof) comprising the steps of growing a host cell of the invention in a nutrient medium and isolating the polypeptide or variant thereof from the cell or the medium. Because nGPCR-x is a seven transmembrane receptor, it will be appreciated that, for some applications, such as certain activity assays, the preferable isolation may involve isolation of cell membranes containing the polypeptide embedded therein, whereas for other applications a more complete isolation may be preferable.
- [00091] According to some aspects of the present invention, transformed host cells having an expression vector comprising any of the nucleic acid molecules described above are provided. Expression of the nucleotide sequence occurs when the expression vector is introduced into an appropriate host cell. Suitable host cells for expression of the polypeptides of the invention include, but are not limited to, prokaryotes, yeast, and eukaryotes. If a prokaryotic expression vector is employed, then the appropriate host cell would be any prokaryotic cell capable of expressing the cloned sequences. Suitable prokaryotic cells include, but are not limited to, bacteria of the genera *Escherichia*, *Bacillus*, *Salmonella*, *Pseudomonas*, *Streptomyces*, and *Staphylococcus*.
- [00092] If an eukaryotic expression vector is employed, then the appropriate host cell would be any eukaryotic cell capable of expressing the cloned sequence. Preferably, eukaryotic cells are

cells of higher eukaryotes. Suitable eukaryotic cells include, but are not limited to, non-human mammalian tissue culture cells and human tissue culture cells. Preferred host cells include, but are not limited to, insect cells, HeLa cells, Chinese hamster ovary cells (CHO cells), African green monkey kidney cells (COS cells), human HEK-293 cells, and murine 3T3 fibroblasts. Propagation of such cells in cell culture has become a routine procedure (*see*, Tissue Culture, Academic Press, Kruse and Patterson, eds. (1973), which is incorporated herein by reference in its entirety).

[00093] In addition, a yeast host may be employed as a host cell. Preferred yeast cells include, but are not limited to, the genera *Saccharomyces*, *Pichia*, and *Kluveromyces*. Preferred yeast hosts are *S. cerevisiae* and *P. pastoris*. Preferred yeast vectors can contain an origin of replication sequence from a 2T yeast plasmid, an autonomously replication sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Shuttle vectors for replication in both yeast and *E. coli* are also included herein.

[00094] Alternatively, insect cells may be used as host cells. In a preferred embodiment, the polypeptides of the invention are expressed using a baculovirus expression system (*see*, Luckow *et al.*, *Bio/Technology*, 1988, 6, 47, Baculovirus Expression Vectors: A Laboratory Manual, O'Rielly *et al.* (Eds.), W.H. Freeman and Company, New York, 1992, and U.S. Patent No. 4,879,236, each of which is incorporated herein by reference in its entirety). In addition, the MAXBAC™ complete baculovirus expression system (Invitrogen) can, for example, be used for production in insect cells.

[00095] Host cells of the invention are a valuable source of immunogen for development of antibodies specifically immunoreactive with nGPCR-x. Host cells of the invention are also useful in methods for the large-scale production of nGPCR-x polypeptides wherein the cells are grown in a suitable culture medium and the desired polypeptide products are isolated from the cells, or from the medium in which the cells are grown, by purification methods known in the art, *e.g.*, conventional chromatographic methods including immunoaffinity chromatography, receptor affinity chromatography, hydrophobic interaction chromatography, lectin affinity chromatography, size exclusion filtration, cation or anion exchange chromatography, high pressure liquid chromatography (HPLC), reverse phase HPLC, and the like. Still other methods of purification include those methods wherein the desired protein is expressed and purified as a fusion protein having a specific tag, label, or chelating moiety that is recognized by a specific binding partner or agent. The purified protein can be cleaved to yield the desired protein, or can be left as an intact fusion protein. Cleavage of the fusion component may produce a form of the desired protein having additional amino acid residues as a result of the cleavage process.

[00096] Knowledge of nGPCR-x DNA sequences allows for modification of cells to permit, or increase, expression of endogenous nGPCR-x. Cells can be modified (e.g., by homologous recombination) to provide increased expression by replacing, in whole or in part, the naturally occurring nGPCR-x promoter with all or part of a heterologous promoter so that the cells express nGPCR-x at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to endogenous nGPCR-x encoding sequences. (See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955.) It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamoyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the nGPCR-x coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the nGPCR-x coding sequences in the cells.

Knock-outs

[00097] The DNA sequence information provided by the present invention also makes possible the development (e.g., by homologous recombination or "knock-out" strategies; see Capecchi, *Science* 244:1288-1292 (1989), which is incorporated herein by reference) of animals that fail to express functional nGPCR-x or that express a variant of nGPCR-x. Such animals (especially small laboratory animals such as rats, rabbits, and mice) are useful as models for studying their *vivo* activities of nGPCR-x and modulators of nGPCR-x.

Antisense

[00098] Also made available by the invention are anti-sense polynucleotides that recognize and hybridize to polynucleotides encoding nGPCR-x. Full-length and fragment anti-sense polynucleotides are provided. Fragment antisense molecules of the invention include (i) those that specifically recognize and hybridize to nGPCR-x RNA (as determined by sequence comparison of DNA encoding nGPCR-x to DNA encoding other known molecules). Identification of sequences unique to nGPCR-x encoding polynucleotides can be deduced through use of any publicly available sequence database, and/or through use of commercially available sequence comparison programs. After identification of the desired sequences, isolation through restriction digestion or amplification using any of the various polymerase chain reaction techniques well known in the art can be performed. Anti-sense polynucleotides are particularly relevant to regulating expression of nGPCR-x by those cells expressing nGPCR-x mRNA.

[00099] Antisense nucleic acids (preferably 10 to 30 base-pair oligonucleotides) capable of specifically binding to nGPCR-x expression control sequences or nGPCR-x RNA are introduced into cells (e.g., by a viral vector or colloidal dispersion system such as a liposome). The antisense nucleic acid binds to the nGPCR-x target nucleotide sequence in the cell and prevents transcription and/or translation of the target sequence. Phosphorothioate and methylphosphonate antisense oligonucleotides are specifically contemplated for therapeutic use by the invention. The antisense oligonucleotides may be further modified by adding poly-L-lysine, transferrin polylysine, or cholesterol moieties at their 5' end. Suppression of nGPCR-x expression at either the transcriptional or translational level is useful to generate cellular or animal models for diseases/conditions characterized by aberrant nGPCR-x expression.

[000100] Antisense oligonucleotides, or fragments of sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or sequences complementary or homologous thereto, derived from the nucleotide sequences of the present invention encoding nGPCR-x are useful as diagnostic tools for probing gene expression in various tissues. For example, tissue can be probed *in situ* with oligonucleotide probes carrying detectable groups by conventional autoradiography techniques to investigate native expression of this enzyme or pathological conditions relating thereto. Antisense oligonucleotides are preferably directed to regulatory regions of sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or mRNA corresponding thereto, including, but not limited to, the initiation codon, TATA box, enhancer sequences, and the like.

Transcription factors

[000101] The nGPCR-x sequences taught in the present invention facilitate the design of novel transcription factors for modulating nGPCR-x expression in native cells and animals, and cells transformed or transfected with nGPCR-x polynucleotides. For example, the Cys₂-His₂ zinc finger proteins, which bind DNA via their zinc finger domains, have been shown to be amenable to structural changes that lead to the recognition of different target sequences. These artificial zinc finger proteins recognize specific target sites with high affinity and low dissociation constants, and are able to act as gene switches to modulate gene expression. Knowledge of the particular nGPCR-x target sequence of the present invention facilitates the engineering of zinc finger proteins specific for the target sequence using known methods such as a combination of structure-based modeling and screening of phage display libraries (Segal *et al.*, Proc. Natl. Acad. Sci. (USA) 96:2758-2763 (1999); Liu *et al.*, Proc. Natl. Acad. Sci. (USA) 94:5525-5530 (1997); Greisman *et al.*, Science 275:657-661 (1997); Choo *et al.*, J. Mol. Biol. 273:525-532 (1997)). Each zinc finger domain usually recognizes three or more base pairs. Since a recognition sequence of 18 base pairs is generally sufficient in length to render it unique in any

known genome, a zinc finger protein consisting of 6 tandem repeats of zinc fingers would be expected to ensure specificity for a particular sequence (Segal *et al.*) The artificial zinc finger repeats, designed based on nGPCR-x sequences, are fused to activation or repression domains to promote or suppress nGPCR-x expression (Liu *et al.*) Alternatively, the zinc finger domains can be fused to the TATA box-binding factor (TBP) with varying lengths of linker region between the zinc finger peptide and the TBP to create either transcriptional activators or repressors (Kim *et al.*, Proc. Natl. Acad. Sci. (USA) 94:3616-3620 (1997). Such proteins and polynucleotides that encode them, have utility for modulating nGPCR-x expression *in vivo* in both native cells, animals and humans; and/or cells transfected with nGPCR-x-encoding sequences. The novel transcription factor can be delivered to the target cells by transfecting constructs that express the transcription factor (gene therapy), or by introducing the protein. Engineered zinc finger proteins can also be designed to bind RNA sequences for use in therapeutics as alternatives to antisense or catalytic RNA methods (McColl *et al.*, Proc. Natl. Acad. Sci. (USA) 96:9521-9526 (1997); Wu *et al.*, Proc. Natl. Acad. Sci. (USA) 92:344-348 (1995)). The present invention contemplates methods of designing such transcription factors based on the gene sequence of the invention, as well as customized zinc finger proteins, that are useful to modulate nGPCR-x expression in cells (native or transformed) whose genetic complement includes these sequences.

Polypeptides

- [000102] The invention also provides purified and isolated mammalian nGPCR-x polypeptides encoded by a polynucleotide of the invention. Presently preferred is a human nGPCR-x polypeptide comprising the amino acid sequence set out in sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, or fragments thereof comprising an epitope specific to the polypeptide. By "epitope specific to" is meant a portion of the nGPCR receptor that is recognizable by an antibody that is specific for the nGPCR, as defined in detail below.
- [000103] Although the sequences provided are particular human sequences, the invention is intended to include within its scope other human allelic variants; non-human mammalian forms of nGPCR-x, and other vertebrate forms of nGPCR-x.
- [000104] It will be appreciated that extracellular epitopes are particularly useful for generating and screening for antibodies and other binding compounds that bind to receptors such as nGPCR-x. Thus, in another preferred embodiment, the invention provides a purified and isolated polypeptide comprising at least one extracellular domain (*e.g.*, the N-terminal extracellular domain or one of the three extracellular loops) of nGPCR-x. Purified and isolated polypeptides comprising the N-terminal extracellular domain of nGPCR-x are highly preferred. Also preferred is a purified and isolated polypeptide comprising a nGPCR-x fragment selected from the group consisting of the N-terminal extracellular domain of nGPCR-x, transmembrane

domains of nGPCR-x, an extracellular loop connecting transmembrane domains of nGPCR-x, an intracellular loop connecting transmembrane domains of nGPCR-x, the C-terminal cytoplasmic region of nGPCR-x, and fusions thereof. Such fragments may be continuous portions of the native receptor. However, it will also be appreciated that knowledge of the nGPCR-x gene and protein sequences as provided herein permits recombining of various domains that are not contiguous in the native protein. Using a FORTRAN computer program called "tmtest.all" [Parodi *et al.*, Comput. Appl. Biosci. 5:527-535 (1994)], nGPCR-x was shown to contain transmembrane-spanning domains.

[000105] The invention also embraces polypeptides that have at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55% or at least 50% identity and/or homology to the preferred polypeptide of the invention. Percent amino acid sequence "identity" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in the nGPCR-x sequence after aligning both sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Percent sequence "homology" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in the nGPCR-x sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and also considering any conservative substitutions as part of the sequence identity.

[000106] In one aspect, percent homology is calculated as the percentage of amino acid residues in the smaller of two sequences which align with identical amino acid residue in the sequence being compared, when four gaps in a length of 100 amino acids may be introduced to maximize alignment (Dayhoff, in Atlas of Protein Sequence and Structure, Vol. 5, p. 124, National Biochemical Research Foundation, Washington, D.C. (1972), incorporated herein by reference).

[000107] Polypeptides of the invention may be isolated from natural cell sources or may be chemically synthesized, but are preferably produced by recombinant procedures involving host cells of the invention. Use of mammalian host cells is expected to provide for such post-translational modifications (*e.g.*, glycosylation, truncation, lipidation, and phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention. Glycosylated and non-glycosylated forms of nGPCR-x polypeptides are embraced by the invention.

[000108] The invention also embraces variant (or analog) nGPCR-x polypeptides. In one example, insertion variants are provided wherein one or more amino acid residues supplement a

nGPCR-x amino acid sequence. Insertions may be located at either or both termini of the protein, or may be positioned within internal regions of the nGPCR-x amino acid sequence. Insertional variants with additional residues at either or both termini can include, for example, fusion proteins and proteins including amino acid tags or labels.

- [000109] Insertion variants include nGPCR-x polypeptides wherein one or more amino acid residues are added to a nGPCR-x acid sequence or to a biologically active fragment thereof.
- [000110] Variant products of the invention also include mature nGPCR-x products, *i.e.*, nGPCR-x products wherein leader or signal sequences are removed, with additional amino terminal residues. The additional amino terminal residues may be derived from another protein, or may include one or more residues that are not identifiable as being derived from specific proteins. nGPCR-x products with an additional methionine residue at position -1 (Met⁻¹-nGPCR-x) are contemplated, as are variants with additional methionine and lysine residues at positions -2 and -1 (Met⁻²-Lys⁻¹-nGPCR-x). Variants of nGPCR-x with additional Met, Met-Lys, Lys residues (or one or more basic residues in general) are particularly useful for enhanced recombinant protein production in bacterial host cells.
- [000111] The invention also embraces nGPCR-x variants having additional amino acid residues that result from use of specific expression systems. For example, use of commercially available vectors that express a desired polypeptide as part of a glutathione-S-transferase (GST) fusion product provides the desired polypeptide having an additional glycine residue at position -1 after cleavage of the GST component from the desired polypeptide. Variants that result from expression in other vector systems are also contemplated.
- [000112] Insertional variants also include fusion proteins wherein the amino terminus and/or the carboxy terminus of nGPCR-x is/are fused to another polypeptide.
- [000113] In another aspect, the invention provides deletion variants wherein one or more amino acid residues in a nGPCR-x polypeptide are removed. Deletions can be effected at one or both termini of the nGPCR-x polypeptide, or with removal of one or more non-terminal amino acid residues of nGPCR-x. Deletion variants, therefore, include all fragments of a nGPCR-x polypeptide.
- [000114] The invention also embraces polypeptide fragments of sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, wherein the fragments maintain biological (*e.g.*, ligand binding and/or intracellular signaling) immunological properties of a nGPCR-x polypeptide.
- [000115] In one preferred embodiment of the invention, an isolated nucleic acid molecule comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID

NO:116, and fragments thereof, wherein the nucleic acid molecule encoding at least a portion of nGPCR-x. In a more preferred embodiment, the isolated nucleic acid molecule comprises a sequence that encodes a polypeptide comprising sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, and fragments thereof.

[000116] As used in the present invention, polypeptide fragments comprise at least 5, 10, 15, 20, 25, 30, 35, or 40 consecutive amino acids of sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116. Preferred polypeptide fragments display antigenic properties unique to, or specific for, human nGPCR-x and its allelic and species homologs. Fragments of the invention having the desired biological and immunological properties can be prepared by any of the methods well known and routinely practiced in the art.

[000117] In still another aspect, the invention provides substitution variants of nGPCR-x polypeptides. Substitution variants include those polypeptides wherein one or more amino acid residues of a nGPCR-x polypeptide are removed and replaced with alternative residues. In one aspect, the substitutions are conservative in nature; however, the invention embraces substitutions that are also non-conservative. Conservative substitutions for this purpose may be defined as set out in Tables 2, 3, or 4 below.

[000118] Variant polypeptides include those wherein conservative substitutions have been introduced by modification of polynucleotides encoding polypeptides of the invention. Amino acids can be classified according to physical properties and contribution to secondary and tertiary protein structure. A conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in Table 2 (from WO 97/09433, page 10, published March 13, 1997 (PCT/GB96/02197, filed 9/6/96), immediately below.

Table 2
Conservative Substitutions I

<u>SIDE CHAIN CHARACTERISTIC</u>	<u>AMINO ACID</u>
Aliphatic	
Non-polar	G A P I L V
Polar - uncharged	C S T M N Q
Polar - charged	D E K R
Aromatic	H F W Y
Other	N Q D E

[000119] Alternatively, conservative amino acids can be grouped as described in Lehninger, [Biochemistry, Second Edition; Worth Publishers, Inc. NY, NY (1975), pp.71-77] as set out in Table 3, below.

Table 3
Conservative Substitutions II

<u>SIDE CHAIN CHARACTERISTIC</u>	<u>AMINO ACID</u>
Non-polar (hydrophobic)	
A. Aliphatic:	A L I V P
B. Aromatic:	F W
C. Sulfur-containing:	M
D. Borderline:	G
Uncharged-polar	
A. Hydroxyl:	S T Y
B. Amides:	N Q
C. Sulfhydryl:	C
D. Borderline:	G
Positively Charged (Basic):	K R H
Negatively Charged (Acidic):	D E

[000120] As still another alternative, exemplary conservative substitutions are set out in Table 4, below.

Table 4
Conservative Substitutions III

<u>Original Residue</u>	<u>Exemplary Substitution</u>
Ala (A)	Val, Leu, Ile
Arg (R)	Lys, Gln, Asn
Asn (N)	Gln, His, Lys, Arg
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (E)	Asp
His (H)	Asn, Gln, Lys, Arg
Ile (I)	Leu, Val, Met, Ala, Phe,
Leu (L)	Ile, Val, Met, Ala, Phe
Lys (K)	Arg, Gln, Asn
Met (M)	Leu, Phe, Ile
Phe (F)	Leu, Val, Ile, Ala
Pro (P)	Gly
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser
Val (V)	Ile, Leu, Met, Phe, Ala

[000121] It should be understood that the definition of polypeptides of the invention is intended to include polypeptides bearing modifications other than insertion, deletion, or substitution of amino acid residues. By way of example, the modifications may be covalent in nature, and include for example, chemical bonding with polymers, lipids, other organic, and inorganic moieties. Such derivatives may be prepared to increase circulating half-life of a polypeptide, or may be designed to improve the targeting capacity of the polypeptide for desired cells, tissues,

or organs. Similarly, the invention further embraces nGPCR-x polypeptides that have been covalently modified to include one or more water-soluble polymer attachments such as polyethylene glycol, polyoxyethylene glycol, or polypropylene glycol. Variants that display ligand binding properties of native nGPCR-x and are expressed at higher levels, as well as variants that provide for constitutively active receptors, are particularly useful in assays of the invention; the variants are also useful in providing cellular, tissue and animal models of diseases/conditions characterized by aberrant nGPCR-x activity.

[000122] In a related embodiment, the present invention provides compositions comprising purified polypeptides of the invention. Preferred compositions comprise, in addition to the polypeptide of the invention, a pharmaceutically acceptable (*i.e.*, sterile and non-toxic) liquid, semisolid, or solid diluent that serves as a pharmaceutical vehicle, excipient, or medium. Any diluent known in the art may be used. Exemplary diluents include, but are not limited to, water, saline solutions, polyoxyethylene sorbitan monolaurate, magnesium stearate, methyl- and propylhydroxybenzoate, talc, alginates, starches, lactose, sucrose, dextrose, sorbitol, mannitol, glycerol, calcium phosphate, mineral oil, and cocoa butter.

[000123] Variants that display ligand binding properties of native nGPCR-x and are expressed at higher levels, as well as variants that provide for constitutively active receptors, are particularly useful in assays of the invention; the variants are also useful in assays of the invention and in providing cellular, tissue and animal models of diseases/conditions characterized by aberrant nGPCR-x activity.

[000124] The G protein-coupled receptor functions through a specific heterotrimeric guanine-nucleotide-binding regulatory protein (G-protein) coupled to the intracellular portion of the G protein-coupled receptor molecule. Accordingly, the G protein-coupled receptor has a specific affinity to G protein. G proteins specifically bind to guanine nucleotides. Isolation of G proteins provides a means to isolate guanine nucleotides. G proteins may be isolated using commercially available anti-G protein antibodies or isolated G protein-coupled receptors. Similarly, G proteins may be detected in a sample isolated using commercially available detectable anti-G protein antibodies or isolated G protein-coupled receptors.

[000125] According to the present invention, the isolated nGPCR-x proteins of the present invention are useful to isolate and purify G proteins from samples such as cell lysates. Example 15 below sets forth an example of isolation of G proteins using isolated nGPCR-x proteins. Such methodology may be used in place of the use of commercially available anti-G protein antibodies which are used to isolate G proteins. Moreover, G proteins may be detected using n-GPCR-x proteins in place of commercially available detectable anti-G protein antibodies. Since nGPCR-x proteins specifically bind to G proteins, they can be employed in any specific use

where G protein specific affinity is required such as those uses where commercially available anti-G protein antibodies are employed.

Antibodies

[000126] Also comprehended by the present invention are antibodies (e.g., monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, and complementary determining region (CDR)-grafted antibodies, including compounds which include CDR sequences which specifically recognize a polypeptide of the invention) specific for nGPCR-x or fragments thereof. Preferred antibodies of the invention are human antibodies that are produced and identified according to methods described in WO93/11236, published June 20, 1993, which is incorporated herein by reference in its entirety. Antibody fragments, including Fab, Fab', F(ab')₂, and F_v, are also provided by the invention. The term "specific for," when used to describe antibodies of the invention, indicates that the variable regions of the antibodies of the invention recognize and bind nGPCR-x polypeptides exclusively (i.e., are able to distinguish nGPCR-x polypeptides from other known GPCR polypeptides by virtue of measurable differences in binding affinity, despite the possible existence of localized sequence identity, homology, or similarity between nGPCR-x and such polypeptides). It will be understood that specific antibodies may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and, in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow et al. (Eds.), Antibodies A Laboratory Manual; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988), Chapter 6. Antibodies that recognize and bind fragments of the nGPCR-x polypeptides of the invention are also contemplated, provided that the antibodies are specific for nGPCR-x polypeptides. Antibodies of the invention can be produced using any method well known and routinely practiced in the art.

[000127] The invention provides an antibody that is specific for the nGPCR-x of the invention. Antibody specificity is described in greater detail below. However, it should be emphasized that antibodies that can be generated from polypeptides that have previously been described in the literature and that are capable of fortuitously cross-reacting with nGPCR-x (e.g., due to the fortuitous existence of a similar epitope in both polypeptides) are considered "cross-reactive" antibodies. Such cross-reactive antibodies are not antibodies that are "specific" for nGPCR-x. The determination of whether an antibody is specific for nGPCR-x or is cross-reactive with another known receptor is made using any of several assays, such as Western blotting assays,

that are well known in the art. For identifying cells that express nGPCR-x and also for modulating nGPCR-x-ligand binding activity, antibodies that specifically bind to an extracellular epitope of the nGPCR-x are preferred.

[000128] In one preferred variation, the invention provides monoclonal antibodies. Hybridomas that produce such antibodies also are intended as aspects of the invention. In yet another variation, the invention provides a humanized antibody. Humanized antibodies are useful for *in vivo* therapeutic indications.

[000129] In another variation, the invention provides a cell-free composition comprising polyclonal antibodies, wherein at least one of the antibodies is an antibody of the invention specific for nGPCR-x. Antisera isolated from an animal is an exemplary composition, as is a composition comprising an antibody fraction of an antisera that has been resuspended in water or in another diluent, excipient, or carrier.

[000130] In still another related embodiment, the invention provides an anti-idiotypic antibody specific for an antibody that is specific for nGPCR-x.

[000131] It is well known that antibodies contain relatively small antigen binding domains that can be isolated chemically or by recombinant techniques. Such domains are useful nGPCR-x binding molecules themselves, and also may be reintroduced into human antibodies, or fused to toxins or other polypeptides. Thus, in still another embodiment, the invention provides a polypeptide comprising a fragment of a nGPCR-x-specific antibody, wherein the fragment and the polypeptide bind to the nGPCR-x. By way of non-limiting example, the invention provides polypeptides that are single chain antibodies and CDR-grafted antibodies.

[000132] Non-human antibodies may be humanized by any of the methods known in the art. In one method, the non-human CDRs are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

[000133] Antibodies of the invention are useful for, *e.g.*, therapeutic purposes (by modulating activity of nGPCR-x), diagnostic purposes to detect or quantitate nGPCR-x, and purification of nGPCR-x. Kits comprising an antibody of the invention for any of the purposes described herein are also comprehended. In general, a kit of the invention also includes a control antigen for which the antibody is immunospecific.

Compositions

[000134] Mutations in the nGPCR-x gene that result in loss of normal function of the nGPCR-x gene product underlie nGPCR-x-related human disease states. The invention comprehends gene therapy to restore nGPCR-x activity to treat those disease states. Delivery of a functional nGPCR-x gene to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and

more particularly viral vectors (e.g., adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (e.g., liposomes or chemical treatments). See, for example, Anderson, *Nature*, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, *Science*, 244: 1275-1281 (1989); Verma, *Scientific American*: 68-84 (1990); and Miller, *Nature*, 357: 455-460 (1992). Alternatively, it is contemplated that in other human disease states, preventing the expression of, or inhibiting the activity of, nGPCR-x will be useful in treating disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of nGPCR-x.

[000135] Another aspect of the present invention is directed to compositions, including pharmaceutical compositions, comprising any of the nucleic acid molecules or recombinant expression vectors described above and an acceptable carrier or diluent. Preferably, the carrier or diluent is pharmaceutically acceptable. Suitable carriers are described in the most recent edition of *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field, which is incorporated herein by reference in its entirety. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The formulations are sterilized by commonly used techniques.

[000136] Also within the scope of the invention are compositions comprising polypeptides, polynucleotides, or antibodies of the invention that have been formulated with, e.g., a pharmaceutically acceptable carrier.

[000137] The invention also provides methods of using antibodies of the invention. For example, the invention provides a method for modulating ligand binding of a nGPCR-x comprising the step of contacting the nGPCR-x with an antibody specific for the nGPCR-x, under conditions wherein the antibody binds the receptor.

[000138] As discussed above, it is well known that GPCRs are expressed in many different tissues and regions, including in the brain. GPCRs that may be expressed in the brain, such as nGPCR-x, provide an indication that aberrant nGPCR-x signaling activity may correlate with one or more neurological or psychological disorders. The invention also provides a method for treating a neurological or psychiatric disorder comprising the step of administering to a mammal in need of such treatment an amount of an antibody-like polypeptide of the invention that is sufficient to modulate ligand binding to a nGPCR-x in neurons of the mammal. nGPCR-x may also be expressed in other tissues, including but not limited to, peripheral blood lymphocytes, pancreas, ovary, uterus, testis, salivary gland, thyroid gland, kidney, adrenal gland, liver, bone marrow, prostate, fetal liver, colon, muscle, and fetal brain, and may be found in many other tissues.

Within the brain, nGPCR-x mRNA transcripts may be found in many tissues, including, but not limited to, frontal lobe, hypothalamus, pons, cerebellum, caudate nucleus, and medulla.

Kits

[000139] The present invention is also directed to kits, including pharmaceutical kits. The kits can comprise any of the nucleic acid molecules described above, any of the polypeptides described above, or any antibody which binds to a polypeptide of the invention as described above, as well as a negative control. The kit preferably comprises additional components, such as, for example, instructions, solid support, reagents helpful for quantification, and the like.

[000140] In another aspect, the invention features methods for detection of a polypeptide in a sample as a diagnostic tool for diseases or disorders, wherein the method comprises the steps of: (a) contacting the sample with a nucleic acid probe which hybridizes under hybridization assay conditions to a nucleic acid target region of a polypeptide having sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, said probe comprising the nucleic acid sequence encoding the polypeptide, fragments thereof, and the complements of the sequences and fragments; and (b) detecting the presence or amount of the probe:target region hybrid as an indication of the disease.

[000141] In preferred embodiments of the invention, the disease is selected from the group consisting of thyroid disorders (*e.g.* thyrotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Crohn's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post-traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); and sexual dysfunction, among others.

[000142] Kits may be designed to detect either expression of polynucleotides encoding nGPCR-x expressed in the brain or the nGPCR-x proteins themselves in order to identify tissue as being

neurological. For example, oligonucleotide hybridization kits can be provided which include a container having an oligonucleotide probe specific for the nGPCR-x-specific DNA and optionally, containers with positive and negative controls and/or instructions. Similarly, PCR kits can be provided which include a container having primers specific for the nGPCR-x-specific sequences, DNA and optionally, containers with size markers, positive and negative controls and/or instructions.

[000143] Hybridization conditions should be such that hybridization occurs only with the genes in the presence of other nucleic acid molecules. Under stringent hybridization conditions only highly complementary nucleic acid sequences hybridize. Preferably, such conditions prevent hybridization of nucleic acids having 1 or 2 mismatches out of 20 contiguous nucleotides. Such conditions are defined supra.

[000144] The diseases for which detection of genes in a sample could be diagnostic include diseases in which nucleic acid (DNA and/or RNA) is amplified in comparison to normal cells. By "amplification" is meant increased numbers of DNA or RNA in a cell compared with normal cells.

[000145] The diseases that could be diagnosed by detection of nucleic acid in a sample preferably include central nervous system and metabolic diseases. The test samples suitable for nucleic acid probing methods of the present invention include, for example, cells or nucleic acid extracts of cells, or biological fluids. The samples used in the above-described methods will vary based on the assay format, the detection method and the nature of the tissues, cells or extracts to be assayed. Methods for preparing nucleic acid extracts of cells are well known in the art and can be readily adapted in order to obtain a sample that is compatible with the method utilized.

[000146] Alternatively, immunoassay kits can be provided which have containers container having antibodies specific for the nGPCR-x-protein and optionally, containers with positive and negative controls and/or instructions.

[000147] Kits may also be provided useful in the identification of GPCR binding partners such as natural ligands or modulators (agonists or antagonists). Substances useful for treatment of disorders or diseases preferably show positive results in one or more *in vitro* assays for an activity corresponding to treatment of the disease or disorder in question. Substances that modulate the activity of the polypeptides preferably include, but are not limited to, antisense oligonucleotides, agonists and antagonists, and inhibitors of protein kinases.

Methods of inducing immune response

[000148] Another aspect of the present invention is directed to methods of inducing an immune response in a mammal against a polypeptide of the invention by administering to the mammal an amount of the polypeptide sufficient to induce an immune response. The amount will be

dependent on the animal species, size of the animal, and the like but can be determined by those skilled in the art.

Methods of identifying ligands

[000149] The invention also provides assays to identify compounds that bind nGPCR-x. One such assay comprises the steps of: (a) contacting a composition comprising a nGPCR-x with a compound suspected of binding nGPCR-x; and (b) measuring binding between the compound and nGPCR-x. In one variation, the composition comprises a cell expressing nGPCR-x on its surface. In another variation, isolated nGPCR-x or cell membranes comprising nGPCR-x are employed. The binding may be measured directly, *e.g.*, by using a labeled compound, or may be measured indirectly by several techniques, including measuring intracellular signaling of nGPCR-x induced by the compound (or measuring changes in the level of nGPCR-x signaling). Following steps (a) and (b), compounds identified as binding nGPCR-x may be tested in other assays including, but not limited to, *in vivo* models, to confirm or quantitate binding to nGPCR-x.

[000150] Specific binding molecules, including natural ligands and synthetic compounds, can be identified or developed using isolated or recombinant nGPCR-x products, nGPCR-x variants, or preferably, cells expressing such products. Binding partners are useful for purifying nGPCR-x products and detection or quantification of nGPCR-x products in fluid and tissue samples using known immunological procedures. Binding molecules are also manifestly useful in modulating (*i.e.*, blocking, inhibiting or stimulating) biological activities of nGPCR-x, especially those activities involved in signal transduction.

[000151] The DNA and amino acid sequence information provided by the present invention also makes possible identification of binding partner compounds with which a nGPCR-x polypeptide or polynucleotide will interact. Methods to identify binding partner compounds include solution assays, *in vitro* assays wherein nGPCR-x polypeptides are immobilized, and cell-based assays. Identification of binding partner compounds of nGPCR-x polypeptides provides candidates for therapeutic or prophylactic intervention in pathologies associated with nGPCR-x normal and aberrant biological activity.

[000152] The invention includes several assay systems for identifying nGPCR-x binding partners. In solution assays, methods of the invention comprise the steps of (a) contacting a nGPCR-x polypeptide with one or more candidate binding partner compounds and (b) identifying the compounds that bind to the nGPCR-x polypeptide. Identification of the compounds that bind the nGPCR-x polypeptide can be achieved by isolating the nGPCR-x polypeptide/binding partner complex, and separating the binding partner compound from the nGPCR-x polypeptide. An additional step of characterizing the physical, biological, and/or biochemical properties of

the binding partner compound is also comprehended in another embodiment of the invention, wherein compounds identified as binding nGPCR-x may be tested in other assays including, but not limited to, *in vivo* models, to confirm or quantitate binding to nGPCR-x. In one aspect, the nGPCR-x polypeptide/binding partner complex is isolated using an antibody immunospecific for either the nGPCR-x polypeptide or the candidate binding partner compound.

[000153] In still other embodiments, either the nGPCR-x polypeptide or the candidate binding partner compound comprises a label or tag that facilitates its isolation, and methods of the invention to identify binding partner compounds include a step of isolating the nGPCR-x polypeptide/binding partner complex through interaction with the label or tag. An exemplary tag of this type is a poly-histidine sequence, generally around six histidine residues, that permits isolation of a compound so labeled using nickel chelation. Other labels and tags, such as the FLAG[®] tag (Eastman Kodak, Rochester, NY), well known and routinely used in the art, are embraced by the invention.

[000154] In one variation of an *in vitro* assay, the invention provides a method comprising the steps of (a) contacting an immobilized nGPCR-x polypeptide with a candidate binding partner compound and (b) detecting binding of the candidate compound to the nGPCR-x polypeptide. In an alternative embodiment, the candidate binding partner compound is immobilized and binding of nGPCR-x is detected. Immobilization is accomplished using any of the methods well known in the art, including covalent bonding to a support, a bead, or a chromatographic resin, as well as non-covalent, high affinity interactions such as antibody binding, or use of streptavidin/biotin binding wherein the immobilized compound includes a biotin moiety. Detection of binding can be accomplished (i) using a radioactive label on the compound that is not immobilized, (ii) using of a fluorescent label on the non-immobilized compound, (iii) using an antibody immunospecific for the non-immobilized compound, (iv) using a label on the non-immobilized compound that excites a fluorescent support to which the immobilized compound is attached, as well as other techniques well known and routinely practiced in the art.

[000155] The invention also provides cell-based assays to identify binding partner compounds of a nGPCR-x polypeptide. In one embodiment, the invention provides a method comprising the steps of contacting a nGPCR-x polypeptide expressed on the surface of a cell with a candidate binding partner compound and detecting binding of the candidate binding partner compound to the nGPCR-x polypeptide. In a preferred embodiment, the detection comprises detecting a calcium flux or other physiological event in the cell caused by the binding of the molecule.

[000156] Another aspect of the present invention is directed to methods of identifying compounds that bind to either nGPCR-x or nucleic acid molecules encoding nGPCR-x, comprising contacting nGPCR-x, or a nucleic acid molecule encoding the same, with a compound, and

determining whether the compound binds nGPCR-x or a nucleic acid molecule encoding the same. Binding can be determined by binding assays which are well known to the skilled artisan, including, but not limited to, gel-shift assays, Western blots, radiolabeled competition assay, phage-based expression cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, ELISA, and the like, which are described in, for example, *Current Protocols in Molecular Biology*, 1999, John Wiley & Sons, NY, which is incorporated herein by reference in its entirety. The compounds to be screened include (which may include compounds which are suspected to bind nGPCR-x, or a nucleic acid molecule encoding the same), but are not limited to, extracellular, intracellular, biologic or chemical origin. The methods of the invention also embrace ligands, especially neuropeptides, that are attached to a label, such as a radiolabel (e.g., ^{125}I , ^{35}S , ^{32}P , ^{33}P , ^3H), a fluorescence label, a chemiluminescent label, an enzymic label and an immunogenic label. Modulators falling within the scope of the invention include, but are not limited to, non-peptide molecules such as non-peptide mimetics, non-peptide allosteric effectors, and peptides. The nGPCR-x polypeptide or polynucleotide employed in such a test may either be free in solution, attached to a solid support, borne on a cell surface or located intracellularly or associated with a portion of a cell. One skilled in the art can, for example, measure the formation of complexes between nGPCR-x and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between nGPCR-x and its substrate caused by the compound being tested.

[000157] In another embodiment of the invention, high throughput screening for compounds having suitable binding affinity to nGPCR-x is employed. Briefly, large numbers of different test compounds are synthesized on a solid substrate. The peptide test compounds are contacted with nGPCR-x and washed. Bound nGPCR-x is then detected by methods well known in the art. Purified polypeptides of the invention can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the protein and immobilize it on the solid support.

[000158] Generally, an expressed nGPCR-x can be used for HTS binding assays in conjunction with its defined ligand, in this case the corresponding neuropeptide that activates it. The identified peptide is labeled with a suitable radioisotope, including, but not limited to, ^{125}I , ^3H , ^{35}S or ^{32}P , by methods that are well known to those skilled in the art. Alternatively, the peptides may be labeled by well-known methods with a suitable fluorescent derivative (Baindur *et al.*, *Drug Dev. Res.*, 1994, 33, 373-398; Rogers, *Drug Discovery Today*, 1997, 2, 156-160). Radioactive ligand specifically bound to the receptor in membrane preparations made from the cell line expressing the recombinant protein can be detected in HTS assays in one of several

standard ways, including filtration of the receptor-ligand complex to separate bound ligand from unbound ligand (Williams, *Med. Res. Rev.*, 1991, 11, 147-184; Sweetnam *et al.*, *J. Natural Products*, 1993, 56, 441-455). Alternative methods include a scintillation proximity assay (SPA) or a FlashPlate format in which such separation is unnecessary (Nakayama, *Cur. Opinion Drug Disc. Dev.*, 1998, 1, 85-91 Bossé *et al.*, *J. Biomolecular Screening*, 1998, 3, 285-292.). Binding of fluorescent ligands can be detected in various ways, including fluorescence energy transfer (FRET), direct spectrophotofluorometric analysis of bound ligand, or fluorescence polarization (Rogers, *Drug Discovery Today*, 1997, 2, 156-160; Hill, *Cur. Opinion Drug Disc. Dev.*, 1998, 1, 92-97).

[000159] Other assays may be used to identify specific ligands of a nGPCR-x receptor, including assays that identify ligands of the target protein through measuring direct binding of test ligands to the target protein, as well as assays that identify ligands of target proteins through affinity ultrafiltration with ion spray mass spectroscopy/HPLC methods or other physical and analytical methods. Alternatively, such binding interactions are evaluated indirectly using the yeast two-hybrid system described in Fields *et al.*, *Nature*, 340:245-246 (1989), and Fieldset *al.*, *Trends in Genetics*, 10:286-292 (1994), both of which are incorporated herein by reference. The two-hybrid system is a genetic assay for detecting interactions between two proteins or polypeptides. It can be used to identify proteins that bind to a known protein of interest, or to delineate domains or residues critical for an interaction. Variations on this methodology have been developed to clone genes that encode DNA binding proteins, to identify peptides that bind to a protein, and to screen for drugs. The two-hybrid system exploits the ability of a pair of interacting proteins to bring a transcription activation domain into close proximity with a DNA binding domain that binds to an upstream activation sequence (UAS) of a reporter gene, and is generally performed in yeast. The assay requires the construction of two hybrid genes encoding (1) a DNA-binding domain that is fused to a first protein and (2) an activation domain fused to a second protein. The DNA-binding domain targets the first hybrid protein to the UAS of the reporter gene; however, because most proteins lack an activation domain, this DNA-binding hybrid protein does not activate transcription of the reporter gene. The second hybrid protein, which contains the activation domain, cannot by itself activate expression of the reporter gene because it does not bind the UAS. However, when both hybrid proteins are present, the noncovalent interaction of the first and second proteins tethers the activation domain to the UAS, activating transcription of the reporter gene. For example, when the first protein is a GPCR gene product, or fragment thereof, that is known to interact with another protein or nucleic acid, this assay can be used to detect agents that interfere with the binding interaction.

Expression of the reporter gene is monitored as different test agents are added to the system.

The presence of an inhibitory agent results in lack of a reporter signal.

[000160] The yeast two-hybrid assay can also be used to identify proteins that bind to the gene product. In an assay to identify proteins that bind to a nGPCR-x receptor, or fragment thereof, a fusion polynucleotide encoding both a nGPCR-x receptor (or fragment) and a UAS binding domain (*i.e.*, a first protein) may be used. In addition, a large number of hybrid genes each encoding a different second protein fused to an activation domain are produced and screened in the assay. Typically, the second protein is encoded by one or more members of a total cDNA or genomic DNA fusion library, with each second protein-coding region being fused to the activation domain. This system is applicable to a wide variety of proteins, and it is not even necessary to know the identity or function of the second binding protein. The system is highly sensitive and can detect interactions not revealed by other methods; even transient interactions may trigger transcription to produce a stable mRNA that can be repeatedly translated to yield the reporter protein.

[000161] Other assays may be used to search for agents that bind to the target protein. One such screening method to identify direct binding of test ligands to a target protein is described in U.S. Patent No. 5,585,277, incorporated herein by reference. This method relies on the principle that proteins generally exist as a mixture of folded and unfolded states, and continually alternate between the two states. When a test ligand binds to the folded form of a target protein (*i.e.*, when the test ligand is a ligand of the target protein), the target protein molecule bound by the ligand remains in its folded state. Thus, the folded target protein is present to a greater extent in the presence of a test ligand which binds the target protein, than in the absence of a ligand. Binding of the ligand to the target protein can be determined by any method that distinguishes between the folded and unfolded states of the target protein. The function of the target protein need not be known in order for this assay to be performed. Virtually any agent can be assessed by this method as a test ligand, including, but not limited to, metals, polypeptides, proteins, lipids, polysaccharides, polynucleotides and small organic molecules.

[000162] Another method for identifying ligands of a target protein is described in Wieboldt *et al.*, Anal. Chem., 69:1683-1691 (1997), incorporated herein by reference. This technique screens combinatorial libraries of 20-30 agents at a time in solution phase for binding to the target protein. Agents that bind to the target protein are separated from other library components by simple membrane washing. The specifically selected molecules that are retained on the filter are subsequently liberated from the target protein and analyzed by HPLC and pneumatically assisted electrospray (ion spray) ionization mass spectroscopy. This procedure selects library

components with the greatest affinity for the target protein, and is particularly useful for small molecule libraries.

[000163] Other embodiments of the invention comprise using competitive screening assays in which neutralizing antibodies capable of binding a polypeptide of the invention specifically compete with a test compound for binding to the polypeptide. In this manner, the antibodies can be used to detect the presence of any peptide that shares one or more antigenic determinants with nGPCR-x. Radiolabeled competitive binding studies are described in A.H. Lin *et al.* *Antimicrobial Agents and Chemotherapy*, 1997, vol. 41, no. 10. pp. 2127-2131, the disclosure of which is incorporated herein by reference in its entirety.

Identification of modulating agents

[000164] The invention also provides methods for identifying a modulator of binding between a nGPCR-x and a nGPCR-x binding partner, comprising the steps of: (a) contacting a nGPCR-x binding partner and a composition comprising a nGPCR-x in the presence and in the absence of a putative modulator compound; (b) detecting binding between the binding partner and the nGPCR-x; and (c) identifying a putative modulator compound or a modulator compound in view of decreased or increased binding between the binding partner and the nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator. Following steps (a) and (b), compounds identified as modulating binding between nGPCR-x and a nGPCR-x binding partner may be tested in other assays including, but not limited to, *in vivo* models, to confirm or quantitate modulation of binding to nGPCR-x.

[000165] nGPCR-x binding partners that stimulate nGPCR-x activity are useful as agonists in disease states or conditions characterized by insufficient nGPCR-x signaling (*e.g.*, as a result of insufficient activity of a nGPCR-x ligand). nGPCR-x binding partners that block ligand-mediated nGPCR-x signaling are useful as nGPCR-x antagonists to treat disease states or conditions characterized by excessive nGPCR-x signaling. In addition nGPCR-x modulators in general, as well as nGPCR-x polynucleotides and polypeptides, are useful in diagnostic assays for such diseases or conditions.

[000166] In another aspect, the invention provides methods for treating a disease or abnormal condition by administering to a patient in need of such treatment a substance that modulates the activity or expression of a polypeptide having sequences selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116.

[000167] Agents that modulate (*i.e.*, increase, decrease, or block) nGPCR-x activity or expression may be identified by incubating a putative modulator with a cell containing a nGPCR-x polypeptide or polynucleotide and determining the effect of the putative modulator on nGPCR-x activity or expression. The selectivity of a compound that modulates the activity of nGPCR-x

can be evaluated by comparing its effects on nGPCR-x to its effect on other GPCR compounds. Following identification of compounds that modulate nGPCR-x activity or expression, such compounds may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity. Selective modulators may include, for example, antibodies and other proteins, peptides, or organic molecules that specifically bind to a nGPCR-x polypeptide or a nGPCR-x-encoding nucleic acid. Modulators of nGPCR-x activity will be therapeutically useful in treatment of diseases and physiological conditions in which normal or aberrant nGPCR-x activity is involved. nGPCR-x polynucleotides, polypeptides, and modulators may be used in the treatment of such diseases and conditions as infections, such as viral infections caused by HIV-1 or HIV-2; pain; cancers; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); Parkinson's disease; and psychotic and neurological disorders, including schizophrenia, migraine, ADHH, major depression, anxiety, mental disorder, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome, among others. nGPCR-x polynucleotides and polypeptides, as well as nGPCR-x modulators, may also be used in diagnostic assays for such diseases or conditions.

[000168] Methods of the invention to identify modulators include variations on any of the methods described above to identify binding partner compounds, the variations including techniques wherein a binding partner compound has been identified and the binding assay is carried out in the presence and absence of a candidate modulator. A modulator is identified in those instances where binding between the nGPCR-x polypeptide and the binding partner compound changes in the presence of the candidate modulator compared to binding in the absence of the candidate modulator compound. A modulator that increases binding between the nGPCR-x polypeptide and the binding partner compound is described as an enhancer or activator, and a modulator that decreases binding between the nGPCR-x polypeptide and the binding partner compound is described as an inhibitor. Following identification of modulators, such compounds may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity as modulators.

[000169] The invention also comprehends high-throughput screening (HTS) assays to identify compounds that interact with or inhibit biological activity (*i.e.*, affect enzymatic activity, binding activity, *etc.*) of a nGPCR-x polypeptide. HTS assays permit screening of large numbers of compounds in an efficient manner. Cell-based HTS systems are contemplated to investigate nGPCR-x receptor-ligand interaction. HTS assays are designed to identify "hits" or "lead compounds" having the desired property, from which modifications can be designed to

improve the desired property. Chemical modification of the "hit" or "lead compound" is often based on an identifiable structure/activity relationship between the "hit" and the nGPCR-x polypeptide.

[000170] Another aspect of the present invention is directed to methods of identifying compounds which modulate (*i.e.*, increase or decrease) an activity of nGPCR-x comprising contacting nGPCR-x with a compound, and determining whether the compound modifies activity of nGPCR-x. The activity in the presence of the test compared is measured to the activity in the absence of the test compound. Where the activity of the sample containing the test compound is higher than the activity in the sample lacking the test compound, the compound will have increased activity. Similarly, where the activity of the sample containing the test compound is lower than the activity in the sample lacking the test compound, the compound will have inhibited activity. Following the identification of compounds that modulate an activity of nGPCR-x, such compounds can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity.

[000171] The present invention is particularly useful for screening compounds by using nGPCR-x in any of a variety of drug screening techniques. The compounds to be screened include (which may include compounds which are suspected to modulate nGPCR-x activity), but are not limited to, extracellular, intracellular, biologic or chemical origin. The nGPCR-x polypeptide employed in such a test may be in any form, preferably, free in solution, attached to a solid support, borne on a cell surface or located intracellularly. One skilled in the art can, for example, measure the formation of complexes between nGPCR-x and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between nGPCR-x and its substrate caused by the compound being tested.

[000172] The activity of nGPCR-x polypeptides of the invention can be determined by, for example, examining the ability to bind or be activated by chemically synthesized peptide ligands. Alternatively, the activity of nGPCR-x polypeptides can be assayed by examining their ability to bind calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and photons. Alternatively, the activity of the nGPCR-x polypeptides can be determined by examining the activity of effector molecules including, but not limited to, adenylate cyclase, phospholipases and ion channels. Thus, modulators of nGPCR-x polypeptide activity may alter a GPCR receptor function, such as a binding property of a receptor or an activity such as G protein-mediated signal transduction or membrane localization. In various embodiments of the method, the assay may take the form of an ion flux assay, a yeast growth assay, a non-hydrolyzable GTP assay such as a [35 S]-GTP γ S assay, a cAMP assay, an inositol triphosphate assay, a diacylglycerol assay, an Aequorin assay, a

Luciferase assay, a FLIPR assay for intracellular Ca^{2+} concentration, a mitogenesis assay, a MAP Kinase activity assay, an arachidonic acid release assay (e.g., using [^3H]-arachidonic acid), and an assay for extracellular acidification rates, as well as other binding or function-based assays of nGPCR-x activity that are generally known in the art. In several of these embodiments, the invention comprehends the inclusion of any of the G proteins known in the art, such as G_{16} , G_{15} , or chimeric G_{q45} , G_{q55} , G_{q65} , G_{q25} , and the like. nGPCR-x activity can be determined by methodologies that are used to assay for FaRP activity, which is well known to those skilled in the art. Biological activities of nGPCR-x receptors according to the invention include, but are not limited to, the binding of a natural or an unnatural ligand, as well as any one of the functional activities of GPCRs known in the art. Non-limiting examples of GPCR activities include transmembrane signaling of various forms, which may involve G protein association and/or the exertion of an influence over G protein binding of various guanidylate nucleotides; another exemplary activity of GPCRs is the binding of accessory proteins or polypeptides that differ from known G proteins.

[000173] The modulators of the invention exhibit a variety of chemical structures, which can be generally grouped into non-peptide mimetics of natural GPCR receptor ligands, peptide and non-peptide allosteric effectors of GPCR receptors, and peptides that may function as activators or inhibitors (competitive, uncompetitive and non-competitive) (e.g., antibody products) of GPCR receptors. The invention does not restrict the sources for suitable modulators, which may be obtained from natural sources such as plant, animal or mineral extracts, or non-natural sources such as small molecule libraries, including the products of combinatorial chemical approaches to library construction, and peptide libraries. Examples of peptide modulators of GPCR receptors exhibit the following primary structures: GLGPRPLRFamide, GNSFLRFamide, GGPQGPLRFamide, GPSGPLRFamide, PDVDHVFLRFamide, and pyro-EDVDHVFLRFamide.

[000174] Other assays can be used to examine enzymatic activity including, but not limited to, photometric, radiometric, HPLC, electrochemical, and the like, which are described in, for example, *Enzyme Assays: A Practical Approach*, eds. R. Eisinger and M. J. Danson, 1992, Oxford University Press, which is incorporated herein by reference in its entirety.

[000175] The use of cDNAs encoding GPCRs in drug discovery programs is well-known; assays capable of testing thousands of unknown compounds per day in high-throughput screens (HTSs) are thoroughly documented. The literature is replete with examples of the use of radiolabeled ligands in HTS binding assays for drug discovery (see Williams, *Medicinal Research Reviews*, 1991, 11, 147-184.; Sweetnam, *et al.*, *J. Natural Products*, 1993, 56, 441-455 for review). Recombinant receptors are preferred for binding assay HTS because they allow for better

specificity (higher relative purity), provide the ability to generate large amounts of receptor material, and can be used in a broad variety of formats (see Hodgson, *Bio/Technology*, 1992, 10, 973-980; each of which is incorporated herein by reference in its entirety).

[000176] A variety of heterologous systems is available for functional expression of recombinant receptors that are well known to those skilled in the art. Such systems include bacteria (Strosberg, *et al.*, *Trends in Pharmacological Sciences*, 1992, 13, 95-98), yeast (Pausch, *Trends in Biotechnology*, 1997, 15, 487-494), several kinds of insect cells (Vanden Broeck, *Int. Rev. Cytology*, 1996, 164, 189-268), amphibian cells (Jayawickreme *et al.*, *Current Opinion in Biotechnology*, 1997, 8, 629-634) and several mammalian cell lines (CHO, HEK-293, COS, etc.; see Gerhardt, *et al.*, *Eur. J. Pharmacology*, 1997, 334, 1-23). These examples do not preclude the use of other possible cell expression systems, including cell lines obtained from nematodes (PCT application WO 98/37177).

[000177] In preferred embodiments of the invention, methods of screening for compounds that modulate nGPCR-x activity comprise contacting test compounds with nGPCR-x and assaying for the presence of a complex between the compound and nGPCR-x. In such assays, the ligand is typically labeled. After suitable incubation, free ligand is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular compound to bind to nGPCR-x.

[000178] It is well known that activation of heterologous receptors expressed in recombinant systems results in a variety of biological responses, which are mediated by G proteins expressed in the host cells. Occupation of a GPCR by an agonist results in exchange of bound GDP for GTP at a binding site on the G_{α} subunit; one can use a radioactive, non-hydrolyzable derivative of GTP, $GTP\gamma[^{35}S]$, to measure binding of an agonist to the receptor (Sim *et al.*, *Neuroreport*, 1996, 7, 729-733). One can also use this binding to measure the ability of antagonists to bind to the receptor by decreasing binding of $GTP\gamma[^{35}S]$ in the presence of a known agonist. One could therefore construct a HTS based on $GTP\gamma[^{35}S]$ binding, though this is not the preferred method.

[000179] The G proteins required for functional expression of heterologous GPCRs can be native constituents of the host cell or can be introduced through well-known recombinant technology. The G proteins can be intact or chimeric. Often, a nearly universally competent G protein (e.g., $G_{\alpha 16}$) is used to couple any given receptor to a detectable response pathway. G protein activation results in the stimulation or inhibition of other native proteins, events that can be linked to a measurable response.

[000180] Examples of such biological responses include, but are not limited to, the following: the ability to survive in the absence of a limiting nutrient in specifically engineered yeast cells (Pausch, *Trends in Biotechnology*, 1997, 15, 487-494); changes in intracellular Ca^{2+}

concentration as measured by fluorescent dyes (Murphy, *et al.*, *Cur. Opinion Drug Disc. Dev.*, 1998, 1, 192-199). Fluorescence changes can also be used to monitor ligand-induced changes in membrane potential or intracellular pH; an automated system suitable for HTS has been described for these purposes (Schroeder, *et al.*, *J. Biomolecular Screening*, 1996, 1, 75-80). Melanophores prepared from *Xenopus laevis* show a ligand-dependent change in pigment organization in response to heterologous GPCR activation; this response is adaptable to HTS formats (Jayawickreme *et al.*, *Cur. Opinion Biotechnology*, 1997, 8, 629-634). Assays are also available for the measurement of common second messengers, including cAMP, phosphoinositides and arachidonic acid, but these are not generally preferred for HTS.

[000181] Preferred methods of HTS employing these receptors include permanently transfected CHO cells, in which agonists and antagonists can be identified by the ability to specifically alter the binding of GTP γ [³⁵S] in membranes prepared from these cells. In another embodiment of the invention, permanently transfected CHO cells could be used for the preparation of membranes which contain significant amounts of the recombinant receptor proteins; these membrane preparations would then be used in receptor binding assays, employing the radiolabeled ligand specific for the particular receptor. Alternatively, a functional assay, such as fluorescent monitoring of ligand-induced changes in internal Ca²⁺ concentration or membrane potential in permanently transfected CHO cells containing each of these receptors individually or in combination would be preferred for HTS. Equally preferred would be an alternative type of mammalian cell, such as HEK-293 or COS cells, in similar formats. More preferred would be permanently transfected insect cell lines, such as *Drosophila* S2 cells. Even more preferred would be recombinant yeast cells expressing the *Drosophila melanogaster* receptors in HTS formats well known to those skilled in the art (*e.g.*, Pausch, *Trends in Biotechnology*, 1997, 15, 487-494).

[000182] The invention contemplates a multitude of assays to screen and identify inhibitors of ligand binding to nGPCR-x receptors. In one example, the nGPCR-x receptor is immobilized and interaction with a binding partner is assessed in the presence and absence of a candidate modulator such as an inhibitor compound. In another example, interaction between the nGPCR-x receptor and its binding partner is assessed in a solution assay, both in the presence and absence of a candidate inhibitor compound. In either assay, an inhibitor is identified as a compound that decreases binding between the nGPCR-x receptor and its binding partner. Following the identification of compounds which inhibit ligand binding to nGPCR-x receptors, such compounds may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity. Another contemplated assay involves a variation of the dihybrid assay wherein an inhibitor of protein/protein interactions is identified

by detection of a positive signal in a transformed or transfected host cell, as described in PCT publication number WO 95/20652, published August 3, 1995.

[000183] Candidate modulators contemplated by the invention include compounds selected from libraries of either potential activators or potential inhibitors. There are a number of different libraries used for the identification of small molecule modulators, including: (1) chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules. Chemical libraries consist of random chemical structures, some of which are analogs of known compounds or analogs of compounds that have been identified as "hits" or "leads" in other drug discovery screens, some of which are derived from natural products, and some of which arise from non-directed synthetic organic chemistry. Natural product libraries are collections of microorganisms, animals, plants, or marine organisms which are used to create mixtures for screening by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of plants or marine organisms. Natural product libraries include polyketides, non-ribosomal peptides, and variants (non-naturally occurring) thereof. For a review, see Science 282:63-68 (1998). Combinatorial libraries are composed of large numbers of peptides, oligonucleotides, or organic compounds as a mixture. These libraries are relatively easy to prepare by traditional automated synthesis methods, PCR, cloning, or proprietary synthetic methods. Of particular interest are non-peptide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created therefrom, see Myers, Curr. Opin. Biotechnol. 8:701-707 (1997). Identification of modulators through use of the various libraries described herein permits modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to modulate activity.

[000184] Still other candidate inhibitors contemplated by the invention can be designed and include soluble forms of binding partners, as well as such binding partners as chimeric, or fusion, proteins. A "binding partner" as used herein broadly encompasses non-peptide modulators, as well as such peptide modulators as neuropeptides other than natural ligands, antibodies, antibody fragments, and modified compounds comprising antibody domains that are immunospecific for the expression product of the identified nGPCR-x gene.

[000185] The polypeptides of the invention are employed as a research tool for identification, characterization and purification of interacting, regulatory proteins. Appropriate labels are incorporated into the polypeptides of the invention by various methods known in the art and the polypeptides are used to capture interacting molecules. For example, molecules are incubated with the labeled polypeptides, washed to remove unbound polypeptides, and the polypeptide

complex is quantified. Data obtained using different concentrations of polypeptide are used to calculate values for the number, affinity, and association of polypeptide with the protein complex.

[000186] Labeled polypeptides are also useful as reagents for the purification of molecules with which the polypeptide interacts including, but not limited to, inhibitors. In one embodiment of affinity purification, a polypeptide is covalently coupled to a chromatography column. Cells and their membranes are extracted, and various cellular subcomponents are passed over the column. Molecules bind to the column by virtue of their affinity to the polypeptide. The polypeptide-complex is recovered from the column, dissociated and the recovered molecule is subjected to protein sequencing. This amino acid sequence is then used to identify the captured molecule or to design degenerate oligonucleotides for cloning the corresponding gene from an appropriate cDNA library.

[000187] Alternatively, compounds may be identified which exhibit similar properties to the ligand for the nGPCR-x of the invention, but which are smaller and exhibit a longer half time than the endogenous ligand in a human or animal body. When an organic compound is designed, a molecule according to the invention is used as a "lead" compound. The design of mimetics to known pharmaceutically active compounds is a well-known approach in the development of pharmaceuticals based on such "lead" compounds. Mimetic design, synthesis and testing are generally used to avoid randomly screening a large number of molecules for a target property. Furthermore, structural data deriving from the analysis of the deduced amino acid sequences encoded by the DNAs of the present invention are useful to design new drugs, more specific and therefore with a higher pharmacological potency.

[000188] Comparison of the protein sequence of the present invention with the sequences present in all the available databases showed a significant homology with the transmembrane portion of G protein coupled receptors. Accordingly, computer modeling can be used to develop a putative tertiary structure of the proteins of the invention based on the available information of the transmembrane domain of other proteins. Thus, novel ligands based on the predicted structure of nGPCR-x can be designed.

[000189] In a particular embodiment, the novel molecules identified by the screening methods according to the invention are low molecular weight organic molecules, in which case a composition or pharmaceutical composition can be prepared thereof for oral intake, such as in tablets. The compositions, or pharmaceutical compositions, comprising the nucleic acid molecules, vectors, polypeptides, antibodies and compounds identified by the screening methods described herein, can be prepared for any route of administration including, but not limited to, oral, intravenous, cutaneous, subcutaneous, nasal, intramuscular or intraperitoneal. The nature

of the carrier or other ingredients will depend on the specific route of administration and particular embodiment of the invention to be administered. Examples of techniques and protocols that are useful in this context are, *inter alia*, found in Remington's Pharmaceutical Sciences, 16th edition, Osol, A (ed.), 1980, which is incorporated herein by reference in its entirety.

[000190] The dosage of these low molecular weight compounds will depend on the disease state or condition to be treated and other clinical factors such as weight and condition of the human or animal and the route of administration of the compound. For treating human or animals, between approximately 0.5 mg/kg of body weight to 500 mg/kg of body weight of the compound can be administered. Therapy is typically administered at lower dosages and is continued until the desired therapeutic outcome is observed.

[000191] The present compounds and methods, including nucleic acid molecules, polypeptides, antibodies, compounds identified by the screening methods described herein, have a variety of pharmaceutical applications and may be used, for example, to treat or prevent unregulated cellular growth, such as cancer cell and tumor growth. In a particular embodiment, the present molecules are used in gene therapy. For a review of gene therapy procedures, see e.g. Anderson, *Science*, 1992, 256, 808-813, which is incorporated herein by reference in its entirety.

[000192] The present invention also encompasses a method of agonizing (stimulating) or antagonizing a nGPCR-x natural binding partner associated activity in a mammal comprising administering to said mammal an agonist or antagonist to one of the above disclosed polypeptides in an amount sufficient to effect said agonism or antagonism. One embodiment of the present invention, then, is a method of treating diseases in a mammal with an agonist or antagonist of the protein of the present invention comprises administering the agonist or antagonist to a mammal in an amount sufficient to agonize or antagonize nGPCR-x-associated functions.

[000193] In an effort to discover novel treatments for diseases, biomedical researchers and chemists have designed, synthesized, and tested molecules that modulate the function of G protein coupled receptors. Some small organic molecules form a class of compounds that modulate the function of G protein coupled receptors.

[000194] Exemplary diseases and conditions amenable to treatment based on the present invention include, but are not limited to, thyroid disorders (e.g. thyrotoxicosis, myxoedema); renal failure; inflammatory conditions (e.g., Chron's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (e.g., pain including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post

traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); sexual dysfunction, among others.

[000195] Methods of determining the dosages of compounds to be administered to a patient and modes of administering compounds to an organism are disclosed in U.S. Application Serial No. 08/702,282, filed August 23, 1996 and International patent publication number WO 96/22976, published August 1 1996, both of which are incorporated herein by reference in their entirety, including any drawings, figures or tables. Those skilled in the art will appreciate that such descriptions are applicable to the present invention and can be easily adapted to it.

[000196] The proper dosage depends on various factors such as the type of disease being treated, the particular composition being used and the size and physiological condition of the patient. Therapeutically effective doses for the compounds described herein can be estimated initially from cell culture and animal models. For example, a dose can be formulated in animal models to achieve a circulating concentration range that initially takes into account the IC_{50} as determined in cell culture assays. The animal model data can be used to more accurately determine useful doses in humans.

[000197] Plasma half-life and biodistribution of the drug and metabolites in the plasma, tumors and major organs can also be determined to facilitate the selection of drugs most appropriate to inhibit a disorder. Such measurements can be carried out. For example, HPLC analysis can be performed on the plasma of animals treated with the drug and the location of radiolabeled compounds can be determined using detection methods such as X-ray, CAT scan and MRI. Compounds that show potent inhibitory activity in the screening assays, but have poor pharmacokinetic characteristics, can be optimized by altering the chemical structure and retesting. In this regard, compounds displaying good pharmacokinetic characteristics can be used as a model.

[000198] Toxicity studies can also be carried out by measuring the blood cell composition. For example, toxicity studies can be carried out in a suitable animal model as follows: 1) the

compound is administered to mice (an untreated control mouse should also be used); 2) blood samples are periodically obtained via the tail vein from one mouse in each treatment group; and 3) the samples are analyzed for red and white blood cell counts, blood cell composition and the percent of lymphocytes versus polymorphonuclear cells. A comparison of results for each dosing regime with the controls indicates if toxicity is present.

[000199] At the termination of each toxicity study, further studies can be carried out by sacrificing the animals (preferably, in accordance with the American Veterinary Medical Association guidelines Report of the American Veterinary Medical Assoc. Panel on Euthanasia, Journal of American Veterinary Medical Assoc., 202:229-249, 1993). Representative animals from each treatment group can then be examined by gross necropsy for immediate evidence of metastasis, unusual illness or toxicity. Gross abnormalities in tissue are noted and tissues are examined histologically. Compounds causing a reduction in body weight or blood components are less preferred, as are compounds having an adverse effect on major organs. In general, the greater the adverse effect the less preferred the compound.

[000200] For the treatment of many diseases, the expected daily dose of a hydrophobic pharmaceutical agent is between 1 to 500 mg/day, preferably 1 to 250 mg/day, and most preferably 1 to 50 mg/day. Drugs can be delivered less frequently provided plasma levels of the active moiety are sufficient to maintain therapeutic effectiveness. Plasma levels should reflect the potency of the drug. Generally, the more potent the compound the lower the plasma levels necessary to achieve efficacy.

[000201] As discussed above, it is well known that GPCRs are expressed in many different tissues and regions, including in the brain. nGPCR-x mRNA transcripts may found in many other tissues, including, but not limited to peripheral blood lymphocytes, pancreas, ovary, uterus, testis, salivary gland, kidney, adrenal gland, liver, bone marrow, prostate, fetal liver, colon, muscle, and fetal brain, and may be found in many other tissues. Within the brain, nGPCR-x mRNA transcripts may be found in many tissues, including, but not limited to, frontal lobe, hypothalamus, pons, cerebellum, cerebrum, caudate nucleus, and medulla.

[000202] Sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 will, as detailed above, enable screening the endogenous neurotransmitters/hormones/ligands which activate, agonize, or antagonize nGPCR-x and for compounds with potential utility in treating disorders including, but not limited to, thyroid disorders (*e.g.* thyreotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Chron's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain including schizophrenia, migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety,

generalized anxiety disorder, post-traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); sexual dysfunction, among others.

[000203] For example, nGPCR-x may be useful in the treatment of respiratory ailments such as asthma, where T cells are implicated by the disease. Contraction of airway smooth muscle is stimulated by thrombin. Cicala *et al* (1999) Br J Pharmacol 126:478-484. Additionally, in bronchiolitis obliterans, it has been noted that activation of thrombin receptors may be deleterious. Hauck *et al.* (1999) Am J Physiol 277:L22-L29. Furthermore, mast cells have also been shown to have thrombin receptors. Cirino *et al* (1996) J Exp Med 183:821-827. nGPCR-x may also be useful in remodeling of airway structure s in chronic pulmonary inflammation via stimulation of fibroblast procollagen synthesis. See, *e.g.*, Chambers *et al.* (1998) Biochem J 333:121-127; Trejo *et al.* (1996) J Biol Chem 271:21536-21541.

[000204] In another example, increased release of sCD40L and expression of CD40L by T cells after activation of thrombin receptors suggests that nGPCR-x may be useful in the treatment of unstable angina due to the role of T cells and inflammation. See Aukrust *et al.* (1999) Circulation 100:614-620.

[000205] A further example is the treatment of inflammatory diseases, such as psoriasis, inflammatory bowel disease, multiple sclerosis, rheumatoid arthritis, and thyroiditis. Due to the tissue expression profile of nGPCR-x, inhibition of thrombin receptors may be beneficial for these diseases. See, *e.g.*, Morris *et al.* (1996) Ann Rheum Dis55:841-843. In addition to T cells, NK cells and monocytes are also critical cell types which contribute to the pathogenesis of these diseases. See, *e.g.*, Naldini & Carney (1996) Cell Immunol 172:35-42; Hoffman & Cooper (1995) Blood Cells Mol Dis 21:156-167; Colotta *et al.* (1994) Am J Pathol 144:975-985.

[000206] Expression of nGPCR-x in bone marrow and spleen may suggest that it may play a role in the proliferation of hematopoietic progenitor cells. See DiCuccio *et al.* (1996) Exp Hematol 24:914-918.

[000207] As another example, nGPCR-x may be useful in the treatment of acute and/or traumatic brain injury. Astrocytes have been demonstrated to express thrombin receptors. Activation of thrombin receptors may be involved in astrogliosis following brain injury. Therefore, inhibition of receptor activity may be beneficial for limiting neuroinflammation. Scar formation mediated by astrocytes may also be limited by inhibiting thrombin receptors. See, *e.g.*, Pindon *et al.* (1998) *Eur J Biochem* 255:766-774; Uhl & Reiser. (1997) *Glia* 21:361-369; Grabham & Cunningham (1995) *J Neurochem* 64:583-591.

[000208] nGPCR-x receptor activation may mediate neuronal and astrocyte apoptosis and prevention of neurite outgrowth. Inhibition would be beneficial in both chronic and acute brain injury. See, *e.g.*, Donovan *et al.* (1997) *J Neurosci* 17:5316-5326; Turgeon *et al.* (1998) *J Neurosci* 18:6882-6891; Smith-Swintosky *et al.* (1997) *J Neurochem* 69:1890-1896; Gill *et al.* (1998) *Brain Res* 797:321-327; Suidan *et al.* (1996) *Semin Thromb Hemost* 22:125-133.

[000209] The attached Sequence Listing contains the sequences of the polynucleotides and polypeptides of the invention and is incorporated herein by reference in its entirety.

Methods of Screening Human Subjects

[000210] Thus in yet another embodiment, the invention provides genetic screening procedures that entail analyzing a person's genome -- in particular their alleles for the nGPCR-x of the invention -- to determine whether the individual possesses a genetic characteristic found in other individuals that are considered to be afflicted with, or at risk for, developing a mental disorder or disease of the brain that is suspected of having a hereditary component. For example, in one embodiment, the invention provides a method for determining a potential for developing a disorder affecting the brain in a human subject comprising the steps of analyzing the coding sequence of one or more nGPCR-x genes from the human subject; and determining development potential for the disorder in said human subject from the analyzing step.

[000211] More particularly, the invention provides a method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of: (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering the amino acid sequence, expression, or biological activity of at least one seven transmembrane receptor that is expressed in the brain, wherein the seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or an allelic variant thereof, and wherein the nucleic acid corresponds to the gene encoding the seven transmembrane receptor; and (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of allele in the nucleic acid correlates with an increased risk of developing the disorder.

[000212] By "human subject" is meant any human being, human embryo, or human fetus. It will be apparent that methods of the present invention will be of particular interest to individuals that have themselves been diagnosed with a disorder affecting the brain or have relatives that have been diagnosed with a disorder affecting the brain.

[000213] By "screening for an increased risk" is meant determination of whether a genetic variation exists in the human subject that correlates with a greater likelihood of developing a disorder affecting the brain than exists for the human population as a whole, or for a relevant racial or ethnic human sub-population to which the individual belongs. Both positive and negative determinations (i.e., determinations that a genetic predisposition marker is present or is absent) are intended to fall within the scope of screening methods of the invention. In preferred embodiments, the presence of a mutation altering the sequence or expression of at least one nGPCR-x seven transmembrane receptor allele in the nucleic acid is correlated with an increased risk of developing mental disorder, whereas the absence of such a mutation is reported as a negative determination.

[000214] The "assaying" step of the invention may involve any techniques available for analyzing nucleic acid to determine its characteristics, including but not limited to well-known techniques such as single-strand conformation polymorphism analysis (SSCP) [Orita *et al.*, *Proc. Natl. Acad. Sci. USA*, 86: 2766-2770 (1989)]; heteroduplex analysis [White *et al.*, *Genomics*, 12: 301-306 (1992)]; denaturing gradient gel electrophoresis analysis [Fischer *et al.*, *Proc. Natl. Acad. Sci. USA*, 80: 1579-1583 (1983)]; and Riesner *et al.*, *Electrophoresis*, 10: 377-389 (1989)]; DNA sequencing; RNase cleavage [Myers *et al.*, *Science*, 230: 1242-1246 (1985)]; chemical cleavage of mismatch techniques [Rowley *et al.*, *Genomics*, 30: 574-582 (1995)]; and Roberts *et al.*, *Nucl. Acids Res.*, 25: 3377-3378 (1997)]; restriction fragment length polymorphism analysis; single nucleotide primer extension analysis [Shumaker *et al.*, *Hum. Mutat.*, 7: 346-354 (1996)]; and Pastinen *et al.*, *Genome Res.*, 7: 606-614 (1997)]; 5' nuclease assays [Pease *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:5022-5026 (1994)]; DNA Microchip analysis [Ramsay, G., *Nature Biotechnology*, 16: 40-48 (1999)]; and Chee *et al.*, U.S. Patent No. 5,837,832]; and ligase chain reaction [Whiteley *et al.*, U.S. Patent No. 5,521,065]. [See generally, Schafer and Hawkins, *Nature Biotechnology*, 16: 33-39 (1998).] All of the foregoing documents are hereby incorporated by reference in their entirety.

[000215] Thus, in one preferred embodiment involving screening nGPCR-x sequences, for example, the assaying step comprises at least one procedure selected from the group consisting of: (a) determining a nucleotide sequence of at least one codon of at least one nGPCR-x allele of the human subject; (b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference

sequences; (c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and (d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

[000216] In a highly preferred embodiment, the assaying involves sequencing of nucleic acid to determine nucleotide sequence thereof, using any available sequencing technique. [See, e.g., Sanger *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 74: 5463-5467 (1977) (dideoxy chain termination method); Mirzabekov, *TIBTECH*, 12: 27-32 (1994) (sequencing by hybridization); Drmanac *et al.*, *Nature Biotechnology*, 16: 54-58 (1998); U.S. Patent No. 5,202,231; and *Science*, 260: 1649-1652 (1993) (sequencing by hybridization); Kieleczawa *et al.*, *Science*, 258: 1787-1791 (1992) (sequencing by primer walking); (Douglas *et al.*, *Biotechniques*, 14: 824-828 (1993) (Direct sequencing of PCR products); and Akane *et al.*, *Biotechniques* 16: 238-241 (1994); Maxam and Gilbert, *Meth. Enzymol.*, 65: 499-560 (1977) (chemical termination sequencing), all incorporated herein by reference.] The analysis may entail sequencing of the entire nGPCR gene genomic DNA sequence, or portions thereof; or sequencing of the entire seven transmembrane receptor coding sequence or portions thereof. In some circumstances, the analysis may involve a determination of whether an individual possesses a particular allelic variant, in which case sequencing of only a small portion of nucleic acid – enough to determine the sequence of a particular codon characterizing the allelic variant – is sufficient. This approach is appropriate, for example, when assaying to determine whether one family member inherited the same allelic variant that has been previously characterized for another family member, or, more generally, whether a person's genome contains an allelic variant that has been previously characterized and correlated with a mental disorder having a heritable component.

[000217] In another highly preferred embodiment, the assaying step comprises performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences. In a preferred embodiment, the hybridization involves a determination of whether nucleic acid derived from the human subject will hybridize with one or more oligonucleotides, wherein the oligonucleotides have nucleotide sequences that correspond identically to a portion of the nGPCR-x gene sequence taught herein, or that correspond identically except for one mismatch. The hybridization conditions are selected to differentiate between perfect sequence complementarity and imperfect matches differing by one or more bases. Such hybridization experiments thereby can provide single nucleotide polymorphism sequence information about

the nucleic acid from the human subject, by virtue of knowing the sequences of the oligonucleotides used in the experiments.

[000218] Several of the techniques outlined above involve an analysis wherein one performs a polynucleotide migration assay, *e.g.*, on a polyacrylamide electrophoresis gel (or in a capillary electrophoresis system), under denaturing or non-denaturing conditions. Nucleic acid derived from the human subject is subjected to gel electrophoresis, usually adjacent to (or co-loaded with) one or more reference nucleic acids, such as reference GPCR-x encoding sequences having a coding sequence identical to all or a portion of SEQ ID NO:1 to SEQ ID NO:58 (or identical except for one known polymorphism). The nucleic acid from the human subject and the reference sequence(s) are subjected to similar chemical or enzymatic treatments and then electrophoresed under conditions whereby the polynucleotides will show a differential migration pattern, unless they contain identical sequences. [See generally Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, New York: John Wiley & Sons, Inc. (1987-1999); and Sambrook *et al.*, (eds.), *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989), both incorporated herein by reference in their entirety.]

[000219] In the context of assaying, the term “nucleic acid of a human subject” is intended to include nucleic acid obtained directly from the human subject (*e.g.*, DNA or RNA obtained from a biological sample such as a blood, tissue, or other cell or fluid sample); and also nucleic acid derived from nucleic acid obtained directly from the human subject. By way of non-limiting examples, well known procedures exist for creating cDNA that is complementary to RNA derived from a biological sample from a human subject, and for amplifying (*e.g.*, via polymerase chain reaction (PCR)) DNA or RNA derived from a biological sample obtained from a human subject. Any such derived polynucleotide which retains relevant nucleotide sequence information of the human subject's own DNA/RNA is intended to fall within the definition of “nucleic acid of a human subject” for the purposes of the present invention.

[000220] In the context of assaying, the term “mutation” includes addition, deletion, and/or substitution of one or more nucleotides in the GPCR gene sequence (*e.g.*, as compared to the seven transmembrane receptor-encoding sequences set forth of SEQ ID NO:1 to SEQ ID NO:58, and other polymorphisms that occur in introns (where introns exist) and that are identifiable via sequencing, restriction fragment length polymorphism, or other techniques. The various activity examples provided herein permit determination of whether a mutation modulates activity of the relevant receptor in the presence or absence of various test substances.

[000221] In a related embodiment, the invention provides methods of screening a person's genotype with respect to the nGPCR-x of the invention, and correlating such genotypes with

diagnoses for disease or with predisposition for disease (for genetic counseling). For example, the invention provides a method of screening for an nGPCR-x hereditary mental disorder genotype in a human patient, comprising the steps of: (a) providing a biological sample comprising nucleic acid from the patient, the nucleic acid including sequences corresponding to said patient's nGPCR-x alleles; (b) analyzing the nucleic acid for the presence of a mutation or mutations; (c) determining a nGPCR-x genotype from the analyzing step; and (d) correlating the presence of a mutation in an nGPCR-x allele with a hereditary mental disorder genotype. In a preferred embodiment, the biological sample is a cell sample containing human cells that contain genomic DNA of the human subject. The analyzing can be performed analogously to the assaying described in preceding paragraphs. For example, the analyzing comprises sequencing a portion of the nucleic acid (e.g., DNA or RNA), the portion comprising at least one codon of the nGPCR-x alleles.

[000222] Although more time consuming and expensive than methods involving nucleic acid analysis, the invention also may be practiced by assaying one or more proteins of a human subject to determine the presence or absence of an amino acid sequence variation in GPCR protein from the human subject. Such protein analyses may be performed, *e.g.*, by fragmenting GPCR protein via chemical or enzymatic methods and sequencing the resultant peptides; or by Western analyses using an antibody having specificity for a particular allelic variant of the GPCR.

[000223] The invention also provides materials that are useful for performing methods of the invention. For example, the present invention provides oligonucleotides useful as probes in the many analyzing techniques described above. In general, such oligonucleotide probes comprise 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides that have a sequence that is identical, or exactly complementary, to a portion of a human GPCR gene sequence taught herein (or allelic variant thereof), or that is identical or exactly complementary except for one nucleotide substitution. In a preferred embodiment, the oligonucleotides have a sequence that corresponds in the foregoing manner to a human GPCR coding sequence taught herein, and in particular, the coding sequences set forth in SEQ ID NO:1 to SEQ ID NO:58. In one variation, an oligonucleotide probe of the invention is purified and isolated. In another variation, the oligonucleotide probe is labeled, *e.g.*, with a radioisotope, chromophore, or fluorophore. In yet another variation, the probe is covalently attached to a solid support. [See generally Ausubel *et al.* and Sambrook *et al.*, *supra*.]

[000224] In a related embodiment, the invention provides kits comprising reagents that are useful for practicing methods of the invention. For example, the invention provides a kit for screening

a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association: (a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-x seven transmembrane receptor gene, the oligonucleotide comprising 6-50 nucleotides that have a sequence that is identical or exactly complementary to a portion of a human nGPCR-x gene sequence or nGPCR-x coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and (b) a media packaged with the oligonucleotide containing information identifying polymorphisms identifiable with the probe that correlate with mental disorder or a genetic predisposition therefor. Exemplary information-containing media include printed paper package inserts or packaging labels; and magnetic and optical storage media that are readable by computers or machines used by practitioners who perform genetic screening and counseling services. The practitioner uses the information provided in the media to correlate the results of the analysis with the oligonucleotide with a diagnosis. In a preferred variation, the oligonucleotide is labeled.

[000225] In still another embodiment, the invention provides methods of identifying those allelic variants of GPCRs of the invention that correlate with mental disorders. For example, the invention provides a method of identifying a seven transmembrane allelic variant that correlates with a mental disorder, comprising steps of: (a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny; (b) analyzing the nucleic acid for the presence of a mutation or mutations in at least one seven transmembrane receptor that is expressed in the brain, wherein the at least one seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding the at least one seven transmembrane receptor; (c) determining a genotype for the patient for the at least one seven transmembrane receptor from said analyzing step; and (d) identifying an allelic variant that correlates with the mental disorder from the determining step. To expedite this process, it may be desirable to perform linkage studies in the patients (and possibly their families) to correlate chromosomal markers with disease states. The chromosomal localization data provided herein facilitates identifying an involved nGPCR with a chromosomal marker.

[000226] The foregoing method can be performed to correlate the nGPCR-x of the invention to a number of disorders having hereditary components that are causative or that predispose persons to the disorder. For example, in one preferred variation, the disorder is a mental disorder.

[000227] Also contemplated as part of the invention are polynucleotides that comprise the allelic variant sequences identified by such methods, and polypeptides encoded by the allelic variant

sequences, and oligonucleotide and oligopeptide fragments thereof that embody the mutations that have been identified. Such materials are useful in *in vitro* cell-free and cell-based assays for identifying lead compounds and therapeutics for treatment of the disorders. For example, the variants are used in activity assays, binding assays, and assays to screen for activity modulators described herein. In one preferred embodiment, the invention provides a purified and isolated polynucleotide comprising a nucleotide sequence encoding a nGPCR-x receptor allelic variant identified according to the methods described above; and an oligonucleotide that comprises the sequences that differentiate the allelic variant from the nGPCR-x sequences set forth in SEQ ID NO:1 to SEQ ID NO:58. The invention also provides a vector comprising the polynucleotide (preferably an expression vector); and a host cell transformed or transfected with the polynucleotide or vector. The invention also provides an isolated cell line that is expressing the allelic variant nGPCR-x polypeptide; purified cell membranes from such cells; purified polypeptide; and synthetic peptides that embody the allelic variation amino acid sequence. In one particular embodiment, the invention provides a purified polynucleotide comprising a nucleotide sequence encoding a nGPCR-x seven transmembrane receptor protein of a human that is affected with a mental disorder; wherein said polynucleotide hybridizes to the complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 under the following hybridization conditions: (a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS; and wherein the polynucleotide encodes a nGPCR-x amino acid sequence that differs from a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, by at least one residue.

[000228] An exemplary assay for using the allelic variants is a method for identifying a modulator of nGPCR-x biological activity, comprising the steps of: (a) contacting a cell expressing the allelic variant in the presence and in the absence of a putative modulator compound; (b) measuring nGPCR-x biological activity in the cell; and (c) identifying a putative modulator compound in view of decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator.

[000229] Additional features of the invention will be apparent from the following Examples. Examples 1 and 2 are actual while the remaining Examples are prophetic. Additional features and variations of the invention will be apparent to those skilled in the art from the entirety of this application, including the detailed description, and all such features are intended as aspects of the invention. Likewise, features of the invention described herein can be re-combined into additional embodiments that also are intended as aspects of the invention, irrespective of

whether the combination of features is specifically mentioned above as an aspect or embodiment of the invention. Also, only such limitations which are described herein as critical to the invention should be viewed as such; variations of the invention lacking limitations which have not been described herein as critical are intended as aspects of the invention.

EXAMPLES

EXAMPLE 1: IDENTIFICATION OF nGPCR-X

A. Database search

- [000230] The Celera database was searched using known GPCR receptors as query sequences to find patterns suggestive of novel G protein-coupled receptors. Positive hits were further analyzed with the GCG program BLAST to determine which ones were the most likely candidates to encode G protein-coupled receptors, using the standard (default) alignment produced by BLAST as a guide.
- [000231] Briefly, the BLAST algorithm, which stands for Basic Local Alignment Search Tool is suitable for determining sequence similarity (Altschul *et al.*, J. Mol. Biol., 1990, 215, 403-410, which is incorporated herein by reference in its entirety). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension for the word hits in each direction are halted when: 1) the cumulative alignment score falls off by the quantity X from its maximum achieved value; 2) the cumulative score goes to zero or below, due to the accumulation of one or more negative scoring residue alignments; or 3) the end of either sequence is reached. The Blast algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The Blast program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff *et al.*, Proc. Natl. Acad. Sci. USA, 1992, 89, 10915-10919, which is incorporated herein by reference in its entirety) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.
- [000232] The BLAST algorithm (Karlin *et al.*, Proc. Natl. Acad. Sci. USA, 1993, 90, 5873-5787, which is incorporated herein by reference in its entirety) and Gapped BLAST perform a statistical analysis of the similarity between two sequences. One measure of similarity provided

by the BLAST algorithm is the smallest sum probability ($P(N)$), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a GPCR gene or cDNA if the smallest sum probability in comparison of the test nucleic acid to a GPCR nucleic acid is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

- [000233] Homology searches are performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.
- [000234] Multiple query sequences may have a significant alignment to the same genomic region, although each alignment may not cover exactly the same DNA region. A procedure is used to determine the region of maximum common overlap between the alignments from several query sequences. This region is called the consensus DNA region. The procedure for determining this consensus involves the automatic parsing of the BLAST output files using the program MSPcrunch to produce a tabular report. From this tabular report the start and end of each alignment in the genomic DNA is extracted. This information is used by a PERL script to derive the maximum common overlap. These regions are reported in the form of a unique sequence identifier, a start and the end position in the sequence. The sequences defined by these regions were extracted from the original genomic sequence file using the program fetchdb.
- [000235] The consensus regions are assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCRs. These sequences were then submitted for further sequence analysis.
- [000236] Further sequence analysis involves the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCR's.
- [000237] nGPCR-cDNAs were sequenced directly using an ABI377 fluorescence-based sequencer (Perkin-Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FS™ polymerase. Each ABI cycle sequencing reaction contained about 0.5 µg of plasmid DNA. Cycle-sequencing was performed using an initial denaturation at 98°C for 1 minute, followed by 50 cycles using the following parameters: 98°C for 30 seconds, annealing at 50°C for 30 seconds, and extension at 60°C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using Centriflex™ gel filtration cartridges (Advanced Genetic

Technologies Corp., Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which is then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 5 µl of a DNA loading solution (83% deionized formamide, 8.3mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90°C for three minutes and loaded into the gel sample wells for sequence analysis using the ABI377 sequencer. Sequence analysis was performed by importing ABI377 files into the Sequencer program (Gene Codes, Ann Arbor, MI). Generally, sequence reads of 700 bp were obtained. Potential sequencing errors were minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers annealing at different locations until all sequencing ambiguities were removed.

[000238] The following Table 5 contains the sequences of the polynucleotides and polypeptides of the invention. The transmembrane domains within the polypeptide sequence are identified by underlining.

Table 5
<p>The following DNA sequence Seq-86 <SEQ ID NO. 1> was identified in <i>H. sapiens</i>:</p> <p>ACACAGTGTGCACACAGTGCAGGGACATACCCCTTCCCCAACTGCCTGGCCTGCACACTTGGCATTTC AGTATTTCTAGGAAGTGATGGCTCTGTGCATCCTGAGCCAATCCAGCTCCGAGCCTCCAAGGCATCCTGGT GATGGGCAGCTGGAAGCTCTGCCTCTGAGGCCTTCACACACCCACCTTCGGTCAAACCTGCTTCTGCTGAG GAACTTGGTGTGTCTTCTCTGGGCAGGAGGTCACATTTGAGAGCACAGGAGCAGTGCCTGCCCCCGGG AATGTGGCTCTGGGTAGAATTGCAGGCTCAGGGGTTTTGGGCAGGAGAGCACCAACCGTGCCACACCCACA CAGACACGGTCACTGGGGCCCTGCAGCAGGGACGACCGCACTTCCCAAAGGGCTGGGAAGCCATGTCCAGA GGAGGCCATGCTCTAGCTCCCTTGGGCAGGGCTGGCTGCAAGGAGGGTGAAGTTGGGCATCTTGAACCCAG AGAAGTAGAGGACTCAGCACCAGCACAAACAGCTCGGCGCATTAAATACACATTCCTCTCCCACTTCTCCCC AAGCCTGAAAAAACCTCAAACAGCCTCTTTGCAGCTCCCTGAGGTGATGACTCACGAACCATGCTCGGGG CAGGGGAAAAGAAAAGCATCCG</p> <p>The following amino acid sequence <SEQ ID NO. 59> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 1:</p> <p>DAFLFPCPEHGSVMTSGSCKEAGLRFFQAWGEVGEECVLMRRAGCAGAESSTSLGSRCPSPSPALPKG ARAWPPLDMASQPFKCGRPCCRAPVTVSVVWHGWCSPAQNPAACNSTQSHIPGGQALLCSQMPPAQKED TPSSSAEASLTGGGCVKASEAELPAAHHQDALEARSWIGSGCTEPLPRNTGNAKCAGQAVGEGGMSLHVC AHC</p>
<p>The following DNA sequence Seq-87 SEQ ID NO. 2> was identified in <i>H. sapiens</i>:</p> <p>CTAAAGGAGGAATAGATGTCTTTAAGAAGAAATGAAAAATAAAGTAAATGTGAAAATTTCCCTTACTTAT TTCCAAACAAGTGCTCCTCCAAAAAATGCAATAATTAAGTTTCTGAAATGGTGAACATATCAGATTAGT AGACATATGGCAGSAGCAGCAATGAGCAGATCAAGTTGAAGTCTAGTATTACCAATCTGTTAATGTTGA CAGGAAGACTCATTGTGACTGTTCTTTTATATCAATAAATGAGTGGATTTCAACTACTCTAAATAGGAAT GCTAAAAGCAGCACTGCTAAAAGTGCATATCAAACCAATAATTTCTGATGCTGTTTGGGTATATCCTACA AACATTTGTAGGACAACAACCTCAGAAGGGAAAAAATATCTTATGCCTTTGAGGTCTGTACTGAATGCTAA TGCATTTGTATATGATGGGTTTAAATACAGAACTGAGAATAAATTAATTTTACGAGCTGCACTCTAGACCTA TAAATCGCTCTGAGTACTACAAATCCATACAAAGGAAGAACAGCTGGATAATTTACACCACAGTATTTG TCAAAAAAAGCTGAAAAATACAGAACCTGATTTTGTCCCTTTTTCGAGTA</p>

The following amino acid sequence <SEQ ID NO. 60> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 2:

LEKGTSGSVSFAFFFFQILVVIQLFFLCMDFFVLRAIYRSRVQLLKVIYSQFCIKPIIYKICISIQYRP
QRHKIFFSLLSCPTNVCRITYQNSIRKLLVYALLAVLLLAFLFRVVEIHSFIDIKGTVKMSLPVNINRLVI
LGLQLDLLICCSCHMSTNLICSPFQKLNLYLHFFGGALVWKVREIFTTLFFHFFLKTSIPPL

The following DNA sequence Seq-88 <SEQ ID NO. 3> was identified in *H. sapiens*:

AGGGGCCCTCCAGCACTGGTCTTGAAGGGGTGACAGGGTCTGGGGTCTGACTCCACCTCCACCACTTCCC
ACCTGAGGGCCCTGGAATGAATCCTTTCTGGATCTGAGCTGCCACATCATCAGTGAAAATGACACCTATA
TGGGACTTCAGTGAGAACACAAATGCAACGTTCTGCCACGGAACAACCATGTACTCACTGGGAGCATTG
AGAGTAGATCCACACTGATTGACACAGGGACTCCAGGCCTGACCCATGATATGTACTGGATACATGGCCAT
GAGTGCTCCACAG

The following amino acid sequence <SEQ ID NO. 61> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 3:

VEHSWPCIQYISWVRPGVPVSISVDLLSMLPVSTWVVPWQERCICVLTEVPYRCHFHCSSDPGKDSFQGP
QVSGGGGSQTPDPVTPSRPVLEGP

The following DNA sequence Seq-89 <SEQ ID NO. 4> was identified in *H. sapiens*:

ACACCAAATACTGCTTTGCTGCCTTAGGCTTCAGCACATTAGCATGGCTTCCCTCCCTTGGCATGGTAACTG
TAGCTGAACTTGGAGGGTTTGTATTACCCATTATAATTATTACTTATTTACATGGAAAACAAGAAAATCT
TTATGGGAATCCAAGTTCCTCCCTAGGAATACCAAGAGAGAGGAAAAGGCTTTGAGGATGGTCTGATGTG
TGAAGTGGTGTTCATTGTGTGTTCACTCCTTACCACCTCAACTCCCATTTTATGATGGTGAAGGAAC
ATGTCTTTTGAAGTCTCTTTTATAAAGATCATTCTCTGTTTCCACATTATTTCCCTGTGTCTTGCAAAAT
CTGAATTGTTGCTTGTATCCAGTTGTATATTATTTATGACCTCAAAATTTTCATGATCAATTTTCAGATCA
TGGCAGCTTGGTTCTTCAGTCATGTATGAGATGTAATAACAGTACCTTAGAAATTCATCAGAGGAAGGGAG
GATCTTCAAATATCTCTCTTGAATGTTTGAAGATTCCAAGACAATATAATCAAATAATTAAGTAGAAAA
ATCGATATGCTCTATTAGTGTATCTATGTCACTTTGAAGATTTTCTTTTTTTTTTTCTTTTTTTTTTATT
ATACTTTAAG

The following amino acid sequence <SEQ ID NO. 62> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 4:

HQILLCCRLLOHISMASSLGMVTVAELGGFVLPIIIITYFTWKTRKSLWEFQVPPRNTKERKKALRMVLMC
EVVFIVCFPTYHLNFPFFMMVKEHVFLNCSFIKIILCFHIIISLCLANLNCCLDPVVYFMTSKFHDQFSDH
GSLVLQSCMRNNSLTLEIHRQKGGSSNYLSMFERFDQNIILKTRKIDMLYCIYVTLKIFLFFFSFFLLYFK

The following DNA sequence Seq-90 <SEQ ID NO. 5> was identified in *H. sapiens*:

AATGCTACTGCTCCTGCATATAATAGCTGCTTTGAAGTGTGTTGCTGCTATGTCCAAAACATAGACTCTTT
CAAAAGCACTTTCTGTTGCTCCTTTTTCTCTTGCATAGGAGTCACATTTTCTGTCTCTTCACATATTTT
ATATTTATTTTTGTTGAAAACAGACATTTTAGATAATGTGTTGTAGCAGTCCAGATACTGATTCTCTCCC
CCAGGAGCTGTTGCTCTTCTTACTTGTATATGTGTTTAGTGACTGGCTGGACTATTTAATAATGTGTAT
TTCCCTGTAGTATATAACCATCTTTTATACTAATGTTACTTTTCCGATAGTGCAGCCTTGGGCATGGACAGA
GTTATCTCTGGGATGACAGTAACCTTTAATAGGGCTCTCTATGACTATCTCTTCCCTGATGTCCTGTAA
GCTATCTGCATCTCTTGGTATCACACCTAGCCTTTGACTTCCACTAATGTTTGATCATTGCCTCACTGTT
TTTGGCAGTGCCCTAAGGCATAAAGTGTTCCACAGTCTGATATAATTAAATTCAGATTCTTACAAGAGTGG
TCTTTGAGGCCAGTCCTTGAGGTTTGTGCTGACTCTGGGAGGGCTCAAATGTTTCCCT

The following amino acid sequence <SEQ ID NO. 63> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 5:

CYCSCILLSVCLLCPKHRLFQKHLLSPFSLAESHSVSSSHISYLFLLKTRHFRVCVAVQILILSPRSCCL
SYLYMCLVTLWLDYFNNVYFPVVYTFYTNVTFPIVQFPAWTELSWDDSNFGSLLSLSLMSLLSYLHLLVSH
LAFDFHLFDHCLTVFGSALRHKVFHSLILNSDSYSGSLRQSLRFVLTGLGLKCFP

The following DNA sequence Seq-91 <SEQ ID NO. 6> was identified in *H. sapiens*:

CCTCACATCCCCCTTCCCTCAAACCTGGCAACCCAACTGTTCTTGACAGCCTCCTTTGGCATTTCCTC
ATTTTGGTGTGAGATCTCACAGCAGAATTTCTTACCTATTATATACCAGTGCCCTCAGTGTGAAGTTCCGGT
TTAACTTCTGTATTACCAGAGCCCACTATCTGCCCCAATAATACCTCCCCCAATTACAAACACACAAG
CATTCCTCCTACAGCTTTGGGCCTCCTATCTGAGTCCTTCAGGAAAGAAGTGTGTGTAAGTCCCTTGGC
AGTGAGTGTAGACTTGGTCCAAGGAAGATGAGCACCAGTCAGGGCAGCTGGGCCCTCTTCTCTCCCTGGCC
ATCAGCAAATCAGCACTGCCATCGATGCCAGGCAATGGGAGCG

The following amino acid sequence <SEQ ID NO. 64> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 6:

PHIPFPSNPGNPKLFLTASFGISFQWQISQONFLPIIYQCLSVKFRFNFLLPRAHYLAPIIPSPNSQTHK
HSLQLWASYLSPSGKKCCVTPPLAVSVDLVQGRAPVRAAGPSSLPGHQQISTAHRCPNGS

The following DNA sequence Seq-92 <SEQ ID NO. 7> was identified in *H. sapiens*:

ATTACTATTTTCAACCTCTTTTACTCCAGGGACTTCTATGCACCCTCTCCCTCACTCCCCCTCAATTTG
TTCTCATATATCCCCATGACCCCCAGTTTTATAACCACTGTGAGGAGCCCAAAGCTGCCATTTCATTCAT
TCCATTAGCATGACTCTTCATGTACTTTGGGGTCTTCAGTCTCTCCCTTCTCCTAATTTCCAGGGTTCCA
TTCTGCTTCTGCTGGCTTCCCTACAAAGCCTGCAACATCATAAGCCATTTTCAGGAAAGAGCTTGATCATCT
TTTGATGAACCTGCAATTCATGACTCACTGCCTTACCTGTCTTTGGCTCTGCATGTCCCCAGTTTCCGTT
TCTTTCTCTGAAAGAGAGATTGCCCAAGAGTCTGCACATCAGCATTACTAGAAATGCATGCAGACCAGC
TTCAGCTGCTTGCCCACTCTTTAAAAAATGAGTAAACAATTTTCTAAAGGGGAAAAAATCTCTTACCTCC
TCACACCAACTATTTGCATAATTCAGTGACCTTTTATAACCGTGCCATTGTATAAGCA

The following amino acid sequence <SEQ ID NO. 65> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 7:

ITIFQPLLLQGLLCTLSLNSPISCSHNPHDPQFYNTTVRSPKLPFIHFHITIFQPLLLQGLLCTLSLNSHD
SSCTLGSSVSPLLLISRVPFCFCLWLPYKACNIISHFRKELDHLMLNPAFMTHCLTCLWLCMSPSFRFLWK
ERLPKSPARHQHYKCMQTSFSCPLTKMSKQFSKGEKISSPPHTNYLHNSVTYKPCHCIS

The following DNA sequence Seq-93 <SEQ ID NO. 8> was identified in *H. sapiens*:

CACCGTCCTCATCATGATCGTCTTCGTATCTGCTGCTGGGGGCCCTACTGCTTCTGGTGCTGCTGGCCG
CCGCCCGGCGAGGCCAGACCATGCAGGCCCTCGCTCCTCAGCGTGGTGGCCGTCTGGCTGACCTGGGGC
AATGGGGCCATCAACCTGTCTATCTACGCCATCCGCAATCCCAACATTTGATGCTCCTAGGGCGCAACCG
CGAGGAGGGCTACCGGACTAGGAATGTGGACGCTTTCTGCCCAGCCAGGGCCCGGGTCTGCAAGCCAGAA
GCCGCACTGCGCTTCGAAACCGCTATGCCAACCGGCTGGGGGCCCTGCAACAGGATGTCTCTTCCAACCG
GCCAGCGGAGTGGCAGGGGACGTGGCCATGTGGGCCCGCAAAATCCAGTTGTACTTTTCTGCCGAGAGGG
ACCACCAGAGCCGGTGACGGCAGTGACCAACAGCCTAAATCCGAAGCTGGGGATACCAGCCTCTAAGACG
GTTGGAATGGCCAGCTTATGAAGGCAATTTCCACTCGCATTATTTAATGATGGAAGATTCTGGGGGAGAG
TTGTGGATTTTCATAAAGCCAAACATTTAAAGCTAGAGACGGGGGAGGCTTACCACTTTCCCCAAACAACAT
AAAAGACAATGTCCCTTCTTTCAAAAAGTGC

The following amino acid sequence <SEQ ID NO. 66> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 8:

TVLIMIVFVICWGPYCFVLVLLAAARQAQTMQAPSLLSVVAVWLTWANGAINPVIYAIRNPNISMLLRNR
EEGYRTRNVDAFLPSQGPGLQARSRLRNRYANRLGACNRMSSSNPASGVAGDVAMWARKNPVVLFCREG
PPEPVTAVTKQPKSEAGDTSLDGWNQQLMKANFHSYLMEDSGGELWISSQTFKARDGGGLPLSPNNIKD
NVPSFKKC

The following DNA sequence nGPCR-93 <SEQ ID NO. 9> was identified in *H. sapiens*:

CGCTCCTGCGTAAACACGCGGTTCCTCGGCAACGCTGGAACCCACGTCAAAGGCTCCGCCAGGTCCCCAG
CGACCGCCACCCCTCCGGCCGAGCCAGCTCCCCGCGGCGGCCGCTAGCCCCCGGCCCGAGCCACCACTC
CGACCTAGCGGCCCGCCGCCCGGTGCGGGATGAGGAGATCCGCGGCCGCCACTGGGGCCCCATGGAGGAGC

CGCAGCCGCCCGCCACCAGCGAGCATGGCCTTACTGGGCAGCCAGCACTCCGGCGCCCCCTCCGCGGCC
 GGCCACCTGGCGGGACTTCTCCGCGGCCACGGCGGCGGTGCTCTCCTTACGACCCGTGGCGACCGCGGC
 GCTGGGGAACCTGAGCGACGCAAGCGGAGGCGGCACAGCTGCCGTCCCGGTGGCGGCGCCCTTGGCGGGT
 CCGGGGACAGCGCGGAGGCGGGGGCGGCGGTGAGGCGGCCGCTAGCGACGGAGGCGGCGCGCTGCTGTCTG
 CACGGAGCTGCAGTGGCGGCCAGGCGCTCGTCTCTGCTCATCTTCTGCTGTCTAGCCTTGGCAACTG
 CGCGGTGATGGGGGTGATTGTGAAGCACCGCAGCTCCGACCCGTACCAACGCCTTCATCTGTCTGCTGT
 CCTATCGGATCTGCTACGGCGCTGCTCTGCTGCCGCGCCCTTCTGGACCTCTTCACTCCGCGCGGG
 GGTTTCGGCGCTGCCGCGCGCGGGGCCCTGGCGCGGCTTCTGCGCGCCAGCCGCTTCTTACGCTCGTG
 CTTCCGGCATCGTGTCCACGCTCAGCGTGGCGCTCATCTCGTTGGACCGTTACTGCGCTATCGTGGCGCCG
 CGCGGAGAGATCGGCGCGCGCGCGCGCTGCAGCTGCTGGCGGGCGCCTGGCTGACGGCCCTGGGCTTC
 TCCTTGCCCTGGGAGCTGCTCGGGGCGCCCCGGGAACCTCGCGGCGCGCGAGAGCTTCCACGGCTGCCCTA
 CCGGACCTCCCCGGACCCCGCGCAGCTGGGCGCGGCCCTTACGCGTGGGGCTGGTGGTGGCTGTACTCTGC
 TGCCCTTCTGCTCATGTGCTTCTGCCACTACCACATCTGCAAGACGGTGCCTGTGCGACGTGCGCGTG
 CGCGCGGTGAACACCTACGCGCGCGTGTGCGCTTCTTACGCGAGGTGCGCACGGCCACCACCGTCTCAT
 CATGATCGTCTTCGTCTGCTGCTGGGGGCCCTACTGCTTCTTGGTGTGCTGGCCGCGCGCGCGGACG
 CACAGACCATGACGGCCCCCTCGTCTCCTCAGCGTGGTGGCGCTGGCTGACCTGGGCGCAATGGGGCCATC
 AACCTGTCTATCTACGCCATCCGAATCCCAACATTTGATGCTCCTAGGCGCAACCGCGAGGAGGGCTA
 CCGGACTAGGAATGTGGACGCTTCTTCCGCGCGCGCGGCTGCAAGCCAGAACCCGCGAGTCCGCG
 TTCGAACCGCTATGCCAACCGCTGGGGGCGCTGCAACAGGATGCTCTTCCAACCGCGCGAGGAGTG
 GCAGGGGACGTGGCCATGTGGGGCCGCAAAATCCAGTTGTACTTTTCTGCCGAGAGGGACACCAGAGCC
 GTGACGCGCAGTGACCAACAGCCTAAATCCGAAGCTGGGGATACCAGCCTCTAAGACGGTTGGAATGGCC
 AGCTTATGAAGGCAATTTCCACTGCGATTATTTAATGATGGAAGATTCTGGGGGAGAGTTGTGGATTTC
 TAAAGCCAAACATTTAAAGCTAGAGACGGGGGAGGCTTACCATTTCCTCCCAACAACATAAAGACAATGT
 CCCTTCTTCAAAAG

The following amino acid sequence <SEQ ID NO. 67> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO.9:

LEPTSKAPPGPQRPPPLRSPAPRGRPPAPSHHSDLAAAAPGAGGDP RPPLGPMEEPQPRPPASMAILLG
 SQHSGAPSAAGPPPGTSSAATAAVLSFSTVATAALGNLSDASGGGTAAAPGGGLGGSGAAREAGAAVRRP
 LATEAAPLLSHGAAVAQALVLLIFLLSSIGNCAVMGVIVKHRQLRTVTNAFILSLSLSDLLTALLCLPA
AFILDLFTPPGGSAPAAAAGPWRGFCCAASRFFSSCGIVSTLSVALISLDRYCAIVRPPREKIGRRRALQLLA
GAWLTALGFSLPWELLGAPRELAQAQSFHGLYRTSPDPAQLGAAFSVGLVVACYLLPFLIMCFHYHICK
 TVRLSDVRVRPVNTYARVLRFFSEVRTATTVLIMIVFVICCWGPYCFLVLLAAARQAQTMAPSLLSVVAV
WLTWANGAINPVYAIRNPNISMLLGRNREEGYRTNVDAFLPSQGPGLQARSRLRNRYANRLGACNRM
 SSSNPASGVAGDVAMWARKNPVVLFCREGPEPVTAVTQPKSEAGDTSLDGWNQMLKANFHSYLMMED
 SGGELWISSQTFKARDGGGLPLSPNNI

The following DNA sequence Seq-2588 <SEQ ID NO. 10> was identified in *H. sapiens*:

TCTCAAAAAATAATAAAACCACTGTACATCAACAAGGCCCTTGGGGGACAGCTGGGGCATAAGTAGGTG
 TCAGCCATACATCAGAGCAGTGTGCCTGCCCTGAGCTGCTTGGGGTTGACCAGCCTGGTGTCCAGAAATGC
 CTGCTGGAGGGAGTCTGGTACAGGAAACCTTGTGCTCTTAGAAGGTCTCCTGAGAGGCCCTGCAAAGCCA
 GAGTCCCTCTTAGCAGCTCAGATCAGTGTATCAAAGTATAGCTCGGGGATTGCTGCCAGCATACAAACCTT
 TACTGGTCTGCAGCGAGATAAGTACAGAAATTGAAAGTAAGCATTTAGAACTTTTATAACAATTTTACA
 AGGTCTTGTCAAATGTTATTAACAAAGCTGAGGCTGGAATTTACCTTTTTCATTTCGTTTTTTTCAAT
 TTAACAAATTTAGTAAATATAGGTAATATAAATGTACCATTTTAGCCATTTTGGAGCGTACAATTTAG
 TAGCAGTAAGTGCTTTCACAATATTGTGTAACCACTAGTATTATATAGTATATATTTTAAATTTTACAG
 AAGTATTAAGTTAGCAGCAGATTAAACATTTTTTCTTAAATTGAGCTTGAGAAGCGCTGGC

The following amino acid sequence <SEQ ID NO. 68> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 10:

ASASQAQFKKMFNLLTYFCKILKIYTIYWLHNIVKALTATKLYAQKWLKWIYIYIYILLQFVIEKNEMK
 KVKEFQQLCFNNIQDLVKLLKFLNAYFQFLYLSRCRPVKVCMIAIPELYFDSTDLSCGLWLCRASQETF
 EHKVSCSTTPSSRHFWTPGWSTPSSSQAHCSVDWLTPYAPAVPQGPCCTVVFYFLR

The following DNA sequence Seq-2589 SEQ ID NO. 11> was identified in *H. sapiens*:

AGAGAGCAGATTGCCCTGTGTAGGTACAGTCTGGGTTCTTTCTAGTCCAGAGTAGGGAAGAAACAGGAAAG
 AGGGCTGGTGTGAAGGACCTTCAGCCACGAGAAGGGCTGTGTACCATGTAGCCCTCTGGGGAGGCACAAA
 AAGGCTCACCATTTTCTGAAAATGACTAGACTGCAGGATCCAGTGAGTGTGACTATTGCATTATGACCT

TATCCACAGGGCCTCACAAGGTGCCTGACATGCAGTAGGCTCCAGATGCATATTTATTATAAAGTGAATAG
 TCCTTAAGCTGCAGGGTCCCTTCTATTGCAATCTAAGAAATAGTCACTTTTATGCCTAATTTGTATTG
 CAGTTTATAAGTTTATAAGAGGGTCTCCCAAATAGTATAAACTTCAAGCCCCACAAAACCTATGTTTGC
 CTCCCATAGGCATGCAATAAATGTTCTGGATCTAATGAGTAACAAGAAAAGAAGGAACAAAACCTAAC
 CCCTCCCTACCCAAACCAAGTGGCAACCGGGGAGGATCAAATTCAACCTTGATCAGTCAGAGGCAGCATTC
 CTAATATTATCCCAAGCAGCAATAGACAATGATTACCTCAATTAATTCAGCCAGTTAAAGCTTAGTTCT
 TACTTGCCAACCGAAGGCTTGAAGGCAAAATGTGTTAAGCCTC

The following amino acid sequence <SEQ ID NO. 69> is the predicted
 amino acid sequence derived from the DNA sequence of SEQ ID NO.11:

RLKHLPSSRLRLASKNAFNWLNLRRIIVYCCGLGIECLLIKVEFDPPRLPLVWVGEGLGFCSEFFLLLR
 TNYICMPMGKKHRCFASLYLGDPLIKLIKLIQNAKFLRMQIEGLQLKDYSLYNKYAS
 GAYCMSGTLGPVDKVMNAIVTLTWILQSSHFKQMVSLFVPPQRTWYTAALLVAEGPSTPALFPVSSLLWTR
 KNPDLTYTGQSAL

The following DNA sequence Seq-2591 <SEQ ID NO. 12> was identified in
H. sapiens:

TTCAGGCAGATGTCAGTTAAAACTTACCTCTGCACACTGCAAAACTGTATAGCCCTGAACAGATACTTT
 TCTTGAGCATAGTTCCTTTGTCTCTAAGCAGGCATAATTGCCAATGTGGGATGATATTAGAAATCTGA
 ACTGATGTTTATTCTCTAGGGTCTTCTCATTGAGCTGGGATTGGAGATGTCTAGTGTCTCAGAGCAGCA
 ATAAGAAAACAGAAACCTCTCCAGCTTCTGACATCCAAATGTCAAGCTCTTAGGAGAAGAATGGAAAGTC
 CTCAGAAATGCAATAGCTTTGGCAGAAATAGCTGATGAAGACCACCTCTCCCCCTCCAGAAAGGCATTG
 GTTCCCATTCATGGAAAAGGAATGTAGAGAGAGATTAGATAATAGTACATCCATAAGGTTCTCTGGAATC
 TGCATCTGAGGAAGAGGGGCGTCAGAGACCCAGCTGTTATCTATAATCCCTCCT

The following amino acid sequence <SEQ ID NO. 70> is the predicted
 amino acid sequence derived from the DNA sequence of SEQ ID NO. 12:

EGLITAGVSDAPLQMQIPGTLWMYYYLISLYIPFSMNGEPMFWRGERWSSSAILPKLFAFLEDFPFSE
 LDIWMSEAGRGFCLIALRHTSPIPAQMRRPLENKHQFRFLNIPTLAIMPALETKELCRSKRVSVQGYTV
 FAVCRGKFLTDICL

The following DNA sequence Seq-2592 <SEQ ID NO. 13> was identified in
H. sapiens:

GCCTAGGGCAAAGTCAAGACATACGGGTGTCCAGTTCTAGCTTTGCAACTAACTGGTTATATATTTTAA
 GTTACAGTCACTCTGCGCCAGTTTCTCATTTTTAATAGAGTGGGTTAGAACTAGATAAATACTTTTCATT
 TGTCAAGCTCTAAATCTGACTTCAGGAAAAACCATAAGGCACTGGAGGTTTATTCATAGGTTTTCTGC
 TGACCCCGTCCCTCTCTGTTCTTCAACCACCACAAGACAATCAACTCCCTGATGGAGATTGGAACAGG
 TGTGTTCTAATCTAAATGCATCACTTAATATTAGTTCCAACCTCTCTGGGGCTTCCTTCAATAGGGGAA
 TTAGACTGGTCTCAATCTCTTTGTACAGATGAGTAACCTTTATTTACCCAAAGATTAGTATTAAACAGTCG
 GGAGCAGGAGGGAGAATACTTATGAGACAACAGCCATTTCCACAGTGGAGAGGAATGGTTTGTCCCAATA
 GAAGTTACCAGATTTCACTCCATTGCCAAATAGATATTATGAGCAAGGAAGAAATCTATAGTAGTAACCT
 AAGACCACCAGAAAGATCAAAGCCAGAGGGTGAAGGATGGAATAAACATTAGACATATCTCTAACCCCT
 CTTTGTGTTGAAATACTCATTACCCTGTGGTACTGGGAATACCTGTGCCTACAA

The following amino acid sequence <SEQ ID NO. 71> is the predicted
 amino acid sequence derived from the DNA sequence of SEQ ID NO. 13:

LAQVFPVPGNEYFKQKRVDRMSNVYCHTLTLWALIFLVLSYYYRFLPCSYLFGNGTEIWLLGTHNHS
 LWKWLSSHKYSPPSCSRLLILNLWNKVTHLYKEIGDQSNPIRKPPQVGTNSVMHLELEHTCSNLQSGKLI
 VLWLLKQRTGSAEKPMNKPVPYGFLLKSEFRAQNESIYLVLTHTSIKNEETGAELLKNIPVSCARTGH
 PYVLTLP

The following DNA sequence Seq-2593 <SEQ ID NO. 14> was identified in
H. sapiens:

TTCTGCCATCGCAAGGGGAGGGAAGAGCACCTAAAGGGCTTATGAGAGGTTTGAAGTACCAAGGGAGGAA
 CAGAACACATTCTTTCCATTGGCTAGGACTCGGTACATGGCTTTCCCTCATTATTAGTGAAGCTTGAG
 AATTCATCTATTTGTGAGCCAGGAAGAAGAGAAAACAATTTGGTTAAACATTTAGCAGTCTCTATGACA
 ATAGTCTGTATGTTGACTGCAAAGGTGGATGAACAAAACCAAGCCTCCTTTAAAGCAATACAATCTGGCAG

AGTCCCTGGGTTATCATCTCTGAACATAGATGCTTATTGTTCAAGAGTTAAGAAAATTAGCATGACTGCATT
CCAGTTCTATAAATTTAATCTTTATTACGATATTGTCCATCCACATGTCTTAAAAATAAAAAAACA
AAAAACCTAGTAACCTACGTTTTATATAGCAAGGAACACTCATATATATCACTTCATTGTATCCTTACAACA
ATCCTGTGCAGTATATGTTTTACTCCCTTTCTTCTATGTTTTGTATATAAAGAAATGAGCCCCAGGGAGTT
GAATGGCTTGCCCCAACTAGTGAAGCTAAACTCCAATCCAGGTCTTTTTATTTCCAAATCCATAATCTAC
AACCATCTGTAGAGAGTTATAATTAAGAGATATGAATGGTCAGGGGCCTTTCCATTTCACTGCAAGTCTGC
CCAGCTCCAACCTACCAGCATCTG

The following amino acid sequence <SEQ ID NO. 72> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 14:

LPSQEGRAPKGLMRGLTDQGREQNTFLSIGDSVTWLSLIISEAWRIHLFVSPGRRENKLWTFSSLYDNSL
YVDCGGTKPSLLSNTIWPSPWVILNIDAYCSRVKKISMTAFQFYKFNLYSAYCHPHVLKNKIKNKKPSN
YVLYSKEHSYISLHCILTTILCSICFTPFLLCFVYKEMSPRELNGLPQLVKLKLQSRSFYFIHNLQPSVE
SYNEIMVRGLSISVQVCPAPTTSI

The following DNA sequence Seq-2594 <SEQ ID NO. 15> was identified in *H. sapiens*:

AATCCTGCATTTCCCATGCTCTGGGGTGAGAAGGAATTAGCTGGGAGCCAATTAGCAATCTTGTGAGAAGC
AAATGAATTTTGAATAAACTGGATACTTAACTGAATGAGACCATTGAAACCAGAAGAGCCTGAGATCCA
TCAGTTAGAGGAAATAAAGAAGTGGCATTTCCTTGCCATCTGGGTGCAGTGTGAGTGATTTTTATAATC
CTACCACATTTTTATCCCTGCTTCCTTAACTGTAGACCCAAGGAACCTGGCTGTTTTGTTCAACATGA
TGTGACCCCATACCTAACAGGCCAGGCACAAAATTGGCCTCCAATAAGTAGTGGATCAAAGTATGAATGG
ATAAACTGAATGAATGAAGCCAACTTGAATTTCTCCATAGCTTATCCAAATGGGAATGGTAAATCATATA
AGCTTTTGAGAAGAGAAGCTTATTAAGAAGCCCTACATCAGTCATGACTGGCATCATTGGTTAGTTACCCA
ATTTTCTCCTTCCCTTCATCTTCTTAATGCAACTCTGGTTTGGGCTGCAGTATACCCAGTTAGAATACGCC
CTCCCCAGAGTCTCATCCAGCTGGAGATGATCAAGTCACCAAGTTCTAGCTAATAAGTGGCAGCCAAAGTA
TTCAGGATGAGACTTCCAGAAAAGCATTGTTTCTGATAACAATGGACAGACC

The following amino acid sequence <SEQ ID NO. 73> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 15:

SVHCYQENNAFSGSLIINTLAGNLLARTGDLIISWMLWGGRIILGYTAAQTRVALGRREGENWVNPMP
VMTDVGLLNKFSQQKLMIFTPIWISYGEIQVWLHSFSLSIHTLIHYLLEANFVPGLVRYGVTSTCKQPGS
LGPTVKGQKCKGRIIKITHAPRWQKCHFFYFLMDLRLFWFQWSHFSLSIQFIQNSFASDKIANWLPAN
SFSPQSMGNAG

The following DNA sequence Seq-2595 <SEQ ID NO. 16> was identified in *H. sapiens*:

GACTGTTATTACTGAAAGTCATTGCTTTTAGACTCTTCCAACCTACAGAGCAAGGAAGTTTATGTGTATATA
GTAATCTGTGAATATACACATACATACATATTTCTATATGTAATCATCCATATTTAAATTAAGTAGAAT
ATGAGTTCATACTGATATCTCCAATCCTAATCAGTTACCACAGGATTATTCGGCCTTTTCCCTTGGA
GTTTGCAACTCCTGCTTCAACAGTTAGAAATCTGGCTTCCATATTCATTGCTTAATTGTTCAATTCCAGT
ACACATAAATGGTGGCTTCAGAATTAATAACTTATACCTCCATGGGAAATAACTTTATTAATAAGTACA
GCACCTTATGTATAGTACTTTTTGAATTTTGTAGACTTAGAGATTCTCTCTTTTCCAAAGTTACTTAGGTC
AGAACCATTTTCATTCTTCAGTGAAGTTGTCTTATGTATTGTAATACAGTTAGATTGTTCTGTCTATG
GTGCATTCCATCCTGGGATTTCTATCTCTTTTTTAACATTTGCATATATTAAGTTTCATTCTTTGTGC
TGTATCATTCTATGGGTTTCAATTAATGCATAGTGCATGAATCTGCCACCATAGGAGCATCATACAGAGT
AGTTTCACCAACTTAAAAAATCCCTATGTTTTAC

The following amino acid sequence <SEQ ID NO. 74> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 16:

LLLLKVIAFRLFQLQSKEVYVYIVICEYHTYTYFYMSSIFKLSRIVHTDISNPQLPQGLFRPFSLSLQ
LLLQLEIWLFPYSFALFNSSTHKWWLQNLIPPWEITLLTKVQHLCCIVLFEFLDLEIPLLFQSYLQNHFFP
FSEVVLICINTVRLFCHMVHSLGFPISFFNICIYVSFFCAVSFYGFQLMHSMVNLPPHHTEFHQLKKFP
MFY

The following DNA sequence Seq-2596 <SEQ ID NO. 17> was identified in *H. sapiens*:

CGTTCCTTCTCTGTGCATTAATGGACATGATGATAGTTGGCTCACTCAGTAAACATTCTGTGTCTGGAAG

GATTTGATTGTCTTTTCTCAGGCAACAATTTGAGGTGATTGAAAAATCTTTCTGAAAAATTAAAA
 AATTTTCTAATTAATAATGCAAGCTCATTAGAAAAAATGAAAAATAATAAGCACAATTTTTC
 TTAACCACTTAAGATGACCATTTGTTAGTTTTTTTTTTTTTTGGTGCCTTTTCCGTTTCCATCTCTTT
 TCTATTAACACTTCTGAAATGTGATTGTAGCAATGACGCATAAGGGGCCCTTGACACATTGAGAAATTTAT
 AAATACGCTGGCTTCTTGTCTTGTCTTTGTCCCCAGCTTAAGTGGGAACCTTTTTCTATATCTTTGAAAC
 TCCAAATCCTAGATAATTCTTCAAGGTCAAGCTCCAATGTCTTGTCTGGATTCTTCTCAGCAGGAATTGATC
 TATTTTCTCTGTATTTTGTGCCACAGGATCTATGACTCTCTTATGGCAACTACCACCTTCTGCCTTATAT
 TACGATTTTGAATCTTCAACAAAGTCTAATTTTTTTTTTTTCAAATGAAGTCTCGCTATATTGCCCAAG
 CTGGAG

The following amino acid sequence <SEQ ID NO. 75> is the predicted
 amino acid sequence derived from the DNA sequence of SEQ ID NO. 17:

FLPLCHNGHDDSWLTQTFCVWKDLICPFLEATILRFEKSFLLKNIKFLIKNNASSLEKNKINKSTIFLNHLK
 MTIVSFFFFLVLSVSNLFSIKTSEMLQIRGPHIEKFINTLASCLAFVPSLTGNSFSISLKLQILDNSSR
 SSSNVLLDSSQOELIYFLCIFVPQDLLSYGNYHLLPYITIFESSNKVFFFFQMKSRVIAQAG

The following DNA sequence Seq-2598 <SEQ ID NO. 18> was identified in
H. sapiens:

ATGTCATGGGAAATGCAAGAATATGTGTCCAGCATGGAAGGAATCAGTATGGAAGTCTTTTGATAAATTG
 TGCCATTATCACTAACATTGCCTCAAACTTTAGACTACCTGCCATATACAAATTAGAGGTGAAAATTAC
 TTCCATGTAATATACAGCCAACACAAAGAATCCTATCCCAGTTTCTTGGATGGATAGGCAAGAATCTGGG
 TAAGGTTTATTGTGCAATAATCCTCTTCTCTCTCTATAGGCCAGGATTTAAGTTTACCTCAAAATGGAA
 AATTTTGGCTGGGAAATTACATGTGGGAAGACATCTTCAGTGGAGATTTAGTAATTACAGTTTCAGCTA
 TGACCTACCCCTTTTCTACTAGATTCTGCCCATGTTGGCCAGAATCCCTAGAAATCAATTATGTTTTGA
 TCATCATCTATGCCCTGATGTTTCTACTGAACGTGATGTGAAACTCCCTGCCGATGCTGGTCATCTTATTC
 AGCTGAGTCAGCCACTGTCAACCGATGTCTACCTGCTGACCCTGGCCTTGCCCGACCTGTTCTTTCCCTGA
 CATTGCCCATCTTGGCTGCCCTCCAAGAATGAATGGCTGGGATTTTTGGCACAACTGTGCCAGGTGGTCTA
 GCTCTGAAGGAAGTCAACTTCTACGGGGGTATTCTACTACTGGCCTGCCGAGCATGGGACTGTTA

The following amino acid sequence <SEQ ID NO. 76> is the predicted
 amino acid sequence derived from the DNA sequence of SEQ ID NO. 18:

VMGNARICVQHGRESVWKSFDKLWHLSTLTPQNFRPLPAIYKLEVKITSMYTSQHKESYPSFLDGARIWVRF
 IVQSSSLFYRPGFKFTSKMENFGWENYMWEDIFSGDFSNSYSYDPTFFLLDSAPCWPESELENYVLIIY
 ALMFLNVMNSLPMVLVILFSVSHCHRCPLPADPGLGRPVLPDIAHLGCLQEMAGIFGTICARWSSSRKSTS
 TGGILLACRSMGLL

The following DNA sequence Seq-2600 <SEQ ID NO. 19> was identified in
H. sapiens:

TATGATATTTTACGCCATGGTGCTGAACATTTCCAAACAGCATAAATGCACCATGTGTGTATGTTTTCTTT
 GGGATGCTGTGCTTAGAGGGTAGCAGACAGGGTGCAAAGTGAGAAGGACCTGGCTCTGCACCAACACTGC
 CAGTATTGAATCCTGACTCCATCATCTGGGAGCTGTGCAACCTACGCAAGGTACTTGGCCTCAGTTTCCTC
 ATCATCCCCATGGCATTTTTGTGAGAATTAATGAGCTGAAACCTTGAACCCCTTCAAACAGCAGCTGGC
 ACAGAGGAAGCACACAATCAATGTGAGCTGTACTCTTCTGGCAGTGTGGAGATCCCAGCTCTGCCCTAG
 CTAGTCACTTCTCTTCTTGAATCTCAGTTTCCTTCATCTGGGAAATGGGAGCAGATGTGAAAGGGGAGG
 GTGAGAATACATATGAAAGTGCTGGCTCCTGGTGATAGCAGGCACCTAATAATGATACACTTTTCCATCT
 TCTGCCTTCCCAGGGATGCATTGTGCCATGTAAGAGAGAGAGCCTCCAGGGTTGGCGAGAGTTTTTGATC
 CAGGCTTTTTTCAAGGTGTCAAAGATGAGCTGGGTGATTCTCCATAGATTTTCTTTCTAACAGGTGACAGTT
 CTGTTTCAGAAATACTGTGGATGTTTACAGGTTTACAGCACAT

The following amino acid sequence <SEQ ID NO. 77> is the predicted
 amino acid sequence derived from the DNA sequence of SEQ ID NO. 19:

VLTTSTVFLKQNCHELLERKIYGESPSSSLTPEKAWIKNSRQFPWRLSLLHGTMHWPGRQKMEKCIILKLLC
 TRSQHFHMYSHFPFHCISHFPDEGTEIPRREVTSGQSWDLHTARKSTADIDCVLPLCQLLEFVSRFQLI
 FSQKCHGDDEETEAKYLAVAQLPDDGVRIQYWQCVVQSQVLLTLHPVCYPLSTASQRKTYTHGAFMLFGNV
 QHHGNII

The following DNA sequence Seq-2601 <SEQ ID NO. 20> was identified in
H. sapiens:

TTTATGCTCATGTAGTTCTTTCCAAGAAGAGAAATTACAGAGTCAAATTGTAGAAATATTTAAAAATCTTT
GGCACACATAAACAGTATCCATATAAATTATACCATCTTTTAGATGAGTTTAAACACCAAATGATAGAAAT
CTCAGTTTCATACAGATTTGGTGGGCTGGAACCAATACTTGCTGATAGGCTGTCCCTCGTCTTTCCCTA
GCTGTTCTGGGAAAGGCAGTTCTGGTAAGAACTCTCCCTACGGCCCTTTTCATCTCACTGTTCTCAGGG
CATAGATAAGTGGGTTGAGCAGTGGGGTTCCCAATGTGTACACCAAGTGAATGATCTTGCTTGGGG
TTGTAGCTGGAGCTGGGGCACAGGTACATGAAGGCACAGCAGCCATACTGCAGCAGCACAGTGAAGT
GGAAGAGCAGGTGGAGAAAGCCGGTGGCGGCCAGCAGCCGAGTGGATCTTGA

The following amino acid sequence <SEQ ID NO. 78> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 20:

KIHSAGRHRAFSTCSSHLTVVLLQYGCCAFMYLCPSSSYNPKQDQFISLVYTLGTPLLNPLIYALRNSEM
KGAVGRVLTRNCLSONSERRGDSLSGKYLVPAPQICMKLRFLSFGVKTHLKDGINYMDTVYVQRFNLNIST
ILCNFSSWKELHEHK

The following DNA sequence Seq-2602 SEQ ID NO. 21> was identified in *H. sapiens*:

TTTAAGCCACCCAGTCTGTGGTGTCTGTTATGGCAGCCCAAGCCAGCTACTACAGGGTGGGACGAGGGGA
GGAGCATGGCCTCTGCTGGAAGTGCAGGCAATGATCCCCAGGAACAATGATGGGAGCTTCTGATTGCTC
TCATTATCTCTGCAAAGTAGGAAGAAAGATTTCATCAGCTGAGCATGAGGATGGTAGAAAACATCTTTGGGA
AATTCAGAAAGTGAAGGAAGGCATAAATAGTCATCTAAAAAAGCAGGAAAGGGAAAAGACAGAGAAATC
CAGTAGTATGCCAGGACTCCAGGAAGCATCAGGACCCACTTGAAATTGCCAATGCTGAATTTAAATGAG
GCCAGTCTGTACAGAAGCACTTCTGGAAATTTGCTAACAGCTAAATAGAGTAGAATCAATACCTTAGAGAAAT
ACGAGTAACCAAGGAATAAAATTAAGTATCACTTTTGTGGTTTTTACTATTAATATTTTCTTCAGTGT
AAATCATAGCTGCTGAATTCCTGAACCCCTCTTATATAAATCTAAAGCTCTGGTTTATCATGGTTGAA
AATTCATGGCTAACTTATCAGGCAAACTGTCCCTAAAGCATTTTTTGAATAGCTTTAGTATCAAGATGGTA
CTGAGTGATACATTTCAATTCCTGCTTAAAGGAAGGCTTAGTTATTTTAAACCAAGTCTTATTTTTATAG

The following amino acid sequence <SEQ ID NO. 79> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 21:

IKIRLGLKLSLPLSREMKCTLSTILILKLFKKCFRDSLDPDKLAMNFQPTRAFIYIRGVQEFRLFTLKKIL
IVKTKVDQLILFLWLLVFSKVLILLYLAVSKFQKCFCTDWPHFKFSIGNFKWVLMPLPGVLGLILDFSVFS
LSCFMTILCLPSLLKFPKDVFYHPHAQLMNLSSYFAEIMRAIRSSHCSWGIICLHFQQRPCSSPRPTLL
AWAATEHHRLLGL

The following DNA sequence Seq-2603 <SEQ ID NO. 22> was identified in *H. sapiens*:

ATTTCTGGATTTATGCCTCCCTGACCCATTCCAGGATTTACCCCAAACCTTCCCACTCTCTTCTAACAG
GGAAAGTTCTGTTATGACACAATAGTACTTATTAAGACAGATTTACCTTCTAAGTCTCAGGACAGCATTT
ACAACCAGAAATAAAGTGGTACATGAAGAACCAGGAGTCTGGTAGTAGTGAATTCATTTTCTTCTTGAA
AAAGTGGATCAAAGGATTCAAACAGCAAGTGGTGAATCAATGAAAAGTGGTAAATGGTGAGGAAAAATG
TTACTAAAAGATGACCTCAAGATTACTGGTGCATATGAATTGCTTTTTTATATAGGAAAATACTGGATAAT
TTCTTATTGTATAGTATAATTAGAAGCAATTTTCATGTGTTTCAATTTTGCACATGAGTTTAAATGGAATAG
ATTTGGTTCCCTCTTAACATGAGTTCAGTGTCTGAACTTGGGCAAAATTTCTAAACAATTCTGAGCTTAC
TACCTCTGCTTGAAAGTGAG

The following amino acid sequence <SEQ ID NO. 80> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 22:

SLSSRGSEAQNCLICPSSDTELMLEPDLFHLNSCGKMNTNCFLYYDNKKLSSIFLYKKAIHMHQSGHL
LVTFPPHFTTFHFTTCCNLPLIHFFKKEHNFHYQTGSSCDQLFLVVKCCPETKVNLSVLLCHNRTFPV
RRECGRFGVNPQMGQGRHKSRL

The following DNA sequence Seq-2604 <SEQ ID NO. 23> was identified in *H. sapiens*:

CTTTGGAATTTTATTCTAAGCATCAATCAAGAGGTATAGTACGAGAAAGGTAGAACATGTAATTATAAAT
CAGGATTCAGGAAGTTTATTTTCTCTCTTTTAAATCTCTCAAAATGATCTTGATTCCTGCAAAGTGT
AGTATATCTGTAAGTAAGAGTCTATTCTTTTAACTTCATCTGTATTAACCAGCTTTATATGACCAAAA

TGCCCCCAAATTTAAATCTTTGCACAGTAAGGCCTTATATGTACACCTGGCCTCATTTCAAAGACTAAA
GCAGTTGTTCTCAAATTCAGCTGCACATTAATATAAAGTGGAAACTGTTTAAGCTCCTGATGACAAAGCC
ACATGTGAGACTAATTTATGCTGAATCACTGGGCCAAGGACCCAGGTATCAGCATTTTTTAAACTATAGA
GGAATAACCAGGGTTGAGAACCCTGCACAAATGGTAAATGCACTTTTATTTAAGTTATTTTTTTAA
TAAATAATGGTTGAATTGATACTGATCTTAGTACCAAGTCATGGCAATTTTTTCAGACTTAGAGAATTCAT
CCTGGCATTGAGATTATTAAAGAACCCTAGAAATCCAAGTGTGTTTATATTTTCTGTAAATATTAG
AGTATGCTAGTGCTCATCCTTATTTGATAATTTGGAAAAATATATTAAACATTT

The following amino acid sequence <SEQ ID NO. 81> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 23:

LEFYSKHQSRGIVRERNMLIQDSGLFFSSFFSQNDLDSCKVLVYLVSKSLFLLNFICINQLYMTKMSPKF
KSLHSAKALYVHLASFQKTKAVVLKFSCTLTITGKLFKLMTKPHVRLIYAESLGQGPYQHFLKLRNNGEP
LHKMVNATFIVIFFKIMVELILILVPSHGNFFRLREFILALRLKNLEIQVFLFIFFLILEYASHPYLI
LEKYIKTF

The following DNA sequence Seq-2606 <SEQ ID NO. 24> was identified in *H. sapiens*:

ATTATCATTTGAATGTTGATATTACATCATATACAAATTGATTGCAACATAGTTATTTAATGTAACATTCT
ATTTTAAAGATAAATTTATCAGAATCATACATTGCTACAGTAGTCTCCCTTATCCACAGGTTTCAATTTCT
ATGGTTTCAGTTACCTAGTGTCAACAAGGATCCAACAATATTACATGGGAAATCACAGAATAAACAGATT
TGTAAGTTTTTAATTTGTGCGCTGTTCTGGGCAACGTGATAAAATCTCATGCTGTTCTCTCTATCTGCGCT
GAACATGAATTATCCTTTGTCCAGTATATCCACACTACATATGCTACCTTCCCATTCATCATTTAGTAGCT
GTTTTGATTATCTGATAGAAAAACACACAGTATATATAGAGTTTTTATGGGGCAAGGAAAACTTTCTC
TTTGTCTCTGAAGATTCACTGAAACTCAACTCACAAGGGCAGACTAATAGGAATGAAGGTAAAAA
AATATATTACATCAATGGAGATAACTACAGAGTGATTATCCATTGCCATCAATGGACTACAGTGGCTTA
AATATCGTTTTTGAGGTTACAAAAAGAGTGGAAGTCTTGGGATCTTGGCAAAACAGGTTATGGGAAGAAGAG
AAGAGAAACCTGGTTAGCAAAGGTCACTCTTGTGATGC

The following amino acid sequence <SEQ ID NO. 82> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 24:

IIIMLILHHIQIDCNIVICNLFKINLSESYIATVVSILHRFIFYGFSYLLSTRIQQYYMGKSQKTVCKFF
VRCSGQRDKISCCSSSLCLNMNYPSSISTLHMLPSHSSFSSCFDYLIKETHSIYRVFYGARENFLFVLR
TENSTHKGRIGMKVKKIYHQWRLQSDYSIAINGLQWLKYRFEVTKRVEVLGSWQNLWEEKEKRNPGORS
SCD

The following DNA sequence Seq-2607 <SEQ ID NO. 25> was identified in *H. sapiens*:

TTTTTCCCCCTGAGTGTCTCTCATGCTTTCTCCAAATGGAGATGGAGAGGTTTCACCTCACTTTTCT
CTAACTCTCCCTAGTTTTTTGGTTTCTTTTCTCCACATCTAAAAGTGTGCAGAATGTCCCTTTAGCACAT
AGAAAAATCTTTTCTTGACCCTGCCACCTACTTAATAAATCCACACTTTTCTTCTTTAAGATTTC
CTTTATAATGGTGTGTGCAATGGCCACATCCACCTTATCCATTCTTCTTAAAGTTCAGAAAAACGGTT
TTGTTTCTGTTACTTTAATGGAATTATTTTCCAAAGATCAACAGGACTTTCCCTCAAGCCCAATCCAGT
CGGTAG

The following amino acid sequence <SEQ ID NO. 83> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 25:

FFPLSVSLMLSSKWRWRGFTSLFNSPFFGFFSSTSKSVQNVPLAHRKSFLDPATYLTKIPHFSSSFKISF
IMVCVNGHIHLIHSFLKFQKNGFVSCYFNGIIFPKINRTFPQAQSSR

The following DNA sequence Seq-2608 <SEQ ID NO. 26> was identified in *H. sapiens*:

ATACATGATAAGGTACATGGATCCAGGGGAAGGATGAAGGGCAGTGTGGGATTGCTTTTGAATTTCTCCAA
ACTCGCCCATAAAAGCAGACAGGACAACTAAGATAACTAAACAAAAAACCCACAGACAAACTATTACA
AACCCCAAAAAGAAGTGTGGTGGGAACAAACATCTGATAGAATCAGACACATTACTGGTGACCGGACATAAG
CCCTGTTAATGAGAAGCTTACATTTAGGAGAGTCAATTAAGTACACGCTATACACAACCTAAAGTGGTAAA
TGCTACCTTGGTTATTCAACTTCACTGTTACATGCCTTGAAGTGTGGGTGCACTGGCCTGAACCATCTG
GTTGTGTTTGATTCTTAGGATGCCACCAACAAATAACATTGAGAAATACCCAGCTACTTTTCATTGTTCT

CCAATGGCAGCAAAGTACAAATGATCTCTATGA

The following amino acid sequence <SEQ ID NO. 84> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 26:

IEIICTLLPLENNEKLGISQCYLLVASGIKHNQNGSGQCTPHFKACNSEVEPRHLPLVVYSVYLIDSPKCK
LLINRAYVRSPVMCLILSDVCSHHTSFGVCNSFVCGFCLVILVCPVCFYGRVWRNSKAIPHCPSSFPWIH
VPYHV

The following DNA sequence Seq-2609 <SEQ ID NO. 27> was identified in *H. sapiens*:

TCCAGGCAGTATTCTCCATGACAATGAGGAAGGTAAGTCTGCAACAGAAGAACAATGGCAGAAATTTTAAG
AAAAGTTTACCGCCTGGGACTATGACTCACCTTTTGGGAGAAAATGTGACTAACCTTTGTAAGAGCTTGT
TGAGAGCTCACTTTCTGGGAGGAGTCGGGAGAAGGGGAGCATCAGCTGACGAAGAGGTGAAGGAGGTACC
CAACAAGAAAAGCGTAGAAGGACCAGGGATTGGGGTCGGGTCTTCTCCTGATTCCAAGGGATGGCATAA
GATATTGCCAAGTGAAGGAAGCGAAGTAGAGCCAGCAAAGGAAGGTGAAGTGTGTTTCATTAGAAATAAT
ATGTTGTGATAATTATACAAAGTACTAATTAGTAAATTCTTTTCCAACTCGACACTCCAAAAATCCCTGT
ACTTATATCCCGAAGGCCTCTTCTTCCCCAAGCTGGAAGACACGGTCACTCATTAGTCACCCACTGTCACA
GGAGTAACAGAGACTACAAATATTGGACAGGACATAAGTGAGGGTCAAGCATCTGGATGCAGATGCATGAC
AGGATGCAAGTCTTCCAGCTCTCATGGACTTTGCGACAGATGCACAGAGTGAGGTA

The following amino acid sequence <SEQ ID NO. 85> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 27:

TSLCASVAKSMRAGKTCILSCICIQMLDPHLCPVQYLSLLLLQWVTNEPCLPAWGRRGLRDISTGIFGVSR
LERNLLISTLYNYHNILFLMKQQFTFLCWLYFASFTWQYLMPSLGIRRKTRPQIPGPSTLFLLGTSFTSSS
ADAPLLPTPPRKVSSQALTKGSHFLPKGESSQAVNFSNFCHCSSVADLPSSLSWRILPG

The following DNA sequence Seq-2610 <SEQ ID NO. 28> was identified in *H. sapiens*:

ATTAAATGCCTGAACTCCCTTCAGCTCTGAACTCTGTGGTGTATTCTCTGAGGACATTACCTCTCTAAGG
GACCCAAATTAACAGCTCACACCATCCATCATTTTTCTGTCTGAGGTTTTATTTCCCTAATCAGATTGG
GTAAATTTTCAGCCTCTCTGTCTCTTACTTCCAATCCAATAAAACCTGTATGGATTGTTTTGTATTTT
ATCTAATGTCATTATTCTTCTAAGAGTCACTGCCTGACCATTTCCCTGCCTATAGCATAATTAGCTATTA
AAAAGCTACACTGGCATGGTTTTCAAACCTGCATCCTCTTTTTCTGAGGTGGATTGATTCTAAACTGATTA
AAATATCTCAGAATTTCCAATACAATTTTTAAATGCAACAGATTTTCAAGACTGCCTCATGACTCTGCCA
AGCCAAGGGAGTTAGCTGCCAAGTCTCTGACTGCCAAGGAAGCCAAATAAATAATCTGATGGTGGTTT
TAAATGAGAGGCAAGTGCCCATTTCTTAGGTTGACAGTGCCACCTACACATTGACTTCTCCAGGGTTTG
TAAGACACCAAGGGTGATGTTTCAGATTTTCCC

The following amino acid sequence <SEQ ID NO. 86> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 28:

LNATPFSSETLWCILGHYLSKGPKLNSSHPSFFCLRFYFPNQIWNVNFQPLSVSYFQSNKTCMDLFCISSN
VIIHSKSHCLTISLPIALAIKKLHWHGFQTCILFFGGLIILNLKYLRISNTIFKMQQIFKTASLCQAKGVSC
QLSLTAKEAKIILMVVLKEASAHFLGQCHPTHLLQLGLDTKGDVSDFP

The following DNA sequence Seq-2611 <SEQ ID NO. 29> was identified in *H. sapiens*:

TCCACATTCTTTCTAAAGTTCTGAGCTTTTCCATGGGCTTCCATGGTAGGGAAAGCACATGGCCTGGGTGT
GGGTAGAGCAGGTGCGGCCATTATATGTATGTTCTTTGCAAGTCTGGCATTGTAAAAATGGGTGATGCT
TGTATTGTGTTTATTATTCAATCATGTAATAGAAGATGCACATAAGATTATTTTGAAAAGTATGCCTTCC
ATTTTCATGCTGAGAATAATGCAGGAAGTTCAGTGTAAATGCAGTTATAATAAAATAGTAGCAAAACAATAT
TTTGCTTTAAATCATGGAATTAGCAAGTAAAGACTAATTGGAAGCCAATCTTTTGCAAATTTTTTAAATGT
AAGTTTATTTGGAGGATATGACTTGTGGCCAGAGTACATATAAAGAACAAAAGAGTATAATTAACAACA
GTTTCAATATGGACTTACCAGGCATCTTGATAAAATCAGTATTGACATGTATGTGAATGCCAACATTGTG
TTTTTCAATTCAATATCTATGTTATGCCATAAACTGGTAGCAGTTATGAAAATTAGAATTGGTTAAAAAC
TGTTGAAATCTTTAAATTTTTCTGTTTA

The following amino acid sequence <SEQ ID NO. 87> is the predicted

amino acid sequence derived from the DNA sequence of SEQ ID NO. 29:

NRKNLKISTVFNQFFSLLPVLWHNIVLNWKNMTLAFTYMSILILSRCLVSPYLKLLLIILFCSLYVLWANK
SYPPNKLTFKKFAKDWLPISLYLLIPFKAKYCFATILLHYTELPALFSKWKAYFSKSYVHLLLDINKH
NTSITHFTNARLAKNHTYKWPHELLYPHPGHVLSLPWPKMEKRLTLERMW

The following DNA sequence Seq-2613 <SEQ ID NO. 30> was identified in *H. sapiens*:

TACGTGGGTTTCCCTATCGTCTCTTCATGAGTTCCTTGTGAAAACAGAAAGACTGAGTCTGCCAATAACC
AGCAAGAGAACAGATAAAATAAAATAAATAAACCATAAGACTTTAACATATGACAAACAACCTGGTAAGGA
TTTTCAAATCTTTGGTCAACTTTGATGGTATTTTCCATACAATGAACCTCTAAATATGAAAAACGTAC
ATCCATATTTTAGATATAAAAGTCTCTTGCACAGGCCAGAAAATGAACTTAAATTTAAGCAATAAAATTC
CCCTTTGTAGACTGCAAAATGGAGAACATGCTATCTAGCTTCATTTTCTTCAACTTACATAAAATGAAAC
AATGGTTAATGTTCTGGCGGCATCTCTAAACATATTCAGTGAAACAAAATTCCTTACAAATGTCAACAGC
TTACAACAAATAACATTTTATCCTGTTTAATTATTAGAAACAAAATCAGTTATGCTGAGATATGTTTGCA
TGGGATTATATACTCTGATCATAGAAACAAATTATTGACATCTGAATCTGAAAGCTGCAAAACATGATAA
AAGACATAATAAAATCACAGATTGTTATTCTCTCAGGAACCTTTTCTAG

The following amino acid sequence <SEQ ID NO. 88> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 30:

KKFLREQICDFIMSFIMFCSFQIQMSIICFYDQSIIPCKHISALILFLNNTGNVICCKLLTFVRKFCFTEY
VRCRQNIHCFIFMVEEKSIACSPFAVYKGEFYCLNSFIFWPVQETFIISKIWMYVFHILEFIVWKNITIKVD
QKILKILTSCLSYVKVLWLILFILSCSLAGYWQTSFCFHKELMKRTIGKPT

The following DNA sequence Seq-2614 SEQ ID NO. 31> was identified in *H. sapiens*:

GGTGCCCATGCTTGGTGGTAGGATGTATGAAGCTCCTTGCTCCTCCAGCTGGGCATCCTGCCACTTGCTGA
GCCAAATAAGGAATGTGGGGAAGCAGCAGGCCACCAGCCAGATGAGGGCCACTGCCTTCCAGGCAGCCCCA
TGGGACATGAAGGAGAGGTAGCGCAGTGGATGGATGACTGCCAGGTAGGTGTGCAGCACAATGGCGGTGAA
GGACAGGATGGTGTCTGGTGCAGGCGGCGAAGACAGCATCAGTGAGAATGCCACAGGCCATGGGCGCCAGCT
CCCAGCCACCCAGGCTGCTGGAGGAGATGAGCATCTGGAGGAGAATGTAGGCCAGGTCTGAGAGCAGGATG
TTAGCCGGGAGCAGGTAGTGGGCTCCTGTGCGAGCCGTTGGTTCGCGAGGATGGTCCACAGCAGCAGGGG
GCTGACAGCCAGTGTGGCTGCAGCCAGCAGGCTTGAGGGAAGGAAAAGCCAGTACAGCATGGAGCTGGGCA
CCCTGAGGTCCCCCAGGCCCAAGGAAGTGTGCTGCTGCTGGGGCATGCAGGGTGTCTTGCTGATGAGC
TGGATCAGGGCCGCGCAAGCTGTAGTGCCACAGGGCAAGGTGCCAGCTCATCCCCCATGCTTCTCTGGCAG
GGATGGCTGGCTTTGT

The following amino acid sequence <SEQ ID NO. 89> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 31:

QSQPSLPGSMGDELAPCPVGTTAWPALIQLISKTPCMPQAAASNTSLGLDLRVPSSMLYWLFLPSSLLAAA
TLAVSPLLLVLTILRNQRLRQEPHYLLPANILLSDLAYILLHMLISSSSLGGWELGRMACGILDAVFAACT
STILSFTAIVLHTYLAVIHPLRYLSFMHGAAWKAVALIWLIVACCFPTFLIWLKWDQALEEQGASYILP
PSMGT

The following DNA sequence Seq-2615 <SEQ ID NO. 32> was identified in *H. sapiens*:

AACACTGACTTCTCTGAAGCAGTTGTCTAAAAGAACCTACACCATTTTATTTAGCAAAAAGGCTTTTGT
AAAAGCAGGGGATAGCAGAAAGAGCTTTGTAAAAAATATGTCATGGATTTTAGGAGTTTCTAAGAGCAAGA
AAACGTTTCTTAAATAGAGGAATGAAGCAATTAGAGTTCATAAAAATCACCTAATGGGCCTTCCAAAAGG
CAAAATGCTAAAGCCCCAGAAATCATCACTGAGGAAGTCTGAAGTAGGAAGAGACCTTGTCTAGAAAGCCG
ACAAGGTAGAAATTAAATGGAACAGGCCCAACTTGAAATTCGAGACCAAAGAGGAGCTGATGACATTG
GTGGGAGACAGGTGTGGGAATAAAGAAATGTTGGTAGATTCTAGAGACATTCCAGCGATAACACAGACAGGA
CTTTGTGACTGACTGTATGGGCGAGCTGCAGGGGTAGGAGAGGAGGAACGATTAAAGACATGATGAAGTGGG
CTATGAGTTGGCAGCTCCATTTACTCCAGAGAACACAGGAGGTGAAAATCATGGGAGACTTGATGAAAACA
CTTTGAGAGGCACCATGGGGATAAAAGCCAGAAATAAGGTGGGAAATGTTGGAAGCTATTCTTAGAAA
AGAGGTTGGGAGGATGAGCATAAGTTAACAGGAACAAGTTAATTTTTTAAAGTGCT

The following amino acid sequence <SEQ ID NO. 90> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 32:

HFKINLFPVNLCSSSHPLFNEPFPPTLFLAFIPMVPLKVFSSSLPFSPPVFSGVNGAANSPPSSSCLNRRSS
SPTPAAAPYSQSQSPVCVIAGMSLESTNILYSHSTCLPPMSSAPLLVSEFQVGPVFFFLPCLRSRTRSLPTS
DFLSDDFWGFSICLLEGLGDFYGTLIASFLYLNRNVFLLETPKIHDIFFTKLFLLSPAFNKSLFAKKWCR
FFTASEKSV

The following DNA sequence Seq-2616 <SEQ ID NO. 33> was identified in
H. sapiens:

TTTCCAGATAAATTGTATGCACAGTAAGTGGTGTGTCAGTATACCATAGCATATATACATCCATTGGCA
CACTGCAGGTGCCAGTGGGACAACATACCAGAGTGTAAAGTCTTCCTGATCATTTTCATGATGTCCCTCAGTT
ATTTACCTTGTAATAAGCTTGTAACGCTCTATGATTGTTTTGAGTCATCCCAATGCAGTCATGTAATAAC
AACATGTATTTTAAATGAACTTGGGGATTTTCTCCATACCTGAATCTCTAGTATTCACATAAATGAAAA
ATCAAAATTAGGATAAGTTAGTGTCAAACATTAATGGATTTTTACAATGCTAATTGGTGTCTCTTTTAAA
TTATTGCTGCCTACAGACACATAGCTATAGTTCATGCACTTTCAACCACCAATGCTGCCAGGCTAGTAAA
GCAGTTAATGTATATTTGGGGTTAATTATCAGAATCACCAGAAACAATTTTTTAATTTTAAAAATATTTT
ATTTTTCCACAGATTATTGGGGTACAGATGCTGTTTGATTACATAAGTCTTTACTGGTGATTGAGAGAT
TTGGGTGCACCCAACATCCGAGCAGTATACATTATTCCTATG

The following amino acid sequence <SEQ ID NO. 91> is the predicted
amino acid sequence derived from the DNA sequence of SEQ ID NO. 33:

FPRIVCTVTGVAVYHSIYTSIWHTAGASGTTYQSVSLPDHFDVLSYLPCKLVNVYDCFVIMQSCNNNM
YFKNLGIPLHTISSIHINEKSKLGVSVKHWIFTMLIGVPFIIAAYRHIAIVPCTFNHQQCQASKAVNVYLG
LIIRITRNNFFNFNILFFHRLLYRCCLITVLYWFERFGCTQHPSSIHYSL

The following DNA sequence Seq-2617 <SEQ ID NO. 34> was identified in
H. sapiens:

CGGGTTATTTAGAGAACCCTTGAAATACCACCTCCTTGGTAACACCAGCTCCCTCCACCCCTGAGCTCA
CGGTCTCTTCCCTGTGAGATGCAGCACCAGGTAAGGTCAATTAACAACCAGGTTTAGAGTAAACAGTGCTGG
GCTGTATTTCTGATCCTGCCTTTCCCTAACTGGGTGCTCTTTGGCAAGTTATTAAGTTACTTTCATCTGTAC
ATGGGTTACATTTATGCCTTTTACATATGGTTGTTGCGAAGATTGAGTGATATGCATACCAAAATGCTG
AGCAGAACACCTTGTCCATATCTTCTCTCTGTTATTAATGGAGGCCTTAAGGTTAAGTAATTTGTTA
TTGTTGTGGTTAATTTTAGTCCTCTGAATTTAATCTAGTACAAATTGTGCTGCATTGGCACATGGTACA
TGTTTCATGAATATTGAGTGTTGTATAAAGGAATGAAAATCAATTACATGAAAAGAAATCCAAATCTTAC
ATTTTACAAACACAGACACAAAGAATACTAAGATTTAACTCAGGGGCAAAAGTTAAGATTGGCCACCAGC
ACGTGGTGAGCTTCTTGAAAGTTGTTTCTGGC

The following amino acid sequence <SEQ ID NO. 92> is the predicted
amino acid sequence derived from the DNA sequence of SEQ ID NO. 34:

GLFREPLEIPPPWHQLPPPPPELTVSSSLDAPGKVINNVQVSKQCWAVFLILFPFNWVLFGLKLLSYFICTMGY
TYAFYIWLRLRLSDMHTKNAEQNTLSISFLSVIKWRPLRLSNLLLLLWLILVLILILIYKLCCIWHMVHVEHYV
LYKGMKNQLHEKKFQILHFTNTDTKNTKILRGKSDLATSTWASLKVCFW

The following DNA sequence Seq-2618 <SEQ ID NO. 35> was identified in
H. sapiens:

ATTCTAATGGCATTAAATCCATAGCATTATCTCCATCTCTGTTTTAAATATCATGCATCCTCATTCTATGAT
TCAATTTAAAGGAATCCTTCAAAGGATCAATCTAAATAAATAACAAGTTAGCTTTCAGGCAAAACAAATAA
ATTTGCTTTGTTTTATTCACCATAAATATTTCACTTAATTACTGAGGTACCTTGTTCAAGAAACACAAA
ACAACATTATAAATTAATTAGCACTGTCCCTGCTGACGTTTTAGTCCTGTGGAATGCAAAAGCTAAAAGTA
AAAACAGGCCATGAAGCCCAACCAGAGCACATCGTATGCAAATGATAAAGCCACAAACATCATGGGAT
CATTCCTGGGACATTCTGAATCACCAAAATTTGTCTTAAATCAAGTATTGCCCTATTTATTTTCAAATTC
AA

The following amino acid sequence <SEQ ID NO. 93> is the predicted
amino acid sequence derived from the DNA sequence of SEQ ID NO. 35:

LNLKINRAILDRQNFQDSECPNDPMMFVGFIIICRCVLWLGFMACFYFLHSTGLKRQGGQCLIYNNVLC
FLNKVPQLSEIFMVNIKQSKFICLPESLVIYLDSPRIPLNIIEGCMIFKTEMEIMLWINAIR

The following DNA sequence Seq-2619 <SEQ ID NO. 36> was identified in *H. sapiens*:

TTTGGGTCTAGAATCCCTTGGTTTGGGAAGCATCCAGAAGGAGCTTCCATCCCCATCCATTCTTGCTCA
CTTCCTCCTCTCTAGCTTTGTTTACATGTCTCTCGATACCTAGACAGAGGCAGAGGCATGGACTTCTGGTT
CTAATCATTCAAGCCTTACACGTCCTCAAGGCTCCATTGAGAAATATCTTCTCGGGGAGTCTGCTTCT
CCTATTTTCAGGATTTACTGACTATTCTCTTATCTCTTGTAAACATTTAATATCCCACTCCTTAGCATTAACT
TTTAAACTTGCTTTCTAATCCTGAGGTTTGTGTCTCTTGCCTTGTAAAATTCTTTGTAAATGGCCAGCCC
AGTACGTAGCCAGTCCCAAGCACCACGTAGGCAATTGAAGGAGCTGGGACAAAAAGAGTTCTTTGTTTGA
ATTTCTTTTACTGCTCTGAGTTTACTCTGTATTTGCACATGAGTTTAAATGTTTGGGGCCATTGAACAT
TTGAGAATCTAGAAGATAATACACCTCTTTTCAGAAAAACACATATGAATACACACACACATGCCACCT
ACACACACAATTTTGCATGTAATTTTAAGGATTCATTACCTTAGCTTACCAGACTGTAAGTTCCTTTGCA
TATT

The following amino acid sequence <SEQ ID NO. 94> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 36:

YAKELTVWAKVNESLKLHAKLCVVACVCVSYVFFKEVYLLDSQIVQWPQNIKTHVQIQSKLRVKEIQT
KNSFCPSSEFNCLRGAWDWATYWAGHLQRILOQKGTQTSGLSKFKSCGVGYMLQEIRESVNP EIG EADSPR
KDNSEWSLEGRVRLELEPEVHASASVVS RDMTKLERRKARNGWGKLLLDASQTKGILDP

The following DNA sequence Seq-2621 <SEQ ID NO. 37> was identified in *H. sapiens*:

GCAAGTTATCTGTATTTATCCCCCTACAAACACACACTCCTAACATACAGTGGTGAGAGAGGAACAACATA
ACTGCAGAGGAAGTAAGTGAGAGACACAAAGCAGTCATTGGTTCATTGCTATAATGAAATTCCTCTAGACA
AATGCTGCCAGGATCTCTTCCCTGGGGATAAGGTCTAGTTATCTTCTGGAAGTGGTTTCAGCTCACTAT
TCTCTACTGTATAATTACAGTGACTCCCTCATCCATCCTTGTCTTCTCAGATCTTAACTTTATCCTCTA
GACTCCAGGCTCCTCCTCTGAGATGTTCTCACTTTTCTGCAACAAAGCTGAGTCTATTTCTCAATCTGTT
TGCTGTCCATAGAAAATGGAAGGTTTTCAGAGGCTTTTATCAATTTTCTCAGTCTCTTTATTGCAAGCTGGG
TCCCATTTACTTATATACTCTTTTAAAAAGTTTTGTGGGCTTTCTATGTATCAGATAATAGACCACTTC
ATTTGATAAAAAGCCACATTCTTTGTTTTCCAGACAAGCTTCTATATTTTGGACAAGTAAGGCCACTTA

The following amino acid sequence <SEQ ID NO. 95> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 37:

KLSVFIPLQHTHPNIQWERNNITAEVSEERHKAVIGSLNLSRPMPLPGSLPWGGLVIFLEVSSSLFSTVL
QLPHPSCLLSRLSLPLDSRLLLDVLTFLLQQLSLFLNLFVHRKWKVQRLLFNFLSLFIASWVPFTYITLL
KSFCEGLSMYQIIDHFIKATFFVFQTSFLYFGQVRPL

The following DNA sequence Seq-2624 <SEQ ID NO. 38> was identified in *H. sapiens*:

TTATGGTGTTTGTAAAGATCTTATTGCCCAAAGAGTCTGTTCTGTCCATCTTATGATATCTGTTTTAACATT
AATGATGCTCAGTTGTGTCTAGACCTAAAAGAAGAAGTTTGTATGACTTCCATGCTGTTATGGTCAGGA
ATTTAGTTTTTAAGCTTTTTTGGGGCTCTAAGCCACAAGGGGATCTGTTCAAGTCAAGTTCAGTAGAGGGCTT
AGGATTTATCATCTTTAATTCACATTCCTCCCATTTTGGTCAAAATATGCCAAAAGTAGCATCAATAGCCAA
GCTCTTATTTCAATCCATATTATTACCAGGTGGTGTGGCTATCTATCTCAGATATATTCTGTTCTTCAATG
GGACCCATATAGCCAAGGGACTTATAGCCAAAAGACTTACAGCCAATTAAACATTCTAGGACAAAAGGGAA
TGGAGGTGGGAAGGCATTCAATTATTCCTTAAAAACCTTTTGAGCAATATAAGAGCCACAAACCAAGCCA
AAAAGTAAGCTTACAAAACCGATTTATCTATAAGTTCTATGTGTTGGGCCATCGGCTCTTAGGCATCTGTG
AGCCCATCTTTTTTGGAGGATCTGAA

The following amino acid sequence <SEQ ID NO. 96> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 38:

MVFVRSYCPKSLFCPSYDIFNINDAQLCLDPKRRSLYDFPCYGOEFSFKLFWGLATRGSVQSVQRADLS
SLIHIPFFWSKYAKSSINSQALISFHIITRWCGYLSQIYSVLQWDPYSQGTYSQKTYSQLNILGQKGMEVG
RHSLELKNLLSNIRATNQPKSKLTKPIYLVLVCGPSALRHLAHLFWRI

The following DNA sequence Seq-2625 <SEQ ID NO. 39> was identified in *H. sapiens*:

AAGGCAGAGGGGGCCAGCAGGGCGGGTTACAGAACCATGATGTGTTTTAACTGGACTCACTTCTGCCAGT
ATCTGCCTGACTCTTCAGCCCATGCTCTTTTCCTTGTGTGAATACTAATGGGGGCATTAGGAGCCAGAG
AAGGGGCTCCGACGCCACTGCTTGTACCTCTGGAGTTACATTTAGCGGCATTTATATTTTGTGATGTGAA
ATTCGAAATCCTCATCCAAATGCAACTGTGGGGGAACCTCATAGGAATTCAGCCAATTCTGGCTCC

The following amino acid sequence <SEQ ID NO. 97> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 39:

GRGGQGGGLQNHDFLTGLTSASICLTLPMSLFLVVILMGALRSQRRGLRRHCLYLWSYIRHLYFVMNSK
SSSRMQLWGNSHRNFSQFWL

The following DNA sequence Seq-2626 <SEQ ID NO. 40> was identified in *H. sapiens*:

CCCTTCTCTCCAGCCATACCGTGACCCACCCATAAGCTGGCCCCCTTAGCTCTGGCTCACCTGGCTCAGA
CTTAGAGGTGGCAGGATTCTCTGCTCAGGAAATAAGGACTGCTCTTGAGCTCCTCACAGGCCCCAGGAA
TCCCAACAAAGCCAACCAAGGCTACCTTCAGGCTTCCAGAAGGGGGTGGTAGTGTCTCATCAGGTTCC
CCAAGTTTAGGAGAGGGCAGCTGGGCCCAGGGCCCTTCTCCTTGTGGCTCAGGATTTAGCCCCACTTACC
ATGGTGCAGCCCCAGCCTTCCAGCCAACCCAGCATTAGAGGCAGTGGCTCCTCTTAATGCCAGGCCCTAGT
TGGCTCAGGCATAATCCAGCCAGGAAACCTCTACCTTTCCACAGCAATGGCCACCAAGTGTGAAAACGGAA
GCCGACACAGACATGCCCTGCACCAAGCCGCTCATCTTGCATGTGGCATTGTGCAAGGGCCACCCTGCAAT
GACAGAGGGCCCCACAGAGTGAGAGATGCCACGCATCAAGAGCCAGAGACTGAAAGCCCTCCAAGCCAGG
TCCCTCTGAGCTTGGATCTTCTCCTCATGACCTGCTAGGTGTTATCTGGTCTCTGCT

The following amino acid sequence <SEQ ID NO. 98> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 40:

SRDQITPSRSWRKDPSSSEGTWLGGLSVSGSCVGISHSVGSVIAGWPFDNATCKMSGSLVQGMSSVSASFVTL
VAIAVEREVSWLDYAANGLALRGATASNAGLAGRLGLHHGKWGILSHKEKGPSPCLPKLGEPDEDTTTP
FWKARPWLA FVGIPGACEELKSSPYFLSSRNPA TSKSEPGEPELRGPAYGWVTVWLGRK

The following DNA sequence Seq-2627 SEQ ID NO. 41> was identified in *H. sapiens*:

AAACTCCCAAACGATAGTAACTTAAATAACTTAGGTCTTTAATACTCTCTTCAGTAAAAGAATTCTAGTAG
TTGGAGAGTCCACCATCCCTAGGAATGTAGTTCTTGTCTCATGGTTCAATATAGCTGCTGCCAGC
CATACAGCCACATTCCGACAGCAAAATATGGAAGAGAATGAAGAGAAGAAGAGCGTGCTAGGAGTCC
CATGTATTATTTCCATATATATTTGGGCAGAACCTAGTCACAGGGCCACTCCATACGTATCTGTTAGCTAT
TGCTACATAGCAACCACAAAATTTCCATGTATACACATATCTGCAGGTTGGCTAGGGTTCAGTTCTCT
CCATGCTGGTCTCAGACAGGCAGTTCTGCTTCGGGTTACAGTGGCTGAGCTGATTCCATTTCTCACTGCAG
GTCTGTGTTTTCAGTTGAGTGACTGTCCCATGTGCCCTTTCATCTTCTTGGGTTGATGAAAGGAAGCCACAT
CTTTCACAGGGCTAGCCACATCTGTTCTCATGGCCCAAGAGACACCAAGAGCAGATAGAAATAGGTG
AGACCTCTTAAGGTCTAGACTCAAACCTGGCACACTGCCACGTCTGTTTACAAGCTATTAGCCAAAGCATA
GATGCATTACCAAGCCCCAAGTCAAGGGCAAGAAGTACAATCCACC

The following amino acid sequence <SEQ ID NO. 99> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 41:

TPKRLKRLSLILSSVKEFLESPPSLGMLSSWFNIAADAPAITATFQTAKYKRMKRRRACLGVPICISY
IWAEPSHRATPYVSVSYCIATTKFPCHTTHICRLARVQFLHAGLRQAVLLRVTVAEIIPFLTAGLCFSVT
VPCAFHLPWVDERKPHLSTGLATSVPHGPKRHQRADNRDLRLSRLKTGTLPRLFTSYPKHRCITKPVKVG
KKYNP

The following DNA sequence Seq-2628 <SEQ ID NO. 42> was identified in *H. sapiens*:

ACAATAATTTGTTGTATATTCCAAAATAGCTAGTAGTGAATGTTTCCAATACAAAGAAAAGATAAATGTT
TGTGGTGATGCATATTTCAAGTACCTGATCTGATAATTGCACATTGTATACATCTATCAAAATATCAGCA
GTACCTCCAAAATATGCTCAATTATTGTATAAGTACAAAAAATTTAAACAATTATAATGTATTATTTATT
TCTAAATGGTTTATTAGATTTAAATTTCTTGGTGTAAATTTTTTCATATATTACCTTATACCTTTAA
CTTCTAAAATATATTAGTCTTTCATATTTTAGAGTAAATTTCTGAAAATCCTTTGAGTATCTGATTTTAC
AATCTTTTCTTCCACTGATTTTCCCTTAGCAATGGCCTGTTTAAAGTGTGTGATGATGTTACTGAGAAAT
GGGCTGGCTACCTGATGCACATAGAAGCCAATACTATGGCAGTGGTTTCTAGAAAAGAAAAGGCTTTACT
GTGAGTCTACTGGCAAGGAGACAGGTGGCAACACTCAAATCTGTCTCCCTGAAGTGAAGATGGTGGGGT

The following amino acid sequence <SEQ ID NO. 100> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 42:

TIICCFQNSCNVSNNTKKRMFVVMHISSTLILHIVYIYQNISSTSKICSIIIVQKNLNNYNVLFISKWFIR
FKIFLVFNFFIYYLIPFNFLKYIRSSYFRVKFSFEYLILQSFLPLIFPQWPVSVVMMLLRNLATCTKPI
LWQWFSRKEKALLVYWQGDWRQHSNLSPTEDGG

The following DNA sequence Seq-2629 <SEQ ID NO. 43> was identified in *H. sapiens*:

CCTCTTACTTGGGCCCCGTTCACTAGTCCTTCAGCCAACTGCCTCACATGCTATTCCCAGTATGAAAATC
TTGCCATTCCCTTTATCTTTTCTCTCTCTCATTACAGCCCTGTGCTAGTTTCTTCATTCCCTTCAAG
TTCCTGGCCAAACTTTATTTACCTCTTGACTGACCACTCCATCTAAAATAGTACTCATCACTGTGTATCCCC
TCAACACACTTTATAGGTCATGGCCATCACCTGATAATGTGTATGTATTTTTTGGTTTACTTGTGTGT
AGTTCATTCTTGCACTTGCTGTAAAGAAATTCCTGAGACTGGGTAATTATAAAGAAAAGAGGTTTAATTGA
CTCACAGTTCTGCAGGCTGTATGGGAAGCATGTTGCTGGCATCTGCTTGGCTTCTGGGGAGGACTCAGGAA
ACTTACAATCATGGGAAGGTGACGGGGGAGCAGGCACATCTGACATAGCAGGAGCAGCAAGTGAGCAAAG
GGGACGTGCCACACACTTCTAAGTAACAGACCTCATGAGAACTCACTATCATGAGAACAGTACCAGGGG
ATGGTGCTAG

The following amino acid sequence <SEQ ID NO. 101> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 43:

SYLGPVHSFSQTASHAIPSMKILPFPLSFFSSLIYSPVLVSSFPSSSGQTLFTSLTPSKIVLITVYPLNT
LYRSWPSPDNVLCIFWFTCCVSSFLHCCKEIPETGFIKKRGLIDSQFCRLYGKHVAGICLASGEDSGNLQS
WGRRSRRIHSRSSKAKGDVPHTSKPDLMRTHYHENSTRGWC

The following DNA sequence Seq-2630 <SEQ ID NO. 44> was identified in *H. sapiens*:

AGTATAACAATTCAGTCTTACATCTCTATATTTTGCTTATCTCAAGTATCCACTTTGTCTGGTATAGTG
TGCTCATTCCACAGTTTTTTGCTGTCTGGGAACAACAATCTAGTGCAACTCCAGCAATGTGAGTTATAGT
GCAATGTCAAACAGAGCAGCATCACCATCTAGAGGTCAAATGATAACTGCAAACTTTCTCACCTTTAT
GAGCCTTCCGTATTCTGTATACATAGCAGTTTATGTGAATGTACAGAAAATAATGTTTGCTATTGTTTTCT
CTCCAGTTGGGTTTCCAGAAAGAGATCATGGCATAAAGCAGGAACCCTGTATTTACAGATGGCATAGGG
AAGCATACATCGCAGAGCCATATATCAGCAGCACTACAGCATGTTTCAACCAAAGATGAGCCTCCACATG
TCAGACAAACCCTACATTGGGACCACAGCAGTGAAGTGTGTTTTAGCACATTCTGATAATGAAATCT
ATGTTGAACCTCAACATGAATGGCTTTTCTTTCTTGGCAGTCAACAGCCTACACCATTCTGCATTTGAC
TGTTTAGTTTATTCTCCCTCTGGAAGGCATGACTATGGAAACAGAGTAGAGGATATTTTGGGGATTAT
GAACTATTAATATAATTTACTCTCATTGCTGTGCTTTCTACAA

The following amino acid sequence <SEQ ID NO. 102> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 44:

YNNSLYISIFCLSQVSTLSGIVCSFHSFWLSWEQSSATPAMVIVQMSNQSSITIRSKLQTFSPLAFRIL
YTQFMMYRKCLLLFSLQLGFQKEIMASRNHLYLQAGSIHRRAIYQQHYSMFQPKMSLPHVRQTTYIGTTA
VTVFFSTFLIMKSMNLNSTMAFFFSWQSTAYTILHLTVFILPSGKALWKQSRGYFGDLNYYNLLSLCLFLQ

The following DNA sequence Seq-2631 <SEQ ID NO. 45> was identified in *H. sapiens*:

TCAGCTTATCTGGTCAATAGCTTTTCGCTCTGTTGCATACCTTGAGCATATGCATCAGCTACAATGTTTAT
AGGTAGCTGTATGGTGTGTTGACACAGCACATGGCGTACCTTTAAACAATTATAGCACTGGGATTTGGATC
TGAATTTATGTTGCCTTGTCAAAGTTTCCTCTTTGTAACATGGTAGCCTTTTAAATATTAGGCAGCTACCT
GCAACACTGGGCATTCACTAACCATCAGGCTTATGGCATCTTGCTCTTCTCGTCCCTCTCTGTGTGT
TGGTACATCATGTTAGGTTTATGCAGTAGACGTAGATAGGAAGCAAGCAATTGGCTACAGGTTATTGAAA
GTCAATTGCTGAGAATGATAAAGACAAGGATAGCCTTCTCTGCAAAGAAGTGCTAAGAAGATTCTAAACG
TATACAAGGATCTCAAGAGAAACAGTCCAGATAGCAACTATTCACTCTTAGACTATGGCTGATACTAT
ACACTTCTCCAGCTCCTCTGCTCCTCAGAGCAGAAAACAGAGATTTTGAAATGAGCACCACCCAGCTCC
TGAATACAATGGTACCTTTTCTATTTCTGTTGACTTTTATTTCTTTTGTGCTGGATCCCTACATAA
TTGTAAGCATATCGCAGGCAAGCACAATGGTAAACAGTGGGTGGACGCTTCCTC

The following amino acid sequence <SEQ ID NO. 103> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 45:

SLSGQLFALLHTLSICISYNVYRLYGVHSTWRFTKIIALGFGSEFMLPCQSFLFVTPFKYAATCNTGHS
DPIRLMASCSSRSLSVCWYIMLGLCSRRREASQATGYKSIAENDKRQGPSLQRSARKILNVYKDLKRNSP
RQHYSVLDYGYITLLQLCSSEQKTEDFEMSTTPAPEYNGTFHFLVTFIFCCWIPYIIIVSISQASTMVN
SGWTLP

The following DNA sequence Seq-2632 <SEQ ID NO. 46> was identified in *H. sapiens*:

ATAGAAGCTTGATTATATTGGTATTTTATTTCAAATTTTCAATTTTGGAAATGGCAGAATGTTGCTATTGA
AAAGTGTCTTAAAGGTCACCACTGTAACCCCTTCATTGTGCTTGAGACCTGCTCAGCTCCTAAATTTAACA
GGGGACGGATCTGAGAACTGACTCCAAGTTGTAACCTCTTGCTAGTTTTCTTTCTAGGGAGATATCCGT
CTCTCCAAACCTGTCGAAATCTAAATTTATTACCTCTTACCTAATACTTGGTCCCCTGTGGACTTCACCTC
ACTGTTTGTGCTAATAGCCTTTTTCATCACCCTCTTGACTTTGGATCTAGAGCATCACCTACTTCCCATT
TTCTGTGACCCCTTACATTCCTCCTGTGCTGACTATGCTGATTATTGTTCTCCCCTATCTTTTGCCCTT
TGCAAATCCTCAAGCCCTCATTCTGGTTGAGACCTTTAAAAGGCTGAGTTACTGGAGTATAGTGTACCCA
AAGTGAGTTGTTCCATAAAAAATAGTAAGTTGGAAAAAAAACAAAAACAAAAAATACCCCTACCCATA
AAGTTGGTAAATGTTCTGTAAAAAGGGTCTTGGCCAGGTACATGTTAGAATAGCTGGTTAAGTTCTT
TGCAGAAAGACTTCTCCTGGCCTTCATTTGTGACTGTG

The following amino acid sequence <SEQ ID NO. 104> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 46:

RTLYWYFYFKFSIFGMAECCYKVSRSPLPLHCADLLSSIQGTDLRLNQVVTSLVFFLGRYPQLQTCRNLN
LLPLTYLVPCLHFTVCANSLFITILTLDSTRASPTSPFSVTLTFLLSVTMSDLLFSPIFCPLQILKPSFWF
RPLKGVTVGCYPKVVPKISKLEKTKNKKIPYPSWMFLKGLQVHVRIAGVSLQKDFSWSFVTV

The following DNA sequence Seq-2633 <SEQ ID NO. 47> was identified in *H. sapiens*:

GCAATTAAGTTTTGTACTGTATGGACAGTGTGAAAAACATTATGGAAAAACAACTTGAAGAAAATGTGAC
AGAATTTCTCCTAACATGTCAATGCTTCAACAGCTACAAATTTCAACCTAGTTTCTTTCTTTGCTGT
TTCTTCTTTTGTCTTTGATACAATCATAACGCTCTCTCCTTGAAGAGATAATAAAGACTAACAGTTAA
AAGATCTGGAAGACTCATATTCTTTCTTTTCACTGGCTACGGTTTTGAAAAGAGGCTGTTGGCTTTTGA
TTTTTCTTTTGGGTTCTTTACATCGCCCAATTCAACAGGCTCTGCTCTCAAAGAAAACAAATCAAAATTG
TCAAGACCTGTGAAGCATGAAAAATAAATTGCTTTTCCAACTCCAAAAGCACCAGAAAAGCATTAAATTT
TGATCTTTTATAAACCTCTATCCCTATCCTCTAATCTATAGATTTTACAGAATGTTTATATATTCTTCTG
TATAATACAGGAGATCAACCTTATTATGAATAAATTGAATTGAACCTGTAATACAACTAATATTTAACT
AGTGTTATTTTGGAGTTCAACTAGACACATATAAACATTTCAAGTGAGATGACACAAATTCCTGGGGCTG
CCAGTATAAAATAAACAGTCCAGTAAGCTGCATCTACCATGCCGTTAAGGGACTCTGCTCTTTAGCTGGT
GGGAGCACAGGCTTCATAA

The following amino acid sequence <SEQ ID NO. 105> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 47:

MKPVLPAPAKRTESLNGMVDAAAYWTVYFILAAPGICVISLEMFYMCLEVELQNNTSLNISCTGSIQFIHNKV
SPVLYRRIYKHSVKSIDRIGDRGLKIKINAFVLVFGVGKSNLFFMLHRSQFFVFESRPVIGRCKEPRKN
QKPTASFQNRSQKRKEYESSRSFNCSFIISRKRGMIVSKTKEETAKERNVGNLLVEAMTLLGEILSHFL
SSCFSIMFFTLISIQYKTL

The following DNA sequence Seq-2634 <SEQ ID NO. 48> was identified in *H. sapiens*:

TCCTGAGAAGACCTGCAGCACAGGGTAAATATGCAAGGGAGGGCCATATACTTTTATCTTTACTTAATT
TATTTTAAATTTACTAATTTTAAAGTATTAACCTATTTTGTTTTTTATTAAATCTCTGTGGTGCACAGAATT
CAAATTGCAGCAAAATCATTACAGGGCTAAACACTGGAAAAATCTCTTAATTCTAAGGTACATGACACAAT
GGACTCAAAAACAGTTGCTGAGTCCCTTTCACTGGAGAAATTTAAAGAAAGGGTATAGAAAAGTTTGACC
AATCCACCCAATCCTGCATCCCCAATTCCAATCTCAAGGACCAAGTTTCCATCTGATCTCTCTCCACCTAC
AGATGGTGGTCTGAATCTCCAATCAACAAACCAAACTGAATCCATCATCTTCTCACACCTGGTTTTT
CCTTCCAATCCCTCATTCTGTGACCTGCCCCATAACCTTACCAGGAATCCAGCCCCCAAGCAGGGTGG
ACTCCTCCTCTGCAATGGACACCAGGGATTACAGGTCTGTTGCTGGCTCCAAAATGCCCAATGCCCTG

TTCTCCCAATCAGCACATTCAACAGT

The following amino acid sequence <SEQ ID NO. 106> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 48:

SEDLQHRVKYAREGHITFIFTFILIYFLSINLFCFYISVVAQNSNCSKNHSGLNTGKISFGTHNGLKNSCV
PFTGEIRKGIEKFPIPPNPASPIPIISRTSFHLISLHLQMVVLNLQINKPKTESIIFSHLVFSPNSLISVTC
PITLPGIQPPKQGGLLPLQWTPGIQVLLAPKCPQCPVLPNQHIQQ

The following DNA sequence Seq-2635 <SEQ ID NO. 49> was identified in *H. sapiens*:

TTTGACTTAACCTTTGTAGCCAGGTAATAAATCCAACTCAGCAAGTATGGGCTGGACCCAGTAGCTC
TGTGGTTGCCACTTTTTGGCCCATATTGAACCGACGTCCCCTTGGCATCTACCAGGGACTCCTCAGGGAGA
GTGTGGGAATGATGGGGGAAGACTCGTCACTCTTTGTAGAGCGTGGGGCAGATGATAGCAGAGACCTTCC
AGGGCCAGGGCTGGGGTCTTGTCTTCTTGGATGTGGTCTAGCGTTGCTCCAGATGGTGGGTTTGTGGCA
GGTGGGGCAGAAGCAGATGATGCAGTTGAGCGGGTCTCTGGTAGAGAGTGATGTCAAAGATGAGCACTCC
TTTTATCCCTGACTCTTCTGAGGATGGCTGCCTCCTTGGTGAGCCACTTGGAGGTCTCAGGCCGATCATG
CGGGATGGTGGCCAGATGAGGAAGGGGATCCAAGCGGTGGCCTTCCAGATGCACTGGGCCCCAGCCCT
TCTTCTAGCTTCGGCTGATTACTGTGGGCTTCAGCAACCAGGGCCTACCTGTAGGTCTCCACATTGTAA
GCACCACAGAACCAGTGCATCTTTGTGCACTCAAATCGCCCTTGATCCAGGGGTATTTTCTCATTAGAA
CACACTTTGAAAGGCGCCATTCCCTTCTGGAGAAAGCCTGGAGAATCTACAGTGCCTTAATTACAGTG

The following amino acid sequence <SEQ ID NO. 107> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 49:

HCNGHCRFSRLSPERGEWPPFKVCSEENTPGSRAIVHKDALGSVVLTVETRYALVAEHSNQPKLGRRAGA
QCIWEGHRLGSPSSSGPPSRMIGLRPPSGSPRRQPSSEESGDKRSAHLHSLPETRLNCIICFCPTCHKPT
IWSNARPHPRKTRPQWALEGLCYHLPHALQKSDDESPPIPTLSLRSPWMPRRFRFMGQKVATTELLGSS
PYLLSLDLLPGLQRVKS

The following DNA sequence Seq-2636 <SEQ ID NO. 50> was identified in *H. sapiens*:

AGATGCCCAGACACCTTCACTTCAAGCAGACAAGGGGCAGAGTCCCTGGAAATCTAGGCAGGGAAGACTTGC
GCCTCTAAGAGTAAAAGGCCTCCAGAGAGGACATGGATGAAAGGAGGACCCTTCCAATGCCACTCTCC
AAAGCAGGAAACATCCAAATAAAGGATGTTGATTTTCAAGACCCCATCCCTTCATGAGTGCTTACACAAC
GGTATATCCTCTCCCGTCTCTTCTCTGGTAGCCAAGACCTTATACCAGTTTGAGTATCCTTTATCCAAAA
TGCTTGGGGTCAGAAGTGTTTTGAATTTAGATATTTTAAATTTGGAATATTTATATCATACCTCTTGG
TTGAACCTTCCAGATACAAAATCTGGAGTCCAGTGAGTATTTCTTTGAGTGTCATGTAGTGCTCAAAA
AGTTTTAGATTTTGGAGCGTTTCAAGTTTCAAGGTTTTGAAATTGGAATACTCAACCTGTACTCTCTGTCC
TTGTTCTACCTCTACCAGACCTCCCCACAGGAATGAATTTAGATCTGAAA

The following amino acid sequence <SEQ ID NO. 108> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 50:

FRSKFIPVGEGLVEVEQQRVQVEYSNFKNLKSETLQNLKLFHHDTQRKYSLSRFLYLEGSTKRYDINI
PKFKNINSKHFPQAFWKDTQTGIRSWLPEEETGEDIPVVALMKGWGPENQHPLFGCFLLWRVALEGGPPF
IHVLSGRPFTLRGASLPCLDFPGLCPLSAEVKVS

The following DNA sequence Seq-2637 <SEQ ID NO. 51> was identified in *H. sapiens*:

TCATCCTCCGCTGTCTATTTTGTAGCTGTGAGTTTATCCACAAAGGAACAGAGCTGAAATGAAACAATTTCA
CCACAGTAACTTGTTAATCGGGCATCCTTTAAGTATGCTGGATTAACACTGGAAGTTCTTTTGAAGACTC
TGAAAGTTTCTTTAATCGTCATGAGATTTTCCAACTAAGTTCATGATATGGATTTTTTCACTGTATC
TAGCTTAAGTCACATTTCAATTCAAATCTAAACCTAACTGATGGAGCTGGAGCTAGTACTTCAGGCAAT
TGGCATCTTTTCGCTGAATACAAACATCCTATTTAAAGACCAACACATGACTCCATTCAAAAATTA
CAGTCATGTGTAGTAAACAGCAAGAACCGGTCTGAGAAACGTGCTTGCACACACAGCGTGAATGCAC
TCACGCAAGCCTAGACCGGTGCGGCTGCCGCACACAGGCCCTGTGGTACAGCCTGTCAATTCAGGCCCA
AGCCTGCATACCATGTTGCTGTGCGGGACGCTGCCGGCGGCTGTAGCACATGCTAAGTATCTGTGTATCT
CAACACAGAAGAGGTAGAGTAAAGTACAGTATTATGATCGTACGGGACGCGTGTGTACACACAGTCTATC
ATTGATGGAAGCATCGTTATATGGCACATTACTGCACTGTAAAAAGACACCAAACTTCGGCCGGCGCAGTG

GCTCATGCCTGTAATCCCAGCACTTTGGGAGGCTGAG

The following amino acid sequence <SEQ ID NO. 109> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 51:

SASQSAGITGMSHCAGRSLSVFSYSAVMCHITMLPSMIDCVYNTRPVRSYCTLLYLFCVEIHRYLALCYSSRR
QRPAAQOHGMOAWGLELTGCTTGPGVROPRLGLRECIHAVCARTRFSDRVLAIVSLHMTVLIFEWSHVFGLL
NRMFVFSEKMPIASHLQLHQFRFRFELKCDLSIQKKSISTFGKISRLLKKTFRVFKRTSSVKSSILKGCPIIN
KLLWNCFISALFLCGTHSSKTAED

The following DNA sequence Seq-2639 <SEQ ID NO. 52> was identified in *H. sapiens*:

TTCTTCCTTTTCTTTTCTTATCATTTTCTTTTGTCTCAAAATAATGAAAAATGCATAAGGGTCTGTAGA
GAGAAGAAAAATGTCCTTGCCCATGAACCTTCTGCAGGTATTTATCTTGCTTCTTATCTTACTAAAAATAG
AATTGAAAGTTTTTCAATTTTGTGTTTCAATTTAGAGGATACAATGGAGATTCAGGAACGAATAGAAAA
TAGTTTAAAGTCTTTACTAGACCAGTTAAAGGTAAGTTTCTTACTGTAGATTCTGTATTGTATCTGG
TTGTATGGCAATAGCTTCGAAGTTCTTTCCCTATTCCCAAGCCCAATCACCCAGAGATAAGTAAGTAGTT
TTAAACACTTTGGAGTCAATACTCCTAGATGCCACCTAAACACATATGTGTGTGAATGAAAAATACAGATAAA
AAGTAATCTTTAAACATAGGAAATGGTGAATCCATGCTTTTTGACTTTAATTTTTTGTATTGTTGGAT
ACCTTTCCATGTCAGTTATATATACCCATTATTTTCAAGACTGCGTAATATTCTATAGTATTGTATTAA
CATTTTTTATGTTATCGCAATTGGTGACATATTATGTATATGAGTTATTTCTTCTACTGATGCTGAAATGA
ATATCTTGGGACAAATTGTTAGGGGTATTATTGAGTCTTCTTGGGATTAAATT

The following amino acid sequence <SEQ ID NO. 110> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 52:

FFFLSLSFSFCLKIMKNAGSVERRKPCPTSCRYLSCFFILLKIELKVFFHFLFFNFRGYNGDSGTNRKFV
FTRPVKRVFLIPVVFVSGMAIASKFFPLFPSPITQVRSSFNLTLESILLDATTHMCVNENDDKKSLNIGNG
VIHAFLTLIFLLEWIPFHVSYIPIYFQDCVIFYISIVLTFMLSQLVITYYVYELFLLMLKISWDKLLGLVL
FESEFLGIK

The following DNA sequence Seq-2640 <SEQ ID NO. 53> was identified in *H. sapiens*:

CTTTTGAGGATTAAAAATTCCTGCTTACTGTGCTTATAACACGGGGATTAAATAAGCACCTTACTGGAATCT
CTCACCTACCATAATTTTAGTATGCTATGTGAGGGAATGAACAGTCTCACACATTAAATAAGTACTACTCA
TATAATGCTTTTAAATTGGTAATGACCTATATGAAACATGATATAGAAAAACATTACAGCTTCTCAAATGA
CCCCATAAGTTAACCAATTGCTTAGGTTTCTGACAAATTTGAATCTGGCCCCATGCACCTTGTCTGGGCC
CCACAAAACAAGGAGGTAGATTATTTATGAAGGTCAACCACTCTGGCAATATCACCATTAATATCAAGCT
CATCTGCCCCATAGCTCCTCCATCTTCAGGTCCAGGACTCTGGATTGGAATGACCTACCTCCACATTCACT
TCTGTAAGTCTATTAGGCATCATCAAGATGGTAGATGATAAATGGACAATGACTTAAGCTTTTTTTTA
CTCTCTCATCCATTCCAATGCTTTCTTCCCTGGTCTTTGCTCATTATTTCATGTTATTTAATATATATTT
GGAAGAATTCATGGCAGTGATAACAATAATGGCTACAATTTTTTATTACCTATGTATGCCAGGCATTGTGC
TAAGTGCTTCAGGTATAAGATCTTGTAAAGGATTGGTTACATTTTACAGATGGTAAGACTGGGATTACAGAT
GTTAGTTGCCTGTTTAAAGTCAATAA

The following amino acid sequence <SEQ ID NO. 111> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 53:

FEDKFLTLTVITRGLISTLLESSTYHNFSMLCEGMNSLTHLIMTTHIMLLIGNDLYETIRKHITASQMTPI
SPIAVSDKFESGPMHLWCWAPQNKEVDYLRSTTLAISPLNLKICPIAPPSSGPGWLWIGMTYLHIQFCKSLG
IIQDRINGQLKFLFLSHPFQCFPLWSLLIISMLFNIYLEEFMAVITIMATIFYLCPGIVLSASGIRSC
KGLVTFYRWDWSDVSCFLFKSI

The following DNA sequence Seq-2641 <SEQ ID NO. 54> was identified in *H. sapiens*:

CTCTTCTCCCTAGGTGGTTTGGCTGGCAATCTTTGGCATTCTTAGCTTGTGGAAGTATCACTCCATCTCTG
TCCTGATTTCTACATGGTGTCTTCTGTGTGCATGTCTCTCCAAATTTCCCATTTTATAAGGACACA
GTCATACTGGATTGGGGCTCATTCTAAAGACCTCATTTAATTTAATCCATAAAGACCCTATCTCCAAATA
ATGTCACATTCTGTGGTACTGGGGGTATGACTTAAACATATAAATTTAGGGAGACAAATTTGAACCTCT
AACAGTACTGAACATCCAGGATGGAAGAATGGTATTAGGTTGAGCCAAACACAGTTGCTTACGTTTGGG

TTTTCCTCACCAGGACAAGAAACCCCAAGTGCAGGAAAATTGGAGACATGGAAACAGGGCTTAAGTAAACA
A

The following amino acid sequence <SEQ ID NO. 112> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 54:

SSPVVCWQSLAFLSLWKYHSISVLISVLCVCLQISPFYKDTVILDSGSFRPHLIFHKDPISKCHIL
WYWGLLKHINFRETNLNLQYTSRMEEHGIRLSQTQLLTFWFSSPGQETPSAGKLETWKTGLKT

The following DNA sequence Seq-2642 <SEQ ID NO. 55> was identified in *H. sapiens*:

TTATTATACTCAACACTGCTAGGAAAGAATCAGTGATGTTGAAGATATATATATATATATTGCTTGTGTA
TTTGTGTGTGAGAGACACACATAGAAAAAGAGAGAGAGAAATATATTGGTTGACACTGGCTTCTTTGAA
AAAAGGCAGTTTAGTAACAATGGCCTTTACTAGACAGACATGTTAGAAGGCAGCAGGAGAAAGGGAATGTG
GTATCAGATATTTTCTGTAAAAGGTTGTTATTAATATTCATGTGGCAAATTGTAGCTGATGTCAAAGTAG
TTATAAAGCAAGGGGAACACAATTCTTTTACAGCAATGTTGAGGTCTAAGAAACATAAAACAAATACCTGG
TAAGTACCATGCATATATACATACATAAACAACTCAATACTCACAAAACATTACATATTTGCAACACTGC
TTTTAGTTTATGCAGTTTATTTTTTGTCTTTTAAAGCTTTTATTATAGTGAATGTCTTATATTTTCATT
AAAAGTTTGTATATTATATGTGAACAACAGTTCTGATAAAGCAATATCTAGATAAAGGCTATTACTTACC
TTTCTCAAATTGATAGATTTTCTCCTTGTAACAAGCTCTGATATAAAATATGATAATTTGTGAAAACCTT
TACACATTCAAACATAAATTATCATATATTTAATGAGACTTTGGGTGTGTATGTGTGAGTGTGTCTGTG
TGT

The following amino acid sequence <SEQ ID NO. 113> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 55:

HTDTHSHIHTQSLIKYMIIFMCKSFQIIIFYIRACYKEKIYQFEKGKPLSRYCFIRTVVSHIISKLLMKY
KTFTIISKLRKTKNKLHLKSSVANMMFCELLIVYVCIYAWYLPGICFMFLRPQHCKRIVFPLLYNYFDI
SYNLPHEYQTFYRKYLIPHSLSPAAPHVCLVKAIVTKLPFFKEASVNQYISLSLFFYVCLSHNTQANIYI
YIFNITDSFLAVLSII

The following DNA sequence Seq-2643 <SEQ ID NO. 56> was identified in *H. sapiens*:

AAAACCTGGTTTTTTAAAGCAAACACAGAAACAATGTAATATAGGTCTTATTACATATGTAGGAAATAAAA
ATAATATGTATGACGACACAGTAGTCTAAAATTTCAGGAGACAGAGAATGGAAGTACATTGTTGCAAGGTT
TTCTAATACACATGTACAAAGTGGTATAATGTTACTTGAAAGATAACTGTGATAAGTTAAAGACGTAATCA
ATGACACTATATCAACCACTAAAATAATACAACAAAGGATATACGAAATATTTTTAAAGTATAATTAACC
CAAAAGAAAGCATAGAGGAAAAAGGGAACAAAGAATAATAGATGGAATAAACAGAAAAACTAGCCAGCTG
GTAAATTTAAACCGATCATATACATATTCACATTAAATACAAAAAGTTTAAACACTTCAAAGTCAAGTCA
GAGGTGTCTATTTGGATAAAAAGAAAGACTCAACTATATGTTACCTATAAGGAATGCACCTTAAATATACA
AACATATTTAAATAAAAAGATGAAAAGTTATATACAATGTTAATACTCATCAAAATAAAGCTAATGAGGCT
ATATTCATATTTAAAGTAGGTTTTAAAGCAAAGATTAC

The following amino acid sequence <SEQ ID NO. 114> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 56:

SLNLLFNMNIASLALFVLTYITFHLFILICLYISAFIGNILSLSFYPIHLLDFEVFKLFVFNVMYMI
GFKFTSWLVFSVYSIYYSLFPSSMLSFGLIILLKIFRISFVVLFWLICHRLRLITVIFQVTLYHFVHVYK
TLQQCTSILCLLNFRLLLSSYLFLFPTYVIRPILHCFVCVCFKKPSF

The following DNA sequence Seq-2644 <SEQ ID NO. 57> was identified in *H. sapiens*:

TCAAGTCCATGCTTTTACGGAAAGACCCCAAGTTCCTGCCTCTTCTATATATTTATCTACCTTGTGGTGAAG
AGCATGTGTGTGCAACACCTTTGCCTGAAATGGTATGGTTTGGCATTAAATGAATTGTGGGTCCATTGAAAA
GAAATCTCCTCTTGTCTCGTGTATGGACAGTTCAAGGTTTGCCTTAGAACTAACTTCAAGGAAAAGTA
GCAGAATCGTAGGAAGGGACAATCTTGCCCTCAGTCCCACCTCTGTTCCGGGCAGGTCTGGGTGGCTATC
TTCTTTCCGGGGCTTTTCTTGCAGAAGAACTCTTCAGCATGTCCTGGATTTCCTTCTTAATGGTCTTGT
GCATGTAGCCATAGACATAGGGGTGGATGCAGCACTGCAGGAAGAAAGCCAGATGATTATGGTGATCACC
CACTGGGGTACCTGGGTTTCGACATCCACCCACACGGCCAGGACTGCTAAAAGCAGTAGGGCCCCAGGGA
TAGCACATAGGAGAAAATGATGATGAAGATCACTTTAGCAGCTTTGCACTGGTAGCACCTGGGCAGAGGAG
GGTTGCTGTTGCTGTTACGACGACTGGGTGGGAGGCTCTCCGGGATGTTCACTGCCTCGACGTCATCTCA

CTGAAATTGATGTCGTCTTCACCAAACCTCCATGTCATCTTCACCCAAGTCAATGCTGCACTGGTTGACCTC
TGTGCGACCCCTGTCTGCCTTCATGCTGTTCTCCTC

The following amino acid sequence <SEQ ID NO. 115> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 57:

EENSMKADKGRTEVNVQCSIDLGEDDMEFGEDDINFSEDDVEAVNIPESLPPSRNSNSNPPLPRCYQCKAA
KVIFIIIFSIVLSLGPYCFILAVLAVWVDVETQVPQWVITIIWLFLLQCCIHPIVYGYMHKTIKKEIQDML
KKFFCKEKPPEKEDSHPDLPGETEGGTEGKIVPSYDSATFPSFGKPTVHNTRNKRRLFNPGPTIHCQTIPFQA
KVLHTHALHHKVDKYIEEAGTGVFPKHGL

The following DNA sequence Seq-2645 <SEQ ID NO. 58> was identified in *H. sapiens*:

AGTGGAAAGACCACACCTAGGAACCGACTCTAGCTCTTACCACCCTGTAAGCCTGAGGCTCAGTTGCTGTC
CCTGGAGAACAGAAAACATAATCATGGCTATTCTGAGGGTCAGGGGCAAGTGCTTTGCAAGTGGGATTGTG
GTGGGCAGTGGGAGGGATTCTGGGGTTCAGTGTCTAGTTGTGTAAGTGGGCAATGCAACCGTGTAAAG
TGTGAGGAAACCTCAATAAGACTGAGCCAGAGGCCAATAAGAAGCCAGCATTTACATGATGTTCTTTTCC
TTTTTGTAACTAGGAAATTCGATTTGCACACTGATTTGGCCACCATTCTGGAGAGATCTCGTGGGATG
TCTCTTTTGTACTTTGAACTTCTTGGTGCCAGGACTGGTCATTGTGATCAGTTACTCCAAATTTTACAG
GTATGTTTTTCTGCAAGTGCTGCCACTGAACCTCACCCAGGCTTGGGGTTATTTCTGCTAGAATCTTAGAAT
TTGGGGTCGGAGAACACCTAAGAGTTACGCCAGCTCAATCTTGATTCACTGCCAGGTCTACAACACTGA
GGAAGGAGAGGATTTTTTTAGAAGTTATATCTTTGTGATTATGTTTTTTGCTCATCACTAAAGTAATACT

The following amino acid sequence <SEQ ID NO. 116> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 58:

SGKTTPRNRLLLPCKPEAQLLSLENRKHNHGYSEGQGVLCCKWDCGGQWEGFWGSLSCLCNWMQPKCKQ
ETLNKTEPEANKKPAFTCSFFFCNEISICTLIWPTIPGEISWDVSEFTLNFLVPLVIVISYSKILQVCFL
QVLPLNFTQAWGYFCNLRIWGRRTPKSSRQLNLDLPRSTTLRKERIFLEVISLLCFLITKVI

EXAMPLE 2: CLONING OF nGPCR-x

[000239] cDNAs may be sequenced directly using an ABI377 or ABI373A fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM Ready Dye-Deoxy Terminator kit with Taq FS polymerase. Each ABI cycle sequencing reaction contains about 0.5µg of plasmid DNA. Cycle-sequencing is performed using an initial denaturation at 98°C for 1 min, followed by 50 cycles: 98°C for 30 sec, annealing at 50°C for 30 sec, and extension at 60°C for 4 min. Temperature cycles and times are controlled by a Perkin-Elmer 9600 thermocycler. Extension products are purified using Centriflex gel filtration (Advanced Genetic Technologies Corp., Gaithersburg, MD). Each reaction product is loaded by pipette onto the column, which is then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B table top centrifuge) at 1500 x g for 4 min at room temperature. Column-purified samples are dried under vacuum for about 40 min and then dissolved in 5µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples are then heated to 90°C for three min and loaded into the gel sample wells for sequence analysis by the ABI377 sequencer. Sequence analysis is performed by importing ABI373A files into the Sequencer program (Gene Codes, Ann Arbor, MI).

Generally, sequence reads of 700 bp are obtained. Potential sequencing errors are minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers at different locations until all sequencing ambiguities are removed.

[000240] To isolate a cDNA clone encoding full length nGPCR, a DNA fragment corresponding to a nucleotide sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58, or a portion thereof, can be used as a probe for hybridization screening of a phage cDNA library. The DNA fragment is amplified by the polymerase chain reaction (PCR) method. The PCR reaction mixture of 50 μ l contains polymerase mixture (0.2mM dNTPs, 1x PCR Buffer and 0.75 μ l Expand High Fidelity Polymerase (Roche Biochemicals)), 1 μ g of 3206491 plasmid, and 50pmoles of forward primer and 50pmoles of reverse primer. The primers are preferably 10 to 25 nucleotides in length and are determined by procedures well known to those skilled in the art. Amplification is performed in an Applied Biosystems PE2400 thermocycler, using the following program: 95°C for 15 seconds, 52°C for 30 seconds and 72°C for 90 seconds; repeated for 25 cycles. The amplified product is separated from the plasmid by agarose gel electrophoresis, and purified by Qiaquick gel extraction kit (Qiagen).

[000241] A lambda phage library containing cDNAs cloned into lambda ZAPII phage-vector is plated with E. coli XL-1 blue host, on 15 cm LB-agar plates at a density of 50,000 pfu per plate, and grown overnight at 37°C; (plated as described by Sambrook *et al.*, *supra*). Phage plaques are transferred to nylon membranes (Amersham Hybond NJ), denatured for 2 minutes in denaturation solution (0.5 M NaOH, 1.5 M NaCl), renatured for 5 minutes in renaturation solution (1 M Tris pH 7.5, 1.5 M NaCl), and washed briefly in 2xSSC (20x SSC: 3 M NaCl, 0.3 M Na-citrate). Filter membranes are dried and incubated at 80°C for 120 minutes to crosslink the phage DNA to the membranes.

[000242] The membranes are hybridized with a DNA probe prepared as described above. A DNA fragment (25ng) is labeled with α -³²P-dCTP (NEN) using Rediprime random priming (Amersham Pharmacia Biotech), according to the manufacturer's instructions. Labeled DNA is separated from unincorporated nucleotides by S200 spin columns (Amersham Pharmacia Biotech), denatured at 95°C for 5 minutes and kept on ice. The DNA-containing membranes (above) are pre-hybridized in 50ml ExpressHyb (Clontech) solution at 68°C for 90 minutes. Subsequently, the labeled DNA probe is added to the hybridization solution, and the probe is left to hybridize to the membranes at 68°C for 70 minutes. The membranes are washed five times in 2x SSC, 0.1% SDS at 42°C for 5 minutes each, and finally washed 30 minutes in 0.1x SSC, 0.2% SDS. Filters are exposed to Kodak XAR film (Eastman Kodak Company, Rochester, N.Y., USA) with an intensifying screen at -80°C for 16 hours. One positive colony is isolated from the plates, and re-plated with about 1000 pfu on a 15 cm LB plate. Plating, plaque lift to

filters and hybridization are performed as described above. About four positive phage plaques are isolated from this secondary screening.

[000243] cDNA containing plasmids (pBluescript SK-) are rescued from the isolated phages by in vivo excision by culturing XL-1 blue cells co-infected with the isolated phages and with the Excision helper phage, as described by the manufacturer (Stratagene). XL-blue cells containing the plasmids are plated on LB plates and grown at 37°C for 16 hours. Colonies (18) from each plate are replated on LB plates and grown. One colony from each plate is stricken onto a nylon filter in an ordered array, and the filter is placed on a LB plate to raise the colonies. The filter is then hybridized with a labeled probe as described above. About three positive colonies are selected and grown up in LB medium. Plasmid DNA is isolated from the three clones by Qiagen Midi Kit (Qiagen) according to the manufacturer's instructions. The size of the insert is determined by digesting the plasmid with the restriction enzymes NotI and SalI, which establishes an insert size. The sequence of the entire insert is determined by automated sequencing on both strands of the plasmids.

EXAMPLE 3: SUBCLONING OF THE CODING REGION OF nGPCR-X VIA PCR

[000244] Additional experiments may be conducted to subclone the coding region of nGPCR and place the isolated coding region into a useful vector. Two additional PCR primers are designed based on the coding region of nGPCR, corresponding to either end. To protect against exonucleolytic attack during subsequent exposure to enzymes, *e.g.*, Taq polymerase, primers are routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target.

[000245] PCR is performed in a 50µl reaction containing 34µl H₂O, 5 µl 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5µl 15mM MgSO₄, 2µl dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 3µl genomic phage DNA (0.25µg/µl), 0.3µl Primer 1 (1µg/µl), 0.3µl Primer 2 (1µg/µl), 0.4µl High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes; followed by 25 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3 minutes.

[000246] The contents from the PCR reaction are loaded onto a 2% agarose gel and fractionated. The DNA band of expected size is excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA is precipitated with ethanol and resuspended in 6µl H₂O for ligation.

[000247] The PCR-amplified DNA fragment containing the coding region is cloned into pCR2.1 using a protocol standard in the art. In particular, the ligation reaction consists of 6µl of GPCR DNA, 1µl 10X ligation buffer, 2µl pCR2.1 (25ng/µl, Invitrogen), and 1µl T4 DNA ligase

(Invitrogen). The reaction mixture is incubated overnight at 14°C and the reaction is then stopped by heating at 65°C for 10 minutes. Two microliters of the ligation reaction are transformed into One Shot cells (Invitrogen) and plated onto ampicillin plates. A single colony containing a recombinant pCR2.1 bearing an insert is used to inoculate a 5ml culture of LB medium. Plasmid DNA is purified using the Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the sequence, a 50 ml culture of LB medium is inoculated with the transformed One Shot cells, cultured, and processed using a Qiagen Plasmid Midi Kit to yield purified pCR-GPCR.

EXAMPLE 4: HYBRIDIZATION ANALYSIS TO DEMONSTRATE nGPCR-X EXPRESSION IN BRAIN

[000248] The expression of nGPCR-x in mammals, such as the rat, may be investigated by *in situ* hybridization histochemistry. To investigate expression in the brain, for example, coronal and sagittal rat brain cryosections (20µm thick) are prepared using a Reichert-Jung cryostat. Individual sections are thaw-mounted onto silanized, nuclease-free slides (CEL Associates, Inc., Houston, TX), and stored at -80°C. Sections are processed starting with post-fixation in cold 4% paraformaldehyde, rinsed in cold phosphate-buffered saline (PBS), acetylated using acetic anhydride in triethanolamine buffer, and dehydrated through a series of alcohol washes in 70%, 95%, and 100% alcohol at room temperature. Subsequently, sections are delipidated in chloroform, followed by rehydration through successive exposure to 100% and 95% alcohol at room temperature. Microscope slides containing processed cryosections are allowed to air dry prior to hybridization. Other tissues may be assayed in a similar fashion.

[000249] A nGPCR-x-specific probe is generated using PCR. Following PCR amplification, the fragment is digested with restriction enzymes and cloned into pBluescript II cleaved with the same enzymes. For production of a probe specific for the sense strand of nGPCR-x, the nGPCR-x clone in pBluescript II is linearized with a suitable restriction enzyme, which provides a substrate for labeled run-off transcripts (*i.e.*, cRNA riboprobes) using the vector-borne T7 promoter and commercially available T7 RNA polymerase. A probe specific for the antisense strand of nGPCR-x is also readily prepared using the nGPCR-x clone in pBluescript II by cleaving the recombinant plasmid with a suitable restriction enzyme to generate a linearized substrate for the production of labeled run-off cRNA transcripts using the T3 promoter and cognate polymerase. The riboprobes are labeled with [³⁵S]-UTP to yield a specific activity of about 0.40 x 10⁶ cpm/pmol for antisense riboprobes and about 0.65 x 10⁶ cpm/pmol for sense-strand riboprobes. Each riboprobe is subsequently denatured and added (2 pmol/ml) to hybridization buffer which contained 50% formamide, 10% dextran, 0.3 M NaCl, 10 mM Tris

(pH 8.0), 1 mM EDTA, 1X Denhardt's Solution, and 10 mM dithiothreitol. Microscope slides containing sequential brain cryosections are independently exposed to 45 μ l of hybridization solution per slide and silanized cover slips are placed over the sections being exposed to hybridization solution. Sections are incubated overnight (15-18 hours) at 52°C to allow hybridization to occur. Equivalent series of cryosections are exposed to sense or antisense nGPCR-x-specific cRNA riboprobes.

[000250] Following the hybridization period, coverslips are washed off the slides in 1X SSC, followed by RNase A treatment involving the exposure of slides to 20 μ g/ml RNase A in a buffer containing 10mM Tris-HCl (pH 7.4), 0.5M EDTA, and 0.5M NaCl for 45 minutes at 37°C. The cryosections are then subjected to three high-stringency washes in 0.1 X SSC at 52°C for 20 minutes each. Following the series of washes, cryosections are dehydrated by consecutive exposure to 70%, 95%, and 100% ammonium acetate in alcohol, followed by air drying and exposure to Kodak BioMax™ MR-1 film. After 13 days of exposure, the film is developed. Based on these results, slides containing tissue that hybridized, as shown by film autoradiograms, are coated with Kodak NTB-2 nuclear track emulsion and the slides are stored in the dark for 32 days. The slides are then developed and counterstained with hematoxylin. Emulsion-coated sections are analyzed microscopically to determine the specificity of labeling. The signal is determined to be specific if autoradiographic grains (generated by antisense probe hybridization) are clearly associated with cresyl violet-stained cell bodies. Autoradiographic grains found between cell bodies indicates non-specific binding of the probe.

[000251] As discussed above, it is well known that GPCRs are expressed in many different tissues and regions, including in the brain. Expression of nGPCR-x in the brain provides an indication that modulators of nGPCR-x activity have utility for treating neurological disorders, including but not limited to, mental disorder, affective disorders, ADHD/ADD (*i.e.*, Attention Deficit Hyperactivity Disorder/Attention Deficit Disorder), and neural disorders such as Alzheimer's disease, Parkinson's disease, migraine, and senile dementia. Some other diseases for which modulators of nGPCR-x may have utility include depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like. Use of nGPCR-x modulators, including nGPCR-x ligands and anti-nGPCR-x antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

EXAMPLE 5: TISSUE EXPRESSION PROFILING

[000252] A PCR-based system (RapidScan™ Gene Expression Panel, OriGene Technologies, Rockville, MD) may be used to generate a comprehensive expression profile of the putative nGPCR-x in human tissue, and in human brain regions. The RapidScan Expression Panel is

comprised of first-strand cDNAs from various human tissues and brain regions that are serially diluted over a 4-log range and arrayed into a multi-well PCR plate. Human tissues in the array may include: brain, heart, kidney, spleen, liver, colon, lung, small intestine, muscle, stomach, testis, placenta, salivary gland, thyroid, adrenal gland, pancreas, ovary, uterus, prostate, skin, PBL, bone marrow, fetal brain, and fetal liver.

[000253] Expression of nGPCR-x in various tissues is detected using PCR primers designed based on the available sequence of the receptor that will prime the synthesis of a predetermined size fragment in the presence of the appropriate cDNA.

[000254] PCR is performed in a 50 μ l reaction containing 34 μ l H₂O, 5 μ l 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 μ l 15mM MgSO₄, 2 μ l dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10mM), 0.3 μ l forward primer (1 μ g/ μ l), 0.3 μ l reverse primer (1 μ g/ μ l), 0.4 μ l High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction mixture is added to each well of the PCR plate. The plate is placed in a MJ Research PTC100 thermocycler, and is then exposed to the following cycling parameters: Pre-soak 94°C for 3 min; denaturation at 94°C for 30 seconds; annealing at primer 57°C for 45 seconds; extension 72°C for 2 minutes; for 35 cycles. PCR productions are then separated and analyzed by electrophoresis on a 1.2% agarose gel stained with ethidium bromide.

[000255] The 4-log dilution range of cDNA deposited on the plate ensures that the amplification reaction is within the linear range and, hence, facilitates semi-quantitative determination of relative mRNA accumulation in the various tissues or brain regions examined.

EXAMPLE 6: NORTHERN BLOT ANALYSIS

[000256] Northern blots are performed to examine the expression of nGPCR-x mRNA. The sense orientation oligonucleotide and the antisense-orientation oligonucleotide, described above, are used as primers to amplify a portion of the GPCR-x cDNA sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58.

[000257] Multiple human tissue northern blots from Clontech (Human II # 7767-1) are hybridized with the probe. Pre-hybridization is carried out at 42 C for 4 hours in 5xSSC, 1X Denhardt's reagent, 0.1% SDS, 50% formamide, 250 mg/ml salmon sperm DNA. Hybridization is performed overnight at 42°C in the same mixture with the addition of about 1.5x10⁶ cpm/ml of labeled probe.

[000258] The probe is labeled with α -³²P-dCTP by Rediprime™ DNA labeling system (Amersham Pharmacia), purified on Nick Column™ (Amersham Pharmacia) and added to the hybridization solution. The filters are washed several times at 42°C in 0.2x SSC, 0.1% SDS.

Filters are exposed to Kodak XAR film (Eastman Kodak Company, Rochester, N.Y., USA) with intensifying screen at -80°C .

EXAMPLE 7: RECOMBINANT EXPRESSION OF nGPCR-X IN EUKARYOTIC HOST CELLS

A. Expression of nGPCR-x in Mammalian Cells

[000259] To produce nGPCR-x protein, a nGPCR-x-encoding polynucleotide is expressed in a suitable host cell using a suitable expression vector and standard genetic engineering techniques. For example, the nGPCR-x-encoding sequence described in Example 1 is subcloned into the commercial expression vector pzeoSV2 (Invitrogen, San Diego, CA) and transfected into Chinese Hamster Ovary (CHO) cells using the transfection reagent FuGENE6™ (Boehringer-Mannheim) and the transfection protocol provided in the product insert. Other eukaryotic cell lines, including human embryonic kidney (HEK-293) and COS cells, are suitable as well. Cells stably expressing nGPCR-x are selected by growth in the presence of $100\mu\text{g/ml}$ zeocin (Stratagene, LaJolla, CA). Optionally, nGPCR-x may be purified from the cells using standard chromatographic techniques. To facilitate purification, antisera is raised against one or more synthetic peptide sequences that correspond to portions of the nGPCR-x amino acid sequence, and the antisera is used to affinity purify nGPCR-x. The nGPCR-x also may be expressed in-frame with a tag sequence (e.g., polyhistidine, hemagglutinin, FLAG) to facilitate purification. Moreover, it will be appreciated that many of the uses for nGPCR-x polypeptides, such as assays described below, do not require purification of nGPCR-x from the host cell.

B. Expression of nGPCR-x in HEK-293 cells

[000260] For expression of nGPCR-x in mammalian cells HEK293 (transformed human, primary embryonic kidney cells), a plasmid bearing the relevant nGPCR-x coding sequence is prepared, using vector pSecTag2A (Invitrogen). Vector pSecTag2A contains the murine IgK chain leader sequence for secretion, the c-myc epitope for detection of the recombinant protein with the anti-myc antibody, a C-terminal polyhistidine for purification with nickel chelate chromatography, and a Zeocin resistant gene for selection of stable transfectants. The forward primer for amplification of this GPCR cDNA is determined by routine procedures and preferably contains a 5' extension of nucleotides to introduce the *HindIII* cloning site and nucleotides matching the GPCR sequence. The reverse primer is also determined by routine procedures and preferably contains a 5' extension of nucleotides to introduce an *XhoI* restriction site for cloning and nucleotides corresponding to the reverse complement of the nGPCR-x sequence. The PCR conditions are 55°C as the annealing temperature. The PCR product is gel purified and cloned into the *HindIII-XhoI* sites of the vector.

[000261] The DNA is purified using Qiagen chromatography columns and transfected into HEK-293 cells using DOTAP™ transfection media (Boehringer Mannheim, Indianapolis, IN). Transiently transfected cells are tested for expression after 24 hours of transfection, using western blots probed with anti-His and anti-nGPCR-x peptide antibodies. Permanently transfected cells are selected with Zeocin and propagated. Production of the recombinant protein is detected from both cells and media by western blots probed with anti-His, anti-Myc or anti-GPCR peptide antibodies.

C. Expression of nGPCR-x in COS cells

[000262] For expression of the nGPCR-x in COS7 cells, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 can be cloned into vector p3-CL. This vector is a pUC18-derived plasmid that contains the HCMV (human cytomegalovirus) promoter-intron located upstream from the bGH (bovine growth hormone) polyadenylation sequence and a multiple cloning site. In addition, the plasmid contains the dhfr (dihydrofolate reductase) gene which provides selection in the presence of the drug methotrexane (MTX) for selection of stable transformants.

[000263] The forward primer is determined by routine procedures and preferably contains a 5' extension which introduces an *XbaI* restriction site for cloning, followed by nucleotides which correspond to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58. The reverse primer is also determined by routine procedures and preferably contains 5'-extension of nucleotides which introduces a *Sall* cloning site followed by nucleotides which correspond to the reverse complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58. The PCR consists of an initial denaturation step of 5 min at 95°C, 30 cycles of 30 sec denaturation at 95°C, 30 sec annealing at 58°C and 30 sec extension at 72°C, followed by 5 min extension at 72°C. The PCR product is gel purified and ligated into the *XbaI* and *Sall* sites of vector p3-CL. This construct is transformed into *E. coli* cells for amplification and DNA purification. The DNA is purified with Qiagen chromatography columns and transfected into COS 7 cells using Lipofectamine™ reagent from BRL, following the manufacturer's protocols. Forty-eight and 72 hours after transfection, the media and the cells are tested for recombinant protein expression.

[000264] nGPCR-x expressed from a COS cell culture can be purified by concentrating the cell-growth media to about 10 mg of protein/ml, and purifying the protein by, for example, chromatography. Purified nGPCR-x is concentrated to 0.5 mg/ml in an Amicon concentrator fitted with a YM-10 membrane and stored at -80°C.

D. Expression of nGPCR-x in Insect Cells

[000265] For expression of nGPCR-x in a baculovirus system, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 can be amplified by PCR. The forward primer is determined by routine procedures and preferably contains a 5' extension which adds the *NdeI* cloning site, followed by nucleotides which correspond to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58. The reverse primer is also determined by routine procedures and preferably contains a 5' extension which introduces the *KpnI* cloning site, followed by nucleotides which correspond to the reverse complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58.

[000266] The PCR product is gel purified, digested with *NdeI* and *KpnI*, and cloned into the corresponding sites of vector pACHTL-A (Pharmingen, San Diego, CA). The pACHTL expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV), and a 6XHis tag upstream from the multiple cloning site. A protein kinase site for phosphorylation and a thrombin site for excision of the recombinant protein precede the multiple cloning site is also present. Of course, many other baculovirus vectors could be used in place of pACHTL-A, such as pAc373, pVL941 and pAcIM1. Other suitable vectors for the expression of GPCR polypeptides can be used, provided that the vector construct includes appropriately located signals for transcription, translation, and trafficking, such as an in-frame AUG and a signal peptide, as required. Such vectors are described in Luckow *et al.*, Virology 170:31-39, among others.

[000267] The virus is grown and isolated using standard baculovirus expression methods, such as those described in Summers *et al.* (A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Texas Agricultural Experimental Station Bulletin No. 1555 (1987)).

[000268] In a preferred embodiment, pACHTL-A containing nGPCR-x gene is introduced into baculovirus using the "BaculoGold™" transfection kit (Pharmingen, San Diego, CA) using methods established by the manufacturer. Individual virus isolates are analyzed for protein production by radiolabeling infected cells with ³⁵S-methionine at 24 hours post infection. Infected cells are harvested at 48 hours post infection, and the labeled proteins are visualized by SDS-PAGE. Viruses exhibiting high expression levels can be isolated and used for scaled up expression.

[000269] For expression of a nGPCR-x polypeptide in a Sf9 cells, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 can be amplified by PCR using the primers and methods described above for baculovirus expression. The nGPCR-x cDNA is cloned into vector pACHTL-A (Pharmingen) for expression in Sf9 insect. The insert is cloned into the *NdeI* and *KpnI* sites, after elimination of an internal *NdeI*

site (using the same primers described above for expression in baculovirus). DNA is purified with Qiagen chromatography columns and expressed in Sf9 cells. Preliminary Western blot experiments from non-purified plaques are tested for the presence of the recombinant protein of the expected size which reacted with the GPCR-specific antibody. These results are confirmed after further purification and expression optimization in HiG5 cells.

EXAMPLE 8: INTERACTION TRAP/TWO-HYBRID SYSTEM

- [000270] In order to assay for nGPCR-x-interacting proteins, the interaction trap/two-hybrid library screening method can be used. This assay was first described in Fields *et al*, *Nature*, 1989, 340, 245, which is incorporated herein by reference in its entirety. A protocol is published in Current Protocols in Molecular Biology 1999, John Wiley & Sons, NY, and Ausubel, F. M. *et al*. 1992, Short protocols in molecular biology, Fourth edition, Greene and Wiley-interscience, NY, each of which is incorporated herein by reference in its entirety. Kits are available from Clontech, Palo Alto, CA (Matchmaker Two-Hybrid System 3).
- [000271] A fusion of the nucleotide sequences encoding all or partial nGPCR-x and the yeast transcription factor GAL4 DNA-binding domain (DNA-BD) is constructed in an appropriate plasmid (*i.e.*, pGBKT7) using standard subcloning techniques. Similarly, a GAL4 active domain (AD) fusion library is constructed in a second plasmid (*i.e.*, pGADT7) from cDNA of potential GPCR-binding proteins (for protocols on forming cDNA libraries, see Sambrook *et al*. 1989, Molecular cloning: a laboratory manual, second edition, Cold Spring Harbor Press, Cold Spring Harbor, NY), which is incorporated herein by reference in its entirety. The DNA-BD/nGPCR-x fusion construct is verified by sequencing, and tested for autonomous reporter gene activation and cell toxicity, both of which would prevent a successful two-hybrid analysis. Similar controls are performed with the AD/library fusion construct to ensure expression in host cells and lack of transcriptional activity. Yeast cells are transformed (*ca* 10⁵ transformants/mg DNA) with both the nGPCR-x and library fusion plasmids according to standard procedures (Ausubel *et al.*, 1992, Short protocols in molecular biology, fourth edition, Greene and Wiley-interscience, NY, which is incorporated herein by reference in its entirety). *In vivo* binding of DNA-BD/nGPCR-x with AD/library proteins results in transcription of specific yeast plasmid reporter genes (*i.e.*, lacZ, HIS3, ADE2, LEU2). Yeast cells are plated on nutrient-deficient media to screen for expression of reporter genes. Colonies are dually assayed for β -galactosidase activity upon growth in Xgal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) supplemented media (filter assay for β -galactosidase activity is described in Breeden *et al.*, Cold Spring Harb. Symp. Quant. Biol., 1985, 50, 643, which is incorporated herein by reference in its entirety). Positive AD-library plasmids are rescued from transformants and reintroduced into

the original yeast strain as well as other strains containing unrelated DNA-BD fusion proteins to confirm specific nGPCR-x/library protein interactions. Insert DNA is sequenced to verify the presence of an open reading frame fused to GAL4 AD and to determine the identity of the nGPCR-x-binding protein.

EXAMPLE 9: MOBILITY SHIFT DNA-BINDING ASSAY USING GEL ELECTROPHORESIS

[000272] A gel electrophoresis mobility shift assay can rapidly detect specific protein-DNA interactions. Protocols are widely available in such manuals as Sambrook *et al.* 1989, *Molecular cloning: a laboratory manual*, second edition, Cold Spring Harbor Press, Cold Spring Harbor, NY and Ausubel, F. M. *et al.*, 1992, *Short Protocols in Molecular Biology*, fourth edition, Greene and Wiley-interscience, NY, each of which is incorporated herein by reference in its entirety.

[000273] Probe DNA (<300 bp) is obtained from synthetic oligonucleotides, restriction endonuclease fragments, or PCR fragments and end-labeled with ³²P. An aliquot of purified nGPCR-x (*ca.* 15 µg) or crude nGPCR-x extract (*ca.* 15 ng) is incubated at constant temperature (in the range 22-37 C) for at least 30 minutes in 10-15 µl of buffer (*i.e.* TAE or TBE, pH 8.0-8.5) containing radiolabeled probe DNA, nonspecific carrier DNA (*ca.* 1 µg), BSA (300 µg/ml), and 10% (v/v) glycerol. The reaction mixture is then loaded onto a polyacrylamide gel and run at 30-35 mA until good separation of free probe DNA from protein-DNA complexes occurs. The gel is then dried and bands corresponding to free DNA and protein-DNA complexes are detected by autoradiography.

EXAMPLE 10: ANTIBODIES TO nGPCR-X

[000274] Standard techniques are employed to generate polyclonal or monoclonal antibodies to the nGPCR-x receptor, and to generate useful antigen-binding fragments thereof or variants thereof, including "humanized" variants. Such protocols can be found, for example, in Sambrook *et al.* (1989) and Harlow *et al.* (Eds.), Antibodies A Laboratory Manual; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988). In one embodiment, recombinant nGPCR-x polypeptides (or cells or cell membranes containing such polypeptides) are used as antigen to generate the antibodies. In another embodiment, one or more peptides having amino acid sequences corresponding to an immunogenic portion of nGPCR-x (*e.g.* 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more amino acids) are used as antigen. Peptides corresponding to extracellular portions of nGPCR-x, especially hydrophilic extracellular portions, are preferred.

The antigen may be mixed with an adjuvant or linked to a hapten to increase antibody production.

A. Polyclonal or Monoclonal antibodies

- [000275] As one exemplary protocol, recombinant nGPCR-x or a synthetic fragment thereof is used to immunize a mouse for generation of monoclonal antibodies (or larger mammal, such as a rabbit, for polyclonal antibodies). To increase antigenicity, peptides are conjugated to Keyhole Limpet Hemocyanin (Pierce), according to the manufacturer's recommendations. For an initial injection, the antigen is emulsified with Freund's Complete Adjuvant and injected subcutaneously. At intervals of two to three weeks, additional aliquots of nGPCR-x antigen are emulsified with Freund's Incomplete Adjuvant and injected subcutaneously. Prior to the final booster injection, a serum sample is taken from the immunized mice and assayed by western blot to confirm the presence of antibodies that immunoreact with nGPCR-x. Serum from the immunized animals may be used as polyclonal antisera or used to isolate polyclonal antibodies that recognize nGPCR-x. Alternatively, the mice are sacrificed and their spleen removed for generation of monoclonal antibodies.
- [000276] To generate monoclonal antibodies, the spleens are placed in 10 ml serum-free RPMI 1640, and single cell suspensions are formed by grinding the spleens in serum-free RPMI 1640, supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, 100 units/ml penicillin, and 100 µg/ml streptomycin (RPMI) (Gibco, Canada). The cell suspensions are filtered and washed by centrifugation and resuspended in serum-free RPMI. Thymocytes taken from three naive Balb/c mice are prepared in a similar manner and used as a Feeder Layer. NS-1 myeloma cells, kept in log phase in RPMI with 10% fetal bovine serum (FBS) (Hyclone Laboratories, Inc., Logan, Utah) for three days prior to fusion, are centrifuged and washed as well.
- [000277] To produce hybridoma fusions, spleen cells from the immunized mice are combined with NS-1 cells and centrifuged, and the supernatant is aspirated. The cell pellet is dislodged by tapping the tube, and 2 ml of 37°C PEG 1500 (50% in 75 mM HEPES, pH 8.0) (Boehringer-Mannheim) is stirred into the pellet, followed by the addition of serum-free RPMI. Thereafter, the cells are centrifuged, resuspended in RPMI containing 15% FBS, 100 µM sodium hypoxanthine, 0.4 µM aminopterin, 16 µM thymidine (HAT) (Gibco), 25 units/ml IL-6 (Boehringer-Mannheim) and 1.5×10^6 thymocytes/ml, and plated into 10 Corning flat-bottom 96-well tissue culture plates (Corning, Corning New York).
- [000278] On days 2, 4, and 6 after the fusion, 100µl of medium is removed from the wells of the fusion plates and replaced with fresh medium. On day 8, the fusions are screened by ELISA, testing for the presence of mouse IgG that binds to nGPCR-x. Selected fusion wells are further cloned by dilution until monoclonal cultures producing anti-nGPCR-x antibodies are obtained.

B. Humanization of anti-nGPCR-x monoclonal antibodies

- [000279] The expression pattern of nGPCR-x as reported herein and the proven track record of GPCRs as targets for therapeutic intervention suggest therapeutic indications for nGPCR-x inhibitors (antagonists). nGPCR-x-neutralizing antibodies comprise one class of therapeutics useful as nGPCR-x antagonists. Following are protocols to improve the utility of anti-nGPCR-x monoclonal antibodies as therapeutics in humans by "humanizing" the monoclonal antibodies to improve their serum half-life and render them less immunogenic in human hosts (*i.e.*, to prevent human antibody response to non-human anti-nGPCR-x antibodies).
- [000280] The principles of humanization have been described in the literature and are facilitated by the modular arrangement of antibody proteins. To minimize the possibility of binding complement, a humanized antibody of the IgG4 isotype is preferred.
- [000281] For example, a level of humanization is achieved by generating chimeric antibodies comprising the variable domains of non-human antibody proteins of interest with the constant domains of human antibody molecules. (See, *e.g.*, Morrison *et al.*, Adv. Immunol., 44:65-92 (1989)). The variable domains of nGPCR-x-neutralizing anti-nGPCR-x antibodies are cloned from the genomic DNA of a B-cell hybridoma or from cDNA generated from mRNA isolated from the hybridoma of interest. The V region gene fragments are linked to exons encoding human antibody constant domains, and the resultant construct is expressed in suitable mammalian host cells (*e.g.*, myeloma or CHO cells).
- [000282] To achieve an even greater level of humanization, only those portions of the variable region gene fragments that encode antigen-binding complementarity determining regions ("CDR") of the non-human monoclonal antibody genes are cloned into human antibody sequences. (See, *e.g.*, Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-327 (1988); Verhoeyen *et al.*, Science 239:1534-36 (1988); and Tempest *et al.*, Bio/Technology 9: 266-71 (1991)). If necessary, the β -sheet framework of the human antibody surrounding the CDR3 regions also is modified to more closely mirror the three dimensional structure of the antigen-binding domain of the original monoclonal antibody. (See Kettleborough *et al.*, Protein Engin., 4:773-783 (1991); and Foote *et al.*, J. Mol. Biol., 224:487-499 (1992)).
- [000283] In an alternative approach, the surface of a non-human monoclonal antibody of interest is humanized by altering selected surface residues of the non-human antibody, *e.g.*, by site-directed mutagenesis, while retaining all of the interior and contacting residues of the non-human antibody. See Padlan, Molecular Immunol., 28(4/5):489-98 (1991).
- [000284] The foregoing approaches are employed using nGPCR-x-neutralizing anti-nGPCR-x monoclonal antibodies and the hybridomas that produce them to generate humanized nGPCR-

x-neutralizing antibodies useful as therapeutics to treat or palliate conditions wherein nGPCR-x expression or ligand-mediated nGPCR-x signaling is detrimental.

C. Human nGPCR-x-Neutralizing Antibodies from Phage Display

[000285] Human nGPCR-x-neutralizing antibodies are generated by phage display techniques such as those described in Aujame *et al.*, Human Antibodies 8(4):155-168 (1997); Hoogenboom, TIBTECH 15:62-70 (1997); and Rader *et al.*, Curr. Opin. Biotechnol. 8:503-508 (1997), all of which are incorporated by reference. For example, antibody variable regions in the form of Fab fragments or linked single chain Fv fragments are fused to the amino terminus of filamentous phage minor coat protein pIII. Expression of the fusion protein and incorporation thereof into the mature phage coat results in phage particles that present an antibody on their surface and contain the genetic material encoding the antibody. A phage library comprising such constructs is expressed in bacteria, and the library is screened for nGPCR-x-specific phage-antibodies using labeled or immobilized nGPCR-x as antigen-probe.

D. Human nGPCR-x-neutralizing antibodies from transgenic mice

[000286] Human nGPCR-x-neutralizing antibodies are generated in transgenic mice essentially as described in Bruggemann *et al.*, Immunol. Today 17(8):391-97 (1996) and Bruggemann *et al.*, Curr. Opin. Biotechnol. 8:455-58 (1997). Transgenic mice carrying human V-gene segments in germline configuration and that express these transgenes in their lymphoid tissue are immunized with a nGPCR-x composition using conventional immunization protocols. Hybridomas are generated using B cells from the immunized mice using conventional protocols and screened to identify hybridomas secreting anti-nGPCR-x human antibodies (*e.g.*, as described above).

EXAMPLE 11: ASSAYS TO IDENTIFY MODULATORS OF nGPCR-X ACTIVITY

[000287] Set forth below are several nonlimiting assays for identifying modulators (agonists and antagonists) of nGPCR-x activity. Among the modulators that can be identified by these assays are natural ligand compounds of the receptor; synthetic analogs and derivatives of natural ligands; antibodies, antibody fragments, and/or antibody-like compounds derived from natural antibodies or from antibody-like combinatorial libraries; and/or synthetic compounds identified by high-throughput screening of libraries; and the like. All modulators that bind nGPCR-x are useful for identifying nGPCR-x in tissue samples (*e.g.*, for diagnostic purposes, pathological purposes, and the like). Agonist and antagonist modulators are useful for up-regulating and down-regulating nGPCR-x activity, respectively, to treat disease states characterized by abnormal levels of nGPCR-x activity. The assays may be performed using single putative modulators, and/or may be performed using a known agonist in combination with candidate antagonists (or *visa versa*).

A. cAMP Assays

[000288] In one type of assay, levels of cyclic adenosine monophosphate (cAMP) are measured in nGPCR-x-transfected cells that have been exposed to candidate modulator compounds.

Protocols for cAMP assays have been described in the literature. (See, *e.g.*, Sutherland *et al.*, *Circulation* 37: 279 (1968); Frandsen *et al.*, *Life Sciences* 18: 529-541 (1976); Dooley *et al.*, *Journal of Pharmacology and Experimental Therapeutics* 283 (2): 735-41 (1997); and George *et al.*, *Journal of Biomolecular Screening* 2 (4): 235-40 (1997)). An exemplary protocol for such an assay, using an Adenylyl Cyclase Activation FlashPlate® Assay from NEN™ Life Science Products, is set forth below.

[000289] Briefly, the nGPCR-x coding sequence (*e.g.*, a cDNA or intronless genomic DNA) is subcloned into a commercial expression vector, such as pzeoSV2 (Invitrogen), and transiently transfected into Chinese Hamster Ovary (CHO) cells using known methods, such as the transfection protocol provided by Boehringer-Mannheim when supplying the FuGENE 6 transfection reagent. Transfected CHO cells are seeded into 96-well microplates from the FlashPlate® assay kit, which are coated with solid scintillant to which antisera to cAMP has been bound. For a control, some wells are seeded with wild type (untransfected) CHO cells. Other wells in the plate receive various amounts of a cAMP standard solution for use in creating a standard curve.

[000290] One or more test compounds (*i.e.*, candidate modulators) are added to the cells in each well, with water and/or compound-free medium/diluent serving as a control or controls. After treatment, cAMP is allowed to accumulate in the cells for exactly 15 minutes at room temperature. The assay is terminated by the addition of lysis buffer containing [¹²⁵I]-labeled cAMP, and the plate is counted using a Packard Topcount™ 96-well microplate scintillation counter. Unlabeled cAMP from the lysed cells (or from standards) and fixed amounts of [¹²⁵I]-cAMP compete for antibody bound to the plate. A standard curve is constructed, and cAMP values for the unknowns are obtained by interpolation. Changes in intracellular cAMP levels of cells in response to exposure to a test compound are indicative of nGPCR-x modulating activity. Modulators that act as agonists of receptors which couple to the G_s subtype of G proteins will stimulate production of cAMP, leading to a measurable 3-10 fold increase in cAMP levels. Agonists of receptors which couple to the G_{i/o} subtype of G proteins will inhibit forskolin-stimulated cAMP production, leading to a measurable decrease in cAMP levels of 50-100%. Modulators that act as inverse agonists will reverse these effects at receptors that are either constitutively active or activated by known agonists.

B. Aequorin Assays

- [000291] In another assay, cells (*e.g.*, CHO cells) are transiently co-transfected with both a nGPCR-x expression construct and a construct that encodes the photoprotein apoaequorin. In the presence of the cofactor coelenterazine, apoaequorin will emit a measurable luminescence that is proportional to the amount of intracellular (cytoplasmic) free calcium. (See generally, Cobbold, *et al.* "Aequorin measurements of cytoplasmic free calcium," *In*: McCormack J.G. and Cobbold P.H., eds., *Cellular Calcium: A Practical Approach*. Oxford:IRL Press (1991); Stables *et al.*, *Analytical Biochemistry* 252: 115-26 (1997); and Haugland, *Handbook of Fluorescent Probes and Research Chemicals*. Sixth edition. Eugene OR: Molecular Probes (1996).)
- [000292] In one exemplary assay, nGPCR-x is subcloned into the commercial expression vector pzeoSV2 (Invitrogen) and transiently co-transfected along with a construct that encodes the photoprotein apoaequorin (Molecular Probes, Eugene, OR) into CHO cells using the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert.
- [000293] The cells are cultured for 24 hours at 37°C in MEM (Gibco/BRL, Gaithersburg, MD) supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10 µg/ml streptomycin, at which time the medium is changed to serum-free MEM containing 5 µM coelenterazine (Molecular Probes, Eugene, OR). Culturing is then continued for two additional hours at 37°C. Subsequently, cells are detached from the plate using VERSEN (Gibco/BRL), washed, and resuspended at 200,000 cells/ml in serum-free MEM.
- [000294] Dilutions of candidate nGPCR-x modulator compounds are prepared in serum-free MEM and dispensed into wells of an opaque 96-well assay plate at 50 µl/well. Plates are then loaded onto an MLX microtiter plate luminometer (Dynex Technologies, Inc., Chantilly, VA). The instrument is programmed to dispense 50µl cell suspensions into each well, one well at a time, and immediately read luminescence for 15 seconds. Dose-response curves for the candidate modulators are constructed using the area under the curve for each light signal peak. Data are analyzed with SlideWrite, using the equation for a one-site ligand, and EC₅₀ values are obtained. Changes in luminescence caused by the compounds are considered indicative of modulatory activity. Modulators that act as agonists at receptors which couple to the G_q subtype of G proteins give an increase in luminescence of up to 100 fold. Modulators that act as inverse agonists will reverse this effect at receptors that are either constitutively active or activated by known agonists.

C. Luciferase Reporter Gene Assay

- [000295] The photoprotein luciferase provides another useful tool for assaying for modulators of nGPCR-x activity. Cells (*e.g.*, CHO cells or COS 7 cells) are transiently co-transfected with both a nGPCR-x expression construct (*e.g.*, nGPCR-x in pzeoSV2) and a reporter construct

which includes a gene for the luciferase protein downstream from a transcription factor binding site, such as the cAMP-response element (CRE), AP-1, or NF-kappa B. Agonist binding to receptors coupled to the G_s subtype of G proteins leads to increases in cAMP, thereby activating the CRE transcription factor and resulting in expression of the luciferase gene. Agonist binding to receptors coupled to the G_q subtype of G protein leads to production of diacylglycerol that activates protein kinase C, which activates the AP-1 or NF-kappa B transcription factors, in turn resulting in expression of the luciferase gene. Expression levels of luciferase reflect the activation status of the signaling events. (See generally, George *et al*, Journal of Biomolecular Screening 2(4): 235-240 (1997); and Stratowa *et al*, Current Opinion in Biotechnology 6: 574-581 (1995)). Luciferase activity may be quantitatively measured using, *e.g.*, luciferase assay reagents that are commercially available from Promega (Madison, WI).

[000296] In one exemplary assay, CHO cells are plated in 24-well culture dishes at a density of 100,000 cells/well one day prior to transfection and cultured at 37°C in MEM (Gibco/BRL) supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10 µg/ml streptomycin. Cells are transiently co-transfected with both a nGPCR-x expression construct and a reporter construct containing the luciferase gene. The reporter plasmids CRE-luciferase, AP-1-luciferase and NF-kappaB-luciferase may be purchased from Stratagene (LaJolla, CA). Transfections are performed using the FuGENE 6 transfection reagent (Boehringer-Mannheim) according to the supplier's instructions. Cells transfected with the reporter construct alone are used as a control. Twenty-four hours after transfection, cells are washed once with PBS pre-warmed to 37°C. Serum-free MEM is then added to the cells either alone (control) or with one or more candidate modulators and the cells are incubated at 37°C for five hours. Thereafter, cells are washed once with ice-cold PBS and lysed by the addition of 100 µl of lysis buffer per well from the luciferase assay kit supplied by Promega. After incubation for 15 minutes at room temperature, 15 µl of the lysate is mixed with 50 µl of substrate solution (Promega) in an opaque-white, 96-well plate, and the luminescence is read immediately on a Wallace model 1450 MicroBeta scintillation and luminescence counter (Wallace Instruments, Gaithersburg, MD).

[000297] Differences in luminescence in the presence versus the absence of a candidate modulator compound are indicative of modulatory activity. Receptors that are either constitutively active or activated by agonists typically give a 3 to 20-fold stimulation of luminescence compared to cells transfected with the reporter gene alone. Modulators that act as inverse agonists will reverse this effect.

D. Intracellular calcium measurement using FLIPR

[000298] Changes in intracellular calcium levels are another recognized indicator of G protein-coupled receptor activity, and such assays can be employed to screen for modulators of nGPCR-x activity. For example, CHO cells stably transfected with a nGPCR-x expression vector are plated at a density of 4×10^4 cells/well in Packard black-walled, 96-well plates specially designed to discriminate fluorescence signals emanating from the various wells on the plate. The cells are incubated for 60 minutes at 37°C in modified Dulbecco's PBS (D-PBS) containing 36 mg/L pyruvate and 1 g/L glucose with the addition of 1% fetal bovine serum and one of four calcium indicator dyes (Fluo-3™ AM, Fluo-4™ AM, Calcium Green™-1 AM, or Oregon Green™ 488 BAPTA-1 AM), each at a concentration of 4 μM. Plates are washed once with modified D-PBS without 1% fetal bovine serum and incubated for 10 minutes at 37°C to remove residual dye from the cellular membrane. In addition, a series of washes with modified D-PBS without 1% fetal bovine serum is performed immediately prior to activation of the calcium response.

[000299] A calcium response is initiated by the addition of one or more candidate receptor agonist compounds, calcium ionophore A23187 (10 μM; positive control), or ATP (4 μM; positive control). Fluorescence is measured by Molecular Device's FLIPR with an argon laser (excitation at 488 nm). (See, e.g., Kuntzweiler *et al.*, Drug Development Research, 44(1):14-20 (1998)). The F-stop for the detector camera was set at 2.5 and the length of exposure was 0.4 milliseconds. Basal fluorescence of cells was measured for 20 seconds prior to addition of candidate agonist, ATP, or A23187, and the basal fluorescence level was subtracted from the response signal. The calcium signal is measured for approximately 200 seconds, taking readings every two seconds. Calcium ionophore A23187 and ATP increase the calcium signal 200% above baseline levels. In general, activated GPCRs increase the calcium signal approximately 10-15% above baseline signal.

E. Mitogenesis Assay

[000300] In a mitogenesis assay, the ability of candidate modulators to induce or inhibit nGPCR-x-mediated cell division is determined. (See, e.g., Lajiness *et al.*, Journal of Pharmacology and Experimental Therapeutics 267(3): 1573-1581 (1993)). For example, CHO cells stably expressing nGPCR-x are seeded into 96-well plates at a density of 5000 cells/well and grown at 37°C in MEM with 10% fetal calf serum for 48 hours, at which time the cells are rinsed twice with serum-free MEM. After rinsing, 80 μl of fresh MEM, or MEM containing a known mitogen, is added along with 20 μl MEM containing varying concentrations of one or more candidate modulators or test compounds diluted in serum-free medium. As controls, some wells on each plate receive serum-free medium alone, and some receive medium containing 10% fetal

bovine serum. Untransfected cells or cells transfected with vector alone also may serve as controls.

[000301] After culture for 16-18 hours, 1 μ Ci of [3 H]-thymidine (2 Ci/mmol) is added to the wells and cells are incubated for an additional 2 hours at 37°C. The cells are trypsinized and collected on filter mats with a cell harvester (Tomtec); the filters are then counted in a Betaplate counter. The incorporation of [3 H]-thymidine in serum-free test wells is compared to the results achieved in cells stimulated with serum (positive control). Use of multiple concentrations of test compounds permits creation and analysis of dose-response curves using the non-linear, least squares fit equation: $A = B \times [C / (D + C)] + G$ where A is the percent of serum stimulation; B is the maximal effect minus baseline; C is the EC₅₀; D is the concentration of the compound; and G is the maximal effect. Parameters B, C and G are determined by Simplex optimization.

[000302] Agonists that bind to the receptor are expected to increase [3 H]-thymidine incorporation into cells, showing up to 80% of the response to serum. Antagonists that bind to the receptor will inhibit the stimulation seen with a known agonist by up to 100%.

F. [35 S]GTP γ S Binding Assay

[000303] Because G protein-coupled receptors signal through intracellular G proteins whose activity involves GTP binding and hydrolysis to yield bound GDP, measurement of binding of the non-hydrolyzable GTP analog [35 S]GTP γ S in the presence and absence of candidate modulators provides another assay for modulator activity. (See, e.g., Kowal *et al*, Neuropharmacology 37:179-187 (1998).)

[000304] In one exemplary assay, cells stably transfected with a nGPCR-x expression vector are grown in 10 cm tissue culture dishes to subconfluence, rinsed once with 5 ml of ice-cold Ca²⁺/Mg²⁺-free phosphate-buffered saline, and scraped into 5 ml of the same buffer. Cells are pelleted by centrifugation (500 x g, 5 minutes), resuspended in TEE buffer (25 mM Tris, pH 7.5, 5 mM EDTA, 5 mM EGTA), and frozen in liquid nitrogen. After thawing, the cells are homogenized using a Dounce homogenizer (one ml TEE per plate of cells), and centrifuged at 1,000 x g for 5 minutes to remove nuclei and unbroken cells.

[000305] The homogenate supernatant is centrifuged at 20,000 x g for 20 minutes to isolate the membrane fraction, and the membrane pellet is washed once with TEE and resuspended in binding buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 10 mM MgCl₂, 1 mM EDTA). The resuspended membranes can be frozen in liquid nitrogen and stored at -70°C until use.

[000306] Aliquots of cell membranes prepared as described above and stored at -70°C are thawed, homogenized, and diluted into buffer containing 20 mM HEPES, 10 mM MgCl₂, 1 mM EDTA, 120 mM NaCl, 10 μ M GDP, and 0.2 mM ascorbate, at a concentration of 10-50 μ g/ml. In a final volume of 90 μ l, homogenates are incubated with varying concentrations of candidate modulator

compounds or 100 μ M GTP for 30 minutes at 30°C and then placed on ice. To each sample, 10 μ l guanosine 5'-O-(3[³⁵S]thio) triphosphate (NEN, 1200 Ci/mmol; [³⁵S]-GTP γ S), was added to a final concentration of 100-200 pM. Samples are incubated at 30°C for an additional 30 minutes, 1 ml of 10mM HEPES, pH 7.4, 10 mM MgCl₂, at 4°C is added and the reaction is stopped by filtration.

[000307] Samples are filtered over Whatman GF/B filters and the filters are washed with 20 ml ice-cold 10 mM HEPES, pH 7.4, 10 mM MgCl₂. Filters are counted by liquid scintillation spectroscopy. Nonspecific binding of [³⁵S]-GTP γ S is measured in the presence of 100 μ M GTP and subtracted from the total. Compounds are selected that modulate the amount of [³⁵S]-GTP γ S binding in the cells, compared to untransfected control cells. Activation of receptors by agonists gives up to a five-fold increase in [³⁵S]GTP γ S binding. This response is blocked by antagonists.

G. MAP Kinase Activity Assay

[000308] Evaluation of MAP kinase activity in cells expressing a GPCR provides another assay to identify modulators of GPCR activity. (See, *e.g.*, Lajiness *et al.*, Journal of Pharmacology and Experimental Therapeutics 267(3):1573-1581 (1993) and Boulton *et al.*, Cell 65:663-675 (1991).)

[000309] In one embodiment, CHO cells stably transfected with nGPCR-x are seeded into 6-well plates at a density of 70,000 cells/well 48 hours prior to the assay. During this 48-hour period, the cells are cultured at 37°C in MEM medium supplemented with 10% fetal bovine serum, 2mM glutamine, 10 U/ml penicillin and 10 μ g/ml streptomycin. The cells are serum-starved for 1-2 hours prior to the addition of stimulants.

[000310] For the assay, the cells are treated with medium alone or medium containing either a candidate agonist or 200 nM Phorbol ester- myristoyl acetate (*i.e.*, PMA, a positive control), and the cells are incubated at 37°C for varying times. To stop the reaction, the plates are placed on ice, the medium is aspirated, and the cells are rinsed with 1 ml of ice-cold PBS containing 1mM EDTA. Thereafter, 200 μ l of cell lysis buffer (12.5 mM MOPS, pH 7.3, 12.5 mM glycerophosphate, 7.5mM MgCl₂, 0.5mM EGTA, 0.5 mM sodium vanadate, 1mM benzamidine, 1mM dithiothreitol, 10 μ g/ml leupeptin, 10 μ g/ml aprotinin, 2 μ g/ml pepstatin A, and 1 μ M okadaic acid) is added to the cells. The cells are scraped from the plates and homogenized by 10 passages through a 23 3/4 G needle, and the cytosol fraction is prepared by centrifugation at 20,000 x g for 15 minutes.

[000311] Aliquots (5-10 μ l containing 1-5 μ g protein) of cytosol are mixed with 1 mM MAPK Substrate Peptide (APRTPGGRR (SEQ ID NO: 117), Upstate Biotechnology, Inc., N.Y.) and 50 μ M [γ -³²P]ATP (NEN, 3000 Ci/mmol), diluted to a final specific activity of ~2000 cpm/pmol,

in a total volume of 25 μ l. The samples are incubated for 5 minutes at 30°C, and reactions are stopped by spotting 20 μ l on 2 cm² squares of Whatman P81 phosphocellulose paper. The filter squares are washed in 4 changes of 1% H₃PO₄, and the squares are subjected to liquid scintillation spectroscopy to quantitate bound label. Equivalent cytosolic extracts are incubated without MAPK substrate peptide, and the bound label from these samples are subtracted from the matched samples with the substrate peptide. The cytosolic extract from each well is used as a separate point. Protein concentrations are determined by a dye binding protein assay (Bio-Rad Laboratories). Agonist activation of the receptor is expected to result in up to a five-fold increase in MAPK enzyme activity. This increase is blocked by antagonists.

H. [³H]Arachidonic Acid Release

[000312] The activation of GPCRs also has been observed to potentiate arachidonic acid release in cells, providing yet another useful assay for modulators of GPCR activity. (See, e.g., Kanterman *et al.*, Molecular Pharmacology 39:364-369 (1991).) For example, CHO cells that are stably transfected with a nGPCR-x expression vector are plated in 24-well plates at a density of 15,000 cells/well and grown in MEM medium supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10 μ g/ml streptomycin for 48 hours at 37°C before use. Cells of each well are labeled by incubation with [³H]-arachidonic acid (Amersham Corp., 210 Ci/mmol) at 0.5 μ Ci/ml in 1 ml MEM supplemented with 10mM HEPES, pH 7.5, and 0.5% fatty-acid-free bovine serum albumin for 2 hours at 37°C. The cells are then washed twice with 1 ml of the same buffer.

[000313] Candidate modulator compounds are added in 1 ml of the same buffer, either alone or with 10 μ M ATP and the cells are incubated at 37°C for 30 minutes. Buffer alone and mock-transfected cells are used as controls. Samples (0.5 ml) from each well are counted by liquid scintillation spectroscopy. Agonists which activate the receptor will lead to potentiation of the ATP-stimulated release of [³H]-arachidonic acid. This potentiation is blocked by antagonists.

I. Extracellular Acidification Rate

[000314] In yet another assay, the effects of candidate modulators of nGPCR-x activity are assayed by monitoring extracellular changes in pH induced by the test compounds. (See, e.g., Dunlop *et al.*, Journal of Pharmacological and Toxicological Methods 40(1):47-55 (1998).) In one embodiment, CHO cells transfected with a nGPCR-x expression vector are seeded into 12 mm capsule cups (Molecular Devices Corp.) at 4 x 10⁵ cells/cup in MEM supplemented with 10% fetal bovine serum, 2 mM L-glutamine, 10 U/ml penicillin, and 10 μ g/ml streptomycin. The cells are incubated in this medium at 37°C in 5% CO₂ for 24 hours.

[000315] Extracellular acidification rates are measured using a Cytosensor microphysiometer (Molecular Devices Corp.). The capsule cups are loaded into the sensor chambers of the

microphysiometer and the chambers are perfused with running buffer (bicarbonate-free MEM supplemented with 4 mM L-glutamine, 10 units/ml penicillin, 10 µg/ml streptomycin, 26 mM NaCl) at a flow rate of 100 µl/minute. Candidate agonists or other agents are diluted into the running buffer and perfused through a second fluid path. During each 60-second pump cycle, the pump is run for 38 seconds and is off for the remaining 22 seconds. The pH of the running buffer in the sensor chamber is recorded during the cycle from 43-58 seconds, and the pump is re-started at 60 seconds to start the next cycle. The rate of acidification of the running buffer during the recording time is calculated by the Cytosoft program. Changes in the rate of acidification are calculated by subtracting the baseline value (the average of 4 rate measurements immediately before addition of a modulator candidate) from the highest rate measurement obtained after addition of a modulator candidate. The selected instrument detects 61mV/pH unit. Modulators that act as agonists of the receptor result in an increase in the rate of extracellular acidification compared to the rate in the absence of agonist. This response is blocked by modulators which act as antagonists of the receptor.

Example 12 - Using nGPCR-x proteins to isolate neurotransmitters

[000316] Isolated nGPCR-x proteins of the present invention can be used to isolate novel or known neurotransmitters (Saito *et al.*, Nature 400: 265-269, 1999). The cDNAs that encode the isolated nGPCR-x can be cloned into mammalian expression vectors and used to stably or transiently transfect mammalian cells including CHO, Cos or HEK293 cells. Receptor expression can be determined by Northern blot analysis of transfected cells and identification of an appropriately sized mRNA band (predicted size from the cDNA). Brain regions shown by mRNA analysis to express each of the nGPCR-x proteins could be processed for peptide extraction using any of several protocols ((Reinscheid R.K. *et al.*, Science 270: 243-247, 1996; Sakurai, T., *et al.*, Cell 92: 573-585, 1998; Hinuma, S., *et al.*, Nature 393: 272-276, 1998). Chromatographic fractions of brain extracts could be tested for ability to activate nGPCR-x proteins by measuring second messenger production such as changes in cAMP production in the presence or absence of forskolin, changes in inositol 3-phosphate levels, changes in intracellular calcium levels or by indirect measures of receptor activation including receptor stimulated mitogenesis, receptor mediated changes in extracellular acidification or receptor mediated changes in reporter gene activation in response to cAMP or calcium (these methods should all be referenced in other sections of the patent). Receptor activation could also be monitored by co-transfecting cells with a chimeric $GI_{q/13}$ to force receptor coupling to a calcium stimulating pathway (Conklin *et al.*, Nature 363: 274-276, 1993). Neurotransmitter mediated activation of receptors could also be monitored by measuring changes in [35 S]-GTPKS binding in membrane

fractions prepared from transfected mammalian cells. This assay could also be performed using baculoviruses containing nGPCR-x proteins infected into SF9 insect cells.

[000317] The neurotransmitter which activates nGPCR-x proteins can be purified to homogeneity through successive rounds of purification using nGPCR-x proteins activation as a measurement of neurotransmitter activity. The composition of the neurotransmitter can be determined by mass spectrometry and Edman degradation if peptidergic. Neurotransmitters isolated in this manner will be bioactive materials which will alter neurotransmission in the central nervous system and will produce behavioral and biochemical changes.

Example 13 - Using nGPCR-x proteins to isolate and purify G proteins

[000318] cDNAs encoding nGPCR-x proteins are epitope-tagged at the amino terminus end of the cDNA with the cleavable influenza-hemagglutinin signal sequence followed by the FLAG epitope (IBI, New Haven, CT). Additionally, these sequences are tagged at the carboxyl terminus with DNA encoding six histidine residues. (Amino and Carboxyl Terminal Modifications to Facilitate the Production and Purification of a G Protein-Coupled Receptor, B.K. Kobilka, *Analytical Biochemistry*, Vol. 231, No. 1, Oct 1995, pp. 269-271). The resulting sequences are cloned into a baculovirus expression vector such as pVL1392 (Invitrogen). The baculovirus expression vectors are used to infect SF-9 insect cells as described (Guan, X. M., Kobilka, T. S., and Kobilka, B. K. (1992) *J. Biol. Chem.* 267, 21995-21998). Infected SF-9 cells could be grown in 1000-ml cultures in SF900 II medium (Life Technologies, Inc.) containing 5% fetal calf serum (Gemini, Calabasas, CA) and 0.1 mg/ml gentamicin (Life Technologies, Inc.) for 48 hours at which time the cells could be harvested. Cell membrane preparations could be separated from soluble proteins following cell lysis. nGPCR-x protein purification is carried out as described for purification of the β_2 receptor (Kobilka, *Anal. Biochem.*, 231 (1): 269-271, 1995) including solubilization of the membranes in 0.8-1.0 % *n*-dodecyl -D-maltoside (DM) (CalBiochem, La Jolla, CA) in buffer containing protease inhibitors followed by Ni-column chromatography using chelating Sepharose™ (Pharmacia, Uppsala, Sweden). The eluate from the Ni-column is further purified on an M1 anti-FLAG antibody column (IBI). Receptor containing fractions are monitored by using receptor specific antibodies following western blot analysis or by SDS-PAGE analysis to look for an appropriate sized protein band (appropriate size would be the predicted molecular weight of the protein).

[000319] This method of purifying G protein is particularly useful to isolate G proteins that bind to the nGPCR-x proteins in the absence of an activating ligand.

[000320] Some of the preferred embodiments of the invention described above are outlined below and include, but are not limited to, the following embodiments. As those skilled in the art will appreciate, numerous changes and modifications may be made to the preferred embodiments of the invention without departing from the spirit of the invention. It is intended that all such variations fall within the scope of the invention.

[000321] The entire disclosure of each publication cited herein is hereby incorporated by reference.

What is claimed is:

1. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of: SEQ ID NO:59 to SEQ ID NO:116; said nucleic acid molecule encoding at least a portion of nGPCR-x.
2. The isolated nucleic acid molecule of claim 1 comprising a sequence that encodes a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116.
3. The isolated nucleic acid molecule of claim 1 comprising a sequence homologous to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58.
4. The isolated nucleic acid molecule of claim 1 comprising a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58.
5. The isolated nucleic acid molecule of claim 1 wherein said nucleic acid molecule is DNA.
6. The isolated nucleic acid molecule of claim 1 wherein said nucleic acid molecule is RNA.
7. An expression vector comprising a nucleic acid molecule of any one of claims 1 to 4.
8. The expression vector of claim 7 wherein said nucleic acid molecule comprises a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58.
9. The expression vector of claim 7 wherein said vector is a plasmid.
10. The expression vector of claim 7 wherein said vector is a viral particle.
11. The expression vector of claim 10 wherein said vector is selected from the group consisting of adenoviruses, baculoviruses, parvoviruses, herpesviruses, poxviruses, adeno-associated viruses, Semliki Forest viruses, vaccinia viruses, and retroviruses.

12. The expression vector of claim 7 wherein said nucleic acid molecule is operably connected to a promoter selected from the group consisting of simian virus 40, mouse mammary tumor virus, long terminal repeat of human immunodeficiency virus, maloney virus, cytomegalovirus immediate early promoter, Epstein Barr virus, rous sarcoma virus, human actin, human myosin, human hemoglobin, human muscle creatine, and human metallothionein.
13. A host cell transformed with an expression vector of claim 7.
14. The transformed host cell of claim 13 wherein said cell is a bacterial cell.
15. The transformed host cell of claim 14 wherein said bacterial cell is *E. coli*.
16. The transformed host cell of claim 13 wherein said cell is yeast.
17. The transformed host cell of claim 16 wherein said yeast is *S. cerevisiae*.
18. The transformed host cell of claim 13 wherein said cell is an insect cell.
19. The transformed host cell of claim 18 wherein said insect cell is *S. frugiperda*.
20. The transformed host cell of claim 13 wherein said cell is a mammalian cell.
21. The transformed host cell of claim 20 wherein mammalian cell is selected from the group consisting of chinese hamster ovary cells, HeLa cells, African green monkey kidney cells, human HEK-293 cells, and murine 3T3 fibroblasts.
22. An isolated nucleic acid molecule comprising at least 10 nucleotides, said nucleic acid molecule comprising a nucleotide sequence complementary to at least a portion of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58.
23. The nucleic acid molecule of claim 22 wherein said molecule is an antisense oligonucleotide directed to a region of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58.

24. The nucleic acid molecule of claim 23 wherein said oligonucleotide is directed to a regulatory region of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58.
25. A composition comprising a nucleic acid molecule of any one of claims 1 to 4 or 22 and an acceptable carrier or diluent.
26. A composition comprising a recombinant expression vector of claim 7 and an acceptable carrier or diluent.
27. A method of producing a polypeptide that comprises a sequence selected from the group of sequences consisting SEQ ID NO:59 to SEQ ID NO:116, and homologs thereof, said method comprising the steps of:
- a) introducing a recombinant expression vector of claim 8 into a compatible host cell;
 - b) growing said host cell under conditions for expression of said polypeptide; and
 - c) recovering said polypeptide.
28. The method of claim 27 wherein said host cell is lysed and said polypeptide is recovered from the lysate of said host cell.
29. The method of claim 27 wherein said polypeptide is recovered by purifying the culture medium without lysing said host cell.
30. An isolated polypeptide encoded by a nucleic acid molecule of claim 1.
31. The polypeptide of claim 30 wherein said polypeptide comprises a sequence selected from the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116.
32. The polypeptide of claim 30 wherein said polypeptide comprises an amino acid sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116.

33. The polypeptide of claim 30 wherein said sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116 comprises at least one conservative amino acid substitution compared to the sequences in the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116.

34. The polypeptide of claim 30 wherein said polypeptide comprises an allelic variant of a polypeptide with a sequence selected from the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116.

35. A composition comprising a polypeptide of claim 34 and an acceptable carrier or diluent.

36. An isolated antibody which binds to an epitope on a polypeptide of claim 30.

37. The antibody of claim 36 wherein said antibody is a monoclonal antibody.

38. A composition comprising an antibody of claim 36 and an acceptable carrier or diluent.

39. A method of inducing an immune response in a mammal against a polypeptide of claim 30 comprising administering to said mammal an amount of said polypeptide sufficient to induce said immune response.

40. A method for identifying a compound which binds nGPCR-x comprising the steps of:

- a) contacting nGPCR-x with a compound; and
- b) determining whether said compound binds nGPCR-x.

41. The method of claim 40 wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116.

42. The method of claim 40 wherein binding of said compound to nGPCR-x is determined by a protein binding assay.

43. The method of claim 40 wherein said protein binding assay is selected from the group consisting of a gel-shift assay, Western blot, radiolabeled competition assay, phage-based expression cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, and ELISA.

44. A compound identified by the method of claim 40.
45. A method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-x comprising the steps of:
- a) contacting said nucleic acid molecule encoding nGPCR-x with a compound; and
 - b) determining whether said compound binds said nucleic acid molecule.
46. The method of claim 45 wherein binding is determined by a gel-shift assay.
47. A compound identified by the method of claim 45.
48. A method for identifying a compound which modulates the activity of nGPCR-x comprising the steps of:
- a) contacting nGPCR-x with a compound; and
 - b) determining whether nGPCR-x activity has been modulated.
49. The method of claim 48 wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116.
50. The method of claim 48 wherein said activity is neuropeptide binding.
51. The method of claim 48 wherein said activity is neuropeptide signaling.
52. A compound identified by the method of claim 48.
53. A method of identifying an animal homolog of nGPCR-x comprising the steps:
- a) comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequence consisting of SEQ ID NO:1 to SEQ ID NO:58, and portions thereof, said portions being at least 10 nucleotides; and
 - b) identifying nucleic acid sequences of the animal that are homologous to said sequence selected from the group sequence consisting of SEQ ID NO:1 to SEQ ID NO:58, and portions thereof, said portions comprising at least 10 nucleotides.

54. The method of claim 53 wherein comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58, and portions thereof, said portions being at least 10 nucleotides, is performed by DNA hybridization.

55. The method of claim 53 wherein comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:58, and portions thereof, said portions being at least 10 nucleotides, is performed by computer homology search.

56. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:

(a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR-x that is expressed in the brain, wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, and allelic variants thereof, and wherein the nucleic acid corresponds to a gene encoding the nGPCR-x; and

(b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR-x in the nucleic acid correlates with an increased risk of developing the disorder.

57. A method according to claim 56, wherein the disease is a mental disorder.

58. A method according to claim 56, wherein the assaying step comprises at least one procedure selected from the group consisting of:

a) comparing nucleotide sequences from the human subject and reference sequences and determining a difference of at least a nucleotide or at least one codon between the nucleotide sequences from the human subject that encodes a nGPCR-x reference sequence;

(b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences;

(c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and

(d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

59. A method according to claim 58 wherein the assaying step comprises: performing a polymerase chain reaction assay to amplify nucleic acid comprising nGPCR-x coding sequence, and determining nucleotide sequence of the amplified nucleic acid.

60. A method of screening for an nGPCR-x hereditary mental disorder genotype in a human patient, comprising the steps of:

(a) providing a biological sample comprising nucleic acid from said patient, said nucleic acid including sequences corresponding to alleles of nGPCR-x; and

(b) detecting the presence of one or more mutations in the nGPCR-x allele;

wherein the presence of a mutation in a nGPCR-x allele is indicative of a hereditary mental disorder genotype.

61. The method according to claim 60 wherein said biological sample is a cell sample.

62. The method according to claim 60 wherein said detecting the presence of a mutation comprises sequencing at least a portion of said nucleic acid, said portion comprising at least one codon of said nGPCR-x allele, said portion comprising at least 10 nucleotides.

63. The method according to claim 60 wherein said nucleic acid is DNA.

64. The method according to claim 60 wherein said nucleic acid is RNA.

65. A kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association:

(a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-x gene, the oligonucleotide comprising 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-x gene sequence or

nGPCR-x coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and

(b) a media packaged with the oligonucleotide, said media containing information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

66. A method of identifying a nGPCR-x allelic variant that correlates with a mental disorder, comprising the steps of:

(a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny;

(b) detecting in the nucleic acid the presence of one or more mutations in an nGPCR-x that is expressed in the brain, wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:59 to SEQ ID NO:116, and allelic variants thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding nGPCR-x;

wherein the one or more mutations detected indicates an allelic variant that correlates with a mental disorder.

67. A purified and isolated polynucleotide comprising a nucleotide sequence encoding a nGPCR-x allelic variant identified according to claim 66.

68. A host cell transformed or transfected with a polynucleotide according to claim 67 or with a vector comprising the polynucleotide.

69. A purified polynucleotide comprising a nucleotide sequence encoding nGPCR-x of a human with a mental disorder;

wherein said polynucleotide hybridizes to the complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 under the following hybridization conditions:

(a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and

(b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS; and

wherein the polynucleotide that encodes nGPCR-x amino acid sequence of the human differs from the sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 by at least one residue.

70. A vector comprising a polynucleotide according to claim 69.

71. A host cell that has been transformed or transfected with a polynucleotide according to claim 69 and that expresses the nGPCR-x protein encoded by the polynucleotide.

72. A host cell according to claim 71 that has been co-transfected with a polynucleotide encoding the nGPCR-x amino acid sequence set forth in a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:58 and that expresses the nGPCR-x protein having the amino acid sequence set forth in SEQ ID NO:59 to SEQ ID NO:116.

73. A method for identifying a modulator of biological activity of nGPCR-x comprising the steps of:

a) contacting a cell according to claim 72 in the presence and in the absence of a putative modulator compound;

b) measuring nGPCR-x biological activity in the cell;

wherein decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

74. A method to identify compounds useful for the treatment of a mental disorder, said method comprising the steps of:

(a) contacting a composition comprising nGPCR-x with a compound suspected of binding nGPCR-x;

(b) detecting binding between nGPCR-x and the compound suspected of binding nGPCR-x;

wherein compounds identified as binding nGPCR-x are candidate compounds useful for the treatment of a mental disorder.

75. A method for identifying a compound useful as a modulator of binding between nGPCR-x and a binding partner of nGPCR-x comprising the steps of:

(a) contacting the binding partner and a composition comprising nGPCR-x in the presence and in the absence of a putative modulator compound;

(b) detecting binding between the binding partner and nGPCR-x; wherein decreased or increased binding between the binding partner and nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a mental disorder.

76. A method according to claim 74 or 75 wherein the composition comprises a cell expressing nGPCR-x on its surface.

77. A method according to claim 76 wherein the composition comprises a cell transformed or transfected with a polynucleotide that encodes nGPCR-x.

78. A method of purifying a G protein from a sample containing said G protein comprising the steps of:

- a) contacting said sample with a polypeptide of claim 1 for a time sufficient to allow said G protein to form a complex with said polypeptide;
- b) isolating said complex from remaining components of said sample;
- c) maintaining said complex under conditions which result in dissociation of said G protein from said polypeptide; and
- d) isolating said G protein from said polypeptide.

79. The method of claim 78 wherein said sample comprises an amino acid sequence selected from the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116.

80. The method of claim 78 wherein said polypeptide comprises an amino acid sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:59 to SEQ ID NO:116.

81. The method of claim 78 wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: SEQ ID NO:59 to SEQ ID NO:116.

82. The isolated nucleic acid molecule of claim 1 comprising a sequence that encodes a polypeptide comprising a sequence of SEQ ID NO:67.

83. The isolated nucleic acid molecule of claim 1 comprising a sequence homologous to a sequence of SEQ ID NO:9.
84. The isolated nucleic acid molecule of claim 1 comprising a sequence of SEQ ID NO: 9.
85. An expression vector comprising a nucleic acid molecule of any one of claims 82 to 84.
86. An isolated nucleic acid molecule comprising at least 10 nucleotides, said nucleic acid molecule comprising a nucleotide sequence complementary to at least a portion of a sequence of SEQ ID NO:9.
87. The nucleic acid molecule of claim 86 wherein said molecule is an antisense oligonucleotide directed to a region of a sequence of SEQ ID NO:9.
88. The nucleic acid molecule of claim 87 wherein said oligonucleotide is directed to a regulatory region of a sequence of SEQ ID NO:9.
89. A method of producing a polypeptide that comprises a sequence of SEQ ID NO:67, and homologs thereof, said method comprising the steps of:
- a) introducing a recombinant expression vector of claim 85 into a compatible host cell;
 - b) growing said host cell under conditions for expression of said polypeptide; and
 - c) recovering said polypeptide.
90. An isolated polypeptide encoded by a nucleic acid molecule of claim 82.
91. The polypeptide of claim 90 wherein said polypeptide comprises a sequence of SEQ ID NO:67.
92. The polypeptide of claim 90 wherein said polypeptide comprises an amino acid sequence homologous to a sequence of SEQ ID NO:67.
93. An isolated antibody which binds to an epitope on a polypeptide of claim 90.

94. The antibody of claim 93 wherein said antibody is a monoclonal antibody.
95. A method of inducing an immune response in a mammal against a polypeptide of claim 90 comprising administering to said mammal an amount of said polypeptide sufficient to induce said immune response.
96. A method for identifying a compound which binds nGPCR-93 comprising the steps of:
- a) contacting nGPCR-93 with a compound; and
 - c) determining whether said compound binds nGPCR-93.
97. The method of claim 96 wherein nGPCR-93 comprises an amino acid sequence of SEQ ID NO:67.
98. The method of claim 96 wherein binding of said compound to nGPCR-x is determined by a protein binding assay.
99. A compound identified by the method of claim 96.
100. A method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-93 comprising the steps of:
- a) contacting said nucleic acid molecule encoding nGPCR-93 with a compound; and
 - b) determining whether said compound binds said nucleic acid molecule.
101. The method of claim 100 wherein binding is determined by a gel-shift assay.
102. A compound identified by the method of claim 100.
103. A method for identifying a compound which modulates the activity of nGPCR-93 comprising the steps of:
- a) contacting nGPCR-93 with a compound; and
 - b) determining whether nGPCR-93 activity has been modulated.
104. The method of claim 103 wherein the nGPCR-x comprises an amino acid sequence of SEQ ID NO:67.

105. The method of claim 103 wherein said activity is neuropeptide binding.
106. The method of claim 103 wherein said activity is neuropeptide signaling.
107. A compound identified by the method of claim 103.
108. 5A method of identifying an animal homolog of nGPCR-93 comprising the steps:
- a) comparing the nucleic acid sequences of the animal with a sequence of SEQ ID NO:9, and portions thereof, said portions being at least 10 nucleotides; and
 - b) identifying nucleic acid sequences of the animal that are homologous to said sequence of SEQ ID NO:9, and portions thereof, said portions comprising at least 10 nucleotides.
109. The method of claim 108 wherein comparing the nucleic acid sequences of the animal with a sequence of SEQ ID NO:9, and portions thereof, said portions being at least 10 nucleotides, is performed by DNA hybridization.
110. The method of claim 108 wherein comparing the nucleic acid sequences of the animal with a sequence of SEQ ID NO:9, and portions thereof, said portions being at least 10 nucleotides, is performed by computer homology search.
111. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:
- (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR-x that is expressed in the brain, wherein the nGPCR-x comprises an amino acid sequence of SEQ ID NO:67, and allelic variants thereof, and wherein the nucleic acid corresponds to a gene encoding the nGPCR-x; and
 - (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR-x in the nucleic acid correlates with an increased risk of developing the disorder.
112. A method according to claim 111, wherein the disease is a mental disorder.

113. A purified polynucleotide comprising a nucleotide sequence encoding nGPCR-x of a human with a mental disorder;

wherein said polynucleotide hybridizes to the complement of a sequence of SEQ ID NO:9 under the following hybridization conditions:

(a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and

(b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS; and

wherein the polynucleotide that encodes nGPCR-x amino acid sequence of the human differs from the sequence of SEQ ID NO:9 by at least one residue.

114. A vector comprising a polynucleotide according to claim 113.

115. A host cell that has been transformed or transfected with a polynucleotide according to claim 113 and that expresses the nGPCR-x protein encoded by the polynucleotide.

116. A host cell according to claim 115 that has been co-transfected with a polynucleotide encoding the nGPCR-x amino acid sequence set forth in a sequence of SEQ ID NO:9 and that expresses the nGPCR-x protein having the amino acid sequence set forth in SEQ ID NO:67.

117. A method for identifying a modulator of biological activity of nGPCR-93 comprising the steps of:

a) contacting a cell according to claim 116 in the presence and in the absence of a putative modulator compound;

b) measuring nGPCR-93 biological activity in the cell;

wherein decreased or increased nGPCR-93 biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

118. A method for identifying a compound useful as a modulator of binding between nGPCR-93 and a binding partner of nGPCR-93 comprising the steps of:

(a) contacting the binding partner and a composition comprising nGPCR-93 in the presence and in the absence of a putative modulator compound;

(b) detecting binding between the binding partner and nGPCR-93;

wherein decreased or increased binding between the binding partner and nGPCR-93 in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a mental disorder.

119. A method according to claim 117 or 118 wherein the composition comprises a cell expressing nGPCR-93 on its surface.

120. A method according to claim 119 wherein the composition comprises a cell transformed or transfected with a polynucleotide that encodes nGPCR-93.

121. A method of purifying a G protein from a sample containing said G protein comprising the steps of:

- a) contacting said sample with a polypeptide of claim 90 for a time sufficient to allow said G protein to form a complex with said polypeptide;
- b) isolating said complex from remaining components of said sample;
- c) maintaining said complex under conditions which result in dissociation of said G protein from said polypeptide; and
- d) isolating said G protein from said polypeptide.

122. The method of claim 121 wherein said sample comprises an amino acid sequence of SEQ ID NO:67.

123. The method of claim 121 wherein said polypeptide comprises an amino acid sequence homologous to a sequence of SEQ ID NO:67.

124. The method of claim 121 wherein said polypeptide comprises an amino acid sequence of SEQ ID NO:67.

00196PCT1.ST25
SEQUENCE LISTING

<110> Pharmacia & Upjohn Inc.
Vogeli, Gabriel

<120> Novel G Protein-Coupled Receptors

<130> 00196PCT1

<150> 60/195,150

<151> 2000-04-06

<150> 60/195,099

<151> 2000-04-06

<150> 60/195,151

<151> 2000-04-06

<150> 60/195,148

<151> 2000-04-06

<150> 60/195,093

<151> 2000-04-06

<150> 60/195,098

<151> 2000-04-06

<150> 60/230,149

<151> 2000-09-05

<160> 117

<170> PatentIn version 3.0

<210> 1

<211> 661

<212> DNA

<213> Homo sapiens

<400> 1

```
acacagtgtg cacacacgtg cagggacata ccccttccc caactgcctg gcctgcacac      60
ttggcatttc cagtatttct aggaagtgat ggctctgtgc atcctgagcc aatccagctc     120
cgagcctcca aggcacacct gtgatgggca gctggaagct ctgcctctga ggccttcaca     180
cacccacctt cgggtcaaac tgcttctgct gaggaacttg gtgtgtcttc cttctgggca     240
ggaggtcaca tttgagagca caggagcagt gcctgcccc cggaatgtg gctctgggta     300
gaattgcagg ctgagggtt ttgggcagga gagcaccaac cgtgccacac ccacacagac     360
acggtcactg gggccctgca gcaggacga ccgcacttcc caaagggtg ggaagccatg     420
tccagaggag gccatgctct agctcccttg ggcagggtg gctgcaagga ggggtgaagt     480
gggcatcttg aaccagaga agtagaggac tcagcaccag cacaaccagc tcggcgcatc     540
aatacacatt cctctccac ttctcccaa gcctgaaaaa acctcaaacc agcctctttg     600
cagctccctg aggtcatgac tcacgaacca tgctcggggc aggggaaaag aaaagcatcc     660
g                                                                           661
```

00196PCT1.ST25

<210> 2
 <211> 627
 <212> DNA
 <213> Homo sapiens

<400> 2
 ctaaaggagg aatagatgtc tttagaaga aatgaaaaaa taaagtaa atgtgaaaattt 60
 cccttactta tttccaaaca agtgctctc caaaaaatg caaataatta agtttctgaa 120
 atggtgaaca tatcagatta gtagacatat ggcaggagca gcaaatgagc agatcaagtt 180
 gaagtcctag tattaccaat ctgttaatgt tgacaggaag actcattttg actgttcctt 240
 ttatatcaat aaatgagtgg atttcaacta ctctaaatag gaatgctaaa agcagcactg 300
 ctaaaagtgc atatcaaacc aataattttc tgatgctgtt ttggtatatc ctacaaacat 360
 ttgtaggaca acaactcaga agggaaaaaa atatcttatg cctttgaggt ctgtactgaa 420
 tgctaatagca ttgttatatg atgggtttta tacagaactg agaataaatt actttcagca 480
 gctgcactct agacctataa atcgctctga gtactacaaa atccatacaa aggaagaaca 540
 gctggataat ttacaccacc agtattttgtc aaaaaaaaaa aaaaaaagct gaaaatacag 600
 aacctgattt tgtccctttt tcgagta 627

<210> 3
 <211> 297
 <212> DNA
 <213> Homo sapiens

<400> 3
 agggggccctc cagcactggt cttgaagggg tgacagggtc tggggctctga ctcccacctc 60
 caccacttcc cacctgaggg ccctggaatg aatcctttcc tggatctgag ctgccacatc 120
 atcagtga aa atgacaccta tatgggactt cagtgagaac acaaatgcaa cgttcctgcc 180
 acggaacaac ccatgtactc actgggagca ttgagagtag atccacactg attgacacag 240
 ggactccagg cctgacccat gatatgtact ggatacatgg ccatgagtgc tccacag 297

<210> 4
 <211> 649
 <212> DNA
 <213> Homo sapiens

<400> 4
 acaccaaata ctgcttttgc gccttaggct tcagcacatt agcatggctt cctcccttgg 60
 catggtaact gtagctgaac ttggagggtt tgtattaccc attataatta ttacttattt 120
 cacatggaaa acaagaaaat ctttatggga attccaagtt ccccttagga ataccaaaga 180
 gaggaagaaag gctttgagga tggctctgat gtgtgaagtg gtgttcattg tgtgtttcac 240
 tccttaccac ctcaacttcc cattctttat gatggtgaag gaacatgtct ttttgaactg 300
 ctcttttata aagatcattc tctgtttcca cattatttcc ctgtgtcttg caaatctgaa 360
 ttgttgtctt gatccagttg tatattattt tatgacctca aaatttcatt atcaattttc 420

00196PCT1.ST25

agatcatggc agcttgggtc ttcagtcatg tatgagatgt aataacagta ccttagaaat 480
 tcatcagagg aaggaggat cttcaaaacta tctctcttga atgtttgaaa gattccaaga 540
 caatataatc aaataattaa ctagaaaaat cgatatgctc tattagtgtg tctatgtcac 600
 tttgaagatt tttctttttt ttttttcttt ttttttatta tactttaag 649

<210> 5
 <211> 626
 <212> DNA
 <213> Homo sapiens

<400> 5
 aatgctactg ctcctgcata taatagctgc tttgaagtgt ttgtctgcta tgtccaaaac 60
 atagactctt tcaaaagcac tttctgttgt ctcctttttc tcttgcatag gagtcacatt 120
 tttctgtctc ttcacatatt tcatatttat ttttgttgaa aaccagacat tttagataat 180
 gtgttgtagc agtcagata ctgattctct cccccaggag ctgttgtctt tcttacttgt 240
 atatgtgttt agtgacttgg ctggactatt ttaataatgt gtatttcctt gtagtatata 300
 ccatctttta tactaatgtt acttttccga tagtgcagcc ttgggcatgg acagagttat 360
 cctgggatga cagtaacttt taatagggtc ctctatgact atctctttcc ctgatgtccc 420
 tgtaagcta tctgcatctc ttggtatcac acctagcctt tgacttcac taattgtttg 480
 atcattgcct cactgttttt ggagtgccc taaggcataa agtggtccac agtctgatat 540
 aattaaattc agattcttac aagagtggtc tttgaggcca gtccttgagg tttgtgctga 600
 ctctgggagg gctcaaagt ttcctt 626

<210> 6
 <211> 400
 <212> DNA
 <213> Homo sapiens

<400> 6
 cctcacatcc ctttcccctc aaaccctggc aaccccaaac tgttcttgac agcctccttt 60
 ggcatcttct cattttggtg tcagatctca cagcagaatt tcttacctat tatataccag 120
 tgctcagtg tgaagttccg gtttaacttc ttgttaccac gagcccacta tcttgcccca 180
 ataataccct cccccaattc acaaacacac aagcattccc tcctacagct ttgggcctcc 240
 tatctgagtc cttcaggaaa gaagtgtgt gtaactccct tggcagttag ttagacttg 300
 gtccaaggaa gatgagcacc agtcagggca gctgggccct cttctctccc tggccatcag 360
 caaatcagca ctgcccacg atgcccaggc aatgggagcg 400

<210> 7
 <211> 556
 <212> DNA
 <213> Homo sapiens

00196PCT1.ST25

<400> 7
 attactatatt ttcaacctct tttactccag ggactttctat gcacctctc cctcaactcc 60
 ccctcaattt gttctcataa tccccatgac cccagtttt ataacaccac tgtcaggagc 120
 ccaaagctgc cattcattca cttccattag catgactctt catgtacttt ggggtcttca 180
 gtctctcccc ttctcctaatt ttccagggtt ccattctgct tctgtctggc tccctacaaa 240
 gcctgcaaca tcataagcca tttcaggaaa gagcttgatc atcttttgat gaacctgca 300
 ttcattgactc actgccttac ctgtctttgg ctctgcatgt cccagttt ccgtttcttt 360
 ctctggaaag agagattgcc caagagtcct gcacatcagc attactagaa atgcatgcag 420
 accagcttca gctgcttgcc aactctttaa aaaaatgagta aacaattttc taaaggggaa 480
 aaaatctctt cacctcctca caccaactat ttgcataatt cagtgcctt ttataaaccg 540
 tgccattgta taagca 556

<210> 8
 <211> 670
 <212> DNA
 <213> Homo sapiens

<400> 8
 caccgtcctc atcatgatcg tcttcgtcat ctgtgctgg gggccctact gcttcctggt 60
 gtgctggcc gccgccggc aggccagac catgcaggcc ccctcgctcc tcagcgtggt 120
 ggccgtctgg ctgacctggg ccaatggggc catcaaccct gtcactctac ccattccgaa 180
 tccaacatt tcgatgctcc tagggcgcaa ccgcgaggag ggctaccgga ctaggaatgt 240
 ggacgcttct ctgccagcc agggccggg tctgcaagcc agaagccgca gtcgccttcg 300
 aaaccgctat gccaacggc tgggggcctg caacaggatg tctctttcca acccgccag 360
 cggagtggca ggggacgtgg ccattgtggc ccgcaaaaat ccagtgtac ttttctgccg 420
 agagggacca ccagagccgg tgacggcagt gaccaaacag cctaaatccg aagctgggga 480
 taccagctc taagacggtt ggaatggcca gcttatgaag gcaaatctc actcgcatta 540
 tttaatgatg gaagattctg ggggagagtt gtgatttca taaagccaaa catttaagc 600
 tagagacggg ggaggttac cactttcccc aaacaacata aaagacaatg tcccttcttt 660
 caaaaagtgc 670

<210> 9
 <211> 1860
 <212> DNA
 <213> Homo sapiens

<400> 9
 cgctcctgcg taaacacgcg gttccctcgg caacgctgga acccacgtca aaggctccgc 60
 cagggtcccca gcgaccgcca cccctccggc cgagccagc tccccgggc ggccgctagc 120
 ccccgcccc gagccaccac tccgacctag cggccgccc ccccggtgcg ggatgaggag 180

00196PCT1.ST25

```

atccgcggcc gccactgggc cccatggagg agccgcagcc gccccgccca ccagcgagca 240
tggccttact gggcagccag cactccggcg cccctccgc ggccggccca cctggcgga 300
cttcctccgc ggccacggcg gccgtgctct ccttcagcac cgtggcgacc gcggcgctgg 360
ggaacctgag cgacgcaagc ggaggcgga cagctgccgc tcccggtggc ggcggccttg 420
gcgggtccgg ggcagcgcg gaggggggg cgcggtgag gcggccgcta gcgacggagg 480
cgcgccgct gctgtcgac ggagctgag tggcgccca ggcgtcgtc ctctgctca 540
tcttctgct gtctagcctt ggcaactgc cggtgatggg ggtgatttg aagcaccggc 600
agctccgac cgtaaccaac gccttcaccc tgctcgtgc cctatcgat ctgtcacgg 660
cgctgctctg cctgcccgc gccttcttg acctctcac tccgcccggg ggttcggcg 720
ctgcccgcgc cgcgggggcc tggcgggct tctgcgcgc cagccgctt ttcagctctg 780
gcttcggcat cgtgtccacg ctacgctgg cgctcatctc gttggaccgt tactgcgcta 840
tcgtgcggcc gccgcgggag aagatcgcc gccgcgcgc gctgcagctg ctggcgggcg 900
cctggctgac ggccctgggc ttctccttg cctgggagct gctcgggcg ccccggaac 960
tcgcggggc gcagagctt cacggctgc tctaccggac ctccccggac cccgcgcagc 1020
tggcgcggc ctacagctg gggctggtg tggcctgcta cctgtgccc ttctgctca 1080
tgtgtctctg cactaccac atctgcaaga cggtgcgcct gtcggacgtg cgcgtgcggc 1140
cggtgaacac ctacgcgcgc gtgctgctt tcttcagca ggtgcgcac gccaccaccg 1200
tctcatcat gatcgtctt gtcactctg gctggggcc ctactgctt ctggtgctgc 1260
tggccgcgc ccggcaggcc cagaccatgc agggccctc gctcctcagc gtggtggccg 1320
tctggctgac ctgggccaat ggggccatca accctgtcat ctacgccatc cgcaatcca 1380
acatttcgat gctcctagg cgcaaccgc agggggcta ccggactagg aatgtggacg 1440
ctttctgcc cagccagggc ccgggtctgc aagccagaag ccgcagtcgc cttcgaaacc 1500
gctatgcaa ccggctgggg gcctgcaaca ggatgtctt ttccaaccg gccagcgag 1560
tggcagggga cgtggccatg tggggccga aaaatccagt tgtactttt tgccgagagg 1620
gaccaccaga gccggtgacg gcagtacca aacagccta atccgaagc ggggatacca 1680
gcctctaaga cggttggaat ggccagctta tgaaggcaaa ttccactcg cattatttaa 1740
tgatggaaga ttctggggga gagttgtgga ttcataaag ccaaacattt aaagctagag 1800
acgggggagg ctaccactt tccccaaaca acataaaaga caatgtccct tctcaaaag 1860

```

<210> 10
<211> 630
<212> DNA
<213> Homo sapiens

```

<400> 10
tctcaaaaaa taaataaaaa cactgtaca tacaaggc cctggggga cagctggggc 60

```


00196PCT1.ST25

```

ataagtaggt gtcagccata catcagagca gtgtgcctgc cctgagctgc ttgggggtga 120
ccagcctggt gtccagaaat gcctgctgga gggagtcgtg gtacaggaaa ccttgtgctc 180
ttagaaggtc tcctgagagg ccctgcaaag ccagagtccc tcttagcagc tcagatcagt 240
gctatcaaag tatagctcgg ggattgctgc cagcatataa acttttactg gtctgcagcg 300
agataagtac agaaattgaa agtaagcatt tagaaacttt tataacaatt ttacaaggtc 360
ttgtcaaatg ttattaaaac aaagctgagg ctggaatttc acctttttca tttcgttttt 420
ttcaatttaa acaaattgta gtaaaatata ggtaatatata atgtaccatt ttagccattt 480
ttgagcgtac aatttagtag cagtaagtgc tttcacaata ttgtgtaacc actagtatta 540
tatagtatat atttttaaaa ttttacagaa gtattaagtt agcagcagat taaacatttt 600
tttcttaaat tgagcttgag aagcgtggc 630

```

```

<210> 11
<211> 683
<212> DNA
<213> Homo sapiens

```

```

<400> 11
agagagcaga ttgccctgtg taggtcaggt ctgggttctt tctagtccag agtagggaag 60
aaacaggaaa gagggctggt gttgaaggac cttcagccac gagaagggct gtgtaccatg 120
tagccctctg gggaggcaca aaaaggctca ccattttctg aaaatgacta gactgcagga 180
tccacgtgag tgtgactatt gcattcatga ccttatccac agggcctcac aaggcgctg 240
acatgcagta ggctccagat gcatatttat tataaagtga atagtcctta agctgcaggg 300
tcccttctat ttgcattcta agaaatagtc acttttatgc ctaattttgt atttgagtt 360
ttataagttt tataagaggg tctcccaaat agtataaact tcaagcccca caaaacctat 420
gtttgcctcc cataggcatg caataaatgt tcgtggatct aatgagtaac aagaaaaaga 480
aggaacaaaa ccctaacccc tcccctaccc aaaccagtgg caaccgggga ggatcaaatt 540
caaccttgat cagtcagagg cagcattcct aaattattcc caagcagcaa tagacaatga 600
tttaacctaa ttaattcagc cagttaaaag cttagttctt acttgccaac cgaaggcttg 660
aaggcaaaat gtgtttaagc ctc 683

```

```

<210> 12
<211> 481
<212> DNA
<213> Homo sapiens

```

```

<400> 12
ttcaggcaga tgtcagttaa aaacttacct ctgcacactg caaaaactgt atagccctga 60
acagatactt ttcttgagca tagttccttt gtctctaaag caggcataat tgccaatgtg 120
gggatgatat ttagaaatct gaactgatgt ttattctcta ggggtcttct catttgagct 180
gggattggag atgtctagtg tctcagagca gcaataagaa aacagaaacc tcttccagct 240

```

00196PCT1.ST25

```

tctgacatcc aaatgtcaag ctcttaggag aagaatggaa agtcctcaag aaatgcaaat 300
agctttggca gaatagctga tgaagaccac ctctcccccc tccagaaagg cattggttcc 360
ccattcatgg aaaagggaaat gtagagagag attagataat agtacatcca taaggttcct 420
ggaatctgca tctgaggaag aggggctgca gagacccag ctgttatcta taatccctcc 480
t 481

```

<210> 13
 <211> 693
 <212> DNA
 <213> Homo sapiens

```

<400> 13
gcactagggc aaagtcaaga catacgggtg tccagttcta gctttgcaac taactggtta 60
tatattttta agttacagtc actctgccc agtttctca tttttaatag agtgggtag 120
aactagataa atactttcat ttgtcaagc tctaaattct gacttcagga aaaaaccata 180
aggcactgga ggtttattca taggttttct tgctgacccc gtccctctct gtttcttcaa 240
ccaccacaag acaatcaact tcctgattg gagattggaa cagggtgtgt ctaattctaa 300
atgcatcact taactattag ttccaactct ctggggcttc cttcaaatag gggaattaga 360
ctgggtctcca atctctttgt acagatgagt aactttattt acccaaagat ttagtattaa 420
cagtcgggag caggagggag aatacttatg agacaacagc catttccaca gtggagagga 480
atggtttggt cccaatagaa gttaccagat ttcagtccca ttgccaaata gatattatga 540
gcaaggaaga aatctatagt agtaacttaa gaccaccaga aagatcaaag ccagaggggt 600
gagggtatgg caataaacat tagacatatc tctaaccctc tttgtttga aatactcatt 660
accctgtggt actgggaata cctgtgccta caa 693

```

<210> 14
 <211> 733
 <212> DNA
 <213> Homo sapiens

```

<400> 14
ttctgccatc gcaaggggag ggaagagcac ctaaagggt tatgagaggt ttgactgacc 60
aagggaggga acagaacaca ttcctttcca ttggctagga ctcggtcaca tggttttccc 120
tcattattag tgaggcttgg agaattcatc tatttgtgag ccaggaaga agagaaaaca 180
aattgtggtg aacatttagc agtctctatg acaatagtct gtatgttgac tgcaaagggt 240
gatgaacaaa accaagcctc ctttaaagca atacaatctg gcagagtcctc tgggttatca 300
ttctgaacat agatgcttat tgtcaagag ttaagaaaat tagcatgact gcattccagt 360
tctataaatt taatctttat tcagcatatt gtcatccaca tgtcttaaaa aataaaataa 420
aaaacaaaaa acctagtaac tacgttttat atagcaagga acactcatat atatcacttc 480

```

00196PCT1.ST25

attgtatcct tacaacaatc ctgtgcagta tatgttttac tccctttctt ctatgttttg 540
 tatataaaga aatgagcccc agggagttga atggcttgcc ccaactagtg aagctaaaac 600
 tccaatccag gtctttttat ttccaaatcc ataatctaca accatctgta gagagttata 660
 attaagagat atgaatggtc aggggccttt ccatttcagt gcaagtctgc ccagctccaa 720
 ctaccagcat ctg 733

<210> 15
 <211> 694
 <212> DNA
 <213> Homo sapiens

<400> 15
 aatcctgcat ttcccatgct ctgggggtgag aaggaattag ctgggagcca attagcaatc 60
 ttgtcagaag caaatgaatt ttgaataaac tggatactta aactgaaatg agaccattga 120
 aaccagaaga gcctgagatc catcagttag aggaaataaa agaagtggca ttttccttgc 180
 catctgggtg cagtgtgagt gattttttata atcctaccac atttttatcc ctgcttcctt 240
 taaactgtag gacccaagga acctggctgt tttgttcaac atgatgtgac ccataaccta 300
 accaggccag gcacaaaatt ggcctccaat aagtagtgga tcaaagtatg aatggataaa 360
 ctgaatgaat gaagccaaac ttgaatttct ccatagctta tccaaatggg aatggtaaaa 420
 atcataagct ttgagaaga gaacttatta agaagcccta catcagtcac gactggcatc 480
 attggttagt ttacccaatt ttctccttcc cttcatcttc ctaatgcaac tctggtttgg 540
 gctgcagtat acccagttag aatacgccct cccagagtc tcatccagct ggagatgac 600
 aagtcaccag ttctagctaa taagttgcca gccaaagtat tcaggatgag acttccagaa 660
 aaagcattgt tttcctgata acaatggaca gacc 694

<210> 16
 <211> 674
 <212> DNA
 <213> Homo sapiens

<400> 16
 gactgttatt actgaaagtc attgctttta gactcttcca actacagagc aaggaagttt 60
 atgtgtatat agtaatctgt gaatatacac atacatacac atatttctat atgtaatcat 120
 ccatatttaa attaagtaga atatgagttc atactgatat ctccaatcct aatcagttac 180
 cacagggatt attccggcct ttttcccttg gaagtttgca actcctgctt caacagttag 240
 aaatctggct tccatattca tttgcttaat tgttcaattc cagtacacat aaatggtggc 300
 ttcagaatta ataacttata cctccatggg aaataacttt attaaactaaa gtacagcact 360
 tatgtatagt actttttgaa ttttttagact tagagattcc tcttcttttc caaagttact 420
 taggtcagaa ccattttcca ttcttcagtg aagttgtctt atgtatttgt aatacagtta 480
 gattgttctg tcatatggtg cattccatcc tgggatttcc tatctctttt tttaacattt 540

00196PCT1.ST25

gcatatatta agtttcattc ttttgtgctg tatcattcta tgggtttcaa ttaatgcata 600
 gtgtcatgaa tctgccacca taggagcatc atacagagta gtttcaccaa cttaaaaaat 660
 tccctatggt ttac 674

<210> 17
 <211> 645
 <212> DNA
 <213> Homo sapiens

<400> 17
 cgttccttcc tctgtgtcat aatggacatg atgatatgtg gctcactcag taaacattct 60
 gtgtctggaa ggatttgatt tgtccttttc ttgaggcaac aattttgagg tgatttgaaa 120
 aatctttctt gaaaaattaa aaaatttttc taattaaaaa taatgcaagc tcattagaaa 180
 aaaattgaaa aataaataaa agcacaattt ttcttaacca cttaaagatg accattgtta 240
 gttttttttt ttttttggtg cttttttccg tttccaatct cttttctatt aaaacttctg 300
 aaatgtgatt gtagcaatga cgcataaggg gcccttgaca cattgagaaa ttataaata 360
 cgctggcttc ttgtcttgct tttgtcccca gcttaactgg gaactctttt tctatatctt 420
 tgaaactcca aatcctagat aattcttcaa ggtcaagctc caatgtcttg ctggattctt 480
 ctcagcagga attgatctat tttctctgta tttttgtgcc acaggatcta tgactctctt 540
 atggcaacta ccaccttctg ccttatatta cgatttttga atcttccaac aaagtctaata 600
 tttttttttt tcaaatgaag tctcgctata ttgcccaagc tggag 645

<210> 18
 <211> 707
 <212> DNA
 <213> Homo sapiens

<400> 18
 atgtcatggg aaatgcaaga atatgtgtcc agcatggaag ggaatcagta tggaagtctt 60
 ttgataaatt gtggcattta tcaactaacat tgcctcaaaa ctttagacta cctgccatat 120
 acaaattaga ggtgaaaatt acttccatgt aatatacaag ccaacacaaa gaatcctatc 180
 ccagtttctt ggatggatag gcaagaatct gggtaagggt tattgtgcaa taatcctctt 240
 ctctcttcta taggccagga tttaagttaa cctcaaaaat ggaaaatttt ggctgggaaa 300
 attacatgtg ggaagacatc ttcagtggag attttagtaa ttacagtttc agctatgacc 360
 ctaccctttt tctactagat tctgccccat gttggccaga atccctagaa atcaattatg 420
 ttttgatcat catctatgcc ctgatgtttc tactgaacgt gatgtgaaac tccctgccga 480
 tgctggtcat cttattcagc tgagtcagcc actgtcaccg atgtctacct gctgaccctg 540
 gccttgcccg acctgttctt ttccctgaca ttgcccatct tggctgcctc caagaatgaa 600
 tggctgggat ttttggcaca atctgtgccca ggtggtctag ctctgaag aagtcaactt 660

00196PCT1.ST25

ctacggggggg tatttacta ctggcctgcc gcagcatggg actgtta 707

<210> 19
 <211> 680
 <212> DNA
 <213> Homo sapiens

<400> 19
 tatgatattt cagccatggt gctgaacatt tccaaacagc ataaatgcac catgtgtgta 60
 tgttttcctt tgggatgctg tgcttagagg gtagcagaca gggtgcaaag tgagaaggac 120
 ctggctctgc acccaacact gccagtattg aatcctgact ccatcatctg ggagctgtgc 180
 aacctacgca aggtacttgg cctcagtttc ctcatcatcc ccatggcatt tttgtgagaa 240
 ttaaatgagc tgaaaccttg aaaccccttc aaacagcagc tggcacagag gaagcacaca 300
 atcaatgtca gctgtactct tcctggcagt gtggagatcc cagctctgcc cctagctagt 360
 cacttctctt cttggaatct cagttccttc atctgggaaa tgggagcaga tgtgaaaagg 420
 ggcagggtga gaatacatat gaaagtgtg gctcctgggt catagcaggc acttaataat 480
 gatacacttt tccatcttct gccttcccca gggatgcatt gtgccatgta agagagagag 540
 cctccagggt tggcgagagt ttttgatcca ggctttttca ggtgtcaaag atgagctggg 600
 tgattctcca tagattttcc tttctaacag gtgacagttc tgtttcagaa atactgtgga 660
 tgttcaggtt tacagcacat 680

<210> 20
 <211> 479
 <212> DNA
 <213> Homo sapiens

<400> 20
 tttatgctca tgtagttctt tccaagaaga gaaattacag agtcaaattg tagaaatatt 60
 taaaaatctt tggcacacat aaacagtatc catataattt ataccatctt ttagatgagt 120
 tttaacacca aatgatagaa atctcagttt catacagatt tggtagggctg gaaccaaata 180
 cttgcctgat aggctgtccc ctctgtcttc cttagctgttc tgggaaaggc agttcctggg 240
 aagaactctc cctacggccc ctttcatctc actgttcctc agggcataga taagtgggtt 300
 gagcagtggg gttcccaatg tgtacaccag tgagatgaac tgatcttgct tggggttgta 360
 gctggagctg gggcacagg acatgaaggc acagcagcca tactgcagca gcaccacagt 420
 gaggtgggaa gagcaggtgg agaaagcccg gtggcggcca gcagccgagt ggatcttga 479

<210> 21
 <211> 709
 <212> DNA
 <213> Homo sapiens

<400> 21
 tttaagccac ccagtctgtg gtgttctgtt atggcagccc aagccagcta ctacagggtg 60

00196PCT1.ST25

```

ggacgagggg aggagcatgg cctctgctgg aagtgcaggc aaatgatccc ccaggaacaa 120
tgatgggagc ttctgattgc tctcattatc tctgcaaagt aggaagaaag attcatcagc 180
tgagcatgag gatggtagaa aacatctttg ggaaatttca gaagtgaagg aaggcataaa 240
atagtcattct aaaaaaagca ggaaaggga aagacagaga aatccagtat gagtcccagg 300
actccaggaa gcatcaggac ccacttgaaa ttgccaatgc tgaatttaaa atgaggccag 360
tctgtacaga agcacttctg gaatttgcta acagctaaat agagtagaat caatacttta 420
gagaatacga gtaaccaaag gaataaaatt aactgatcaa cttttgtggt ttttactatt 480
aatattttct tcagtgtaaa tcatagctgc ctgaattcct gaaccctct tatataaatc 540
taaaaagctc tggtttatca tggttgaaaa ttcatggcta acttatcagg caaactgtcc 600
ctaaagcatt ttttgaatag ctttagtadc aagatggtac tgagtgtaca tttcatttcc 660
ctgcttaaag gaaggcttag ttattttaaa ccaagtctta tttttatag 709

```

<210> 22
<211> 517
<212> DNA
<213> Homo sapiens

```

<400> 22
atttctggat ttatgcctcc cctgacccat tccaggattt accccaaacc ttccacactc 60
tcttctaaca gggaaagttc tgttatgaca caatagtact tattaagaca gatttacctt 120
ctaagtctca ggacagcatt tcacaaccag aaataactgg tcacatgaag aaccaggagt 180
ctggtagtag tgaaattcat tttccttctt gaaaaagtgg atcaaaggat tcaaacagca 240
agtgggtgaat caatgaaaag tggtaaaatg gtgaggaaaa aatgttacta aaagatgacc 300
tcaagattac tgggtgcatat gaattgcttt tttatatagg aaaatactgg ataatttctt 360
attgtcatag tataattaga agcaatttca tgtgttcatt ttgccacatg agtttaaatg 420
gaatagattt gggtccctct ctaacatgag ttcagtgtct gaacttgggc aaatttctaa 480
acaattctga gcttcactac ctctgcttga aagtgag 517

```

<210> 23
<211> 695
<212> DNA
<213> Homo sapiens

```

<400> 23
ctttggaatt ttattctaag catcaatcaa gaggtatagt acgagaaagg tagaacatgt 60
aattataaat tcaggattca ggaagtttat ttttctcttc tttttaattc tctcaaatg 120
atcttgattc ctgcaaagtg ttagtatatc tggtaagtaa gagtctatct cttttaaact 180
tcatctgtat taaccagctt tatatgacca aaatgtcccc caaatttaaa tctttgcaca 240
gtaaggcctt atatgtacac ctggcctcat ttcaaaagac taaagcagtt gttctcaaat 300
tcagctgcac attaataata actggaaaac tgtttaagct cctgatgaca aagccacatg 360

```

00196PCT1.ST25

tgagactaat ttatgctgaa tcaactgggcc aaggacccag gtatcagcat tttttaaaac 420
 tatagaggaa taaccagggt tgagaaccac tgcacaaaat ggtaaatagca actttttattt 480
 aagttatttt ttttaaataa ataatgggtg aattgatact gatcttagta ccaagtcagt 540
 gcaatttttt cagacttaga gaattcatcc tggcattgag attattaaag aacctagaaa 600
 tccaagtgtt tttgtttata tttttcctgt aaatattaga gtatgctagt gctcatcctt 660
 atttgataat tttggaaaaa tatattaaaa cattt 695

<210> 24
 <211> 677
 <212> DNA
 <213> Homo sapiens

<400> 24
 attatcattt gaatgttgat attacatcat atacaaattg attgcaacat agttatttaa 60
 tgtaacattc ttttttaaaa gataaattta tcagaatcat acattgctac agtagtctcc 120
 cttatccaca ggttcatttt ctatggtttc agttacctac tgtcaacaag gatccaacaa 180
 tattacatgg gaaaatcaca gaaataaaca gtttgtaagt ttttaattgt gcgctgttct 240
 gggcaacgtg ataaaatctc atgctgttcc tctctatctt gcctgaacat gaattatcct 300
 ttgtccagta tatccacact acatatgcta ccttcccatt catcatttag tagctgtttt 360
 gattatctga tagaaaaaac acacagtata tatagagttt tttatggggc aagggaaaac 420
 tttctctttg tcctctgaag attcactgaa aactcaactc acaagggcag actaatagga 480
 atgaaggtaa aaaaaaaaaa atattaacat caatggagat aactacagag tgattattcc 540
 attgccatca atggactaca gtggcttaaa tatcgttttg aggttacaaa aagagtggaa 600
 gtcttgggat cttggcaaaa caggttatgg gaagaagaga agagaaaccc tggtagcaa 660
 aggtcatctt gtgatgc 677

<210> 25
 <211> 361
 <212> DNA
 <213> Homo sapiens

<400> 25
 ttttttcccc ctgagtgttt ctctcatgct ttcctccaaa tggagatgga gaggtttcac 60
 ctcaactttc tctaactctc cctagttttt tggtttcttt tcctccacat ctaaaagtgt 120
 gcgaatgtc ccttttagcac atagaaaatc ttttcttgac cctgccacct acttaactaa 180
 aatccacac ttttcttctt cttttaagat ttcttttata atggtgtgtg tcaatggcca 240
 catccacctt atccattcct tcttaaagt ccagaaaaac ggttttgttt cctgttactt 300
 taatggaatt atttttccaa agatcaacag gactttccct caagcccaat ccagtcggtg 360
 g 361

00196PCT1.ST25

<210> 26
 <211> 459
 <212> DNA
 <213> Homo sapiens

<400> 26
 atacatgata aggtacatgg atccagggga aggatgaagg gcagtgtggg attgcttttg 60
 aatttctcca aactcgccca taaaagcaga caggacaaac taagataact aaacaaaaaa 120
 acccacagac aaaactatta caaaccccaa aagaagtgtg gtgggaacaa acatctgata 180
 gaatcagaca cattactggt gaccggacat aagccctgtt aatgagaagc ttacatttag 240
 gagagtcaat taagtacacg ctatacacia cctaaagtgg taaatgctac cttgggttatt 300
 caacttcact gttacatgcc ttgaagtgtg ggtgactg gcctgaacca ttctggttgt 360
 gtttgattcc ttaggatgcc accaacaat aacattgaga aatacccagc tacttttcat 420
 tgttctccaa tggcagcaaa gtacaaatga tctctatga 459

<210> 27
 <211> 625
 <212> DNA
 <213> Homo sapiens

<400> 27
 tccaggcagt atttccatg acaatgagga aggtaagtct gcaacagaag aacaatggca 60
 gaaattttta gaaaagttta ccgcctggga ctatgactca ctttttggga gaaaatgtga 120
 ctaacccttt gtaagagctt gttgagagct cactttcctg ggaggagtcg ggagaagggg 180
 agcatcagct gacgaagagg tgaaggaggt acccaacaag aaaagcgtag aaggaccagg 240
 gatttggggg cggttcttcc tcctgattcc aagggatggc ataagatatt gccaagtga 300
 ggaagcgaag tagagccagc aaaggaaggt gaactgttgt ttcattagaa ataatatgtt 360
 gtgataatta tacaagtac taattagtaa atttcttcc aacctcgaca ctccaaaaat 420
 ccctgtactt atatcccgaa ggcctcttct tcccaagct ggaagacagc gtcactcatt 480
 agtcaccac tgacacagga gtaacagaga ctacaaatat tggacaggac ataagtggg 540
 gtcaagcatc tggatgcaga tgcacagag gatgcaagtc ttccagctc tcatggactt 600
 tgcgacagat gcacagagtg aggta 625

<210> 28
 <211> 601
 <212> DNA
 <213> Homo sapiens

<400> 28
 attaaatgcc tgaactccct tcagctctga aactctgtgg tgtattctct gaggacatta 60
 cctctctaag ggacccaaat taaacagctc acaccatcca tcatTTTTct gtctgaggtt 120
 ttatttcctt aatcagattt gggtaaattt tcagcctctc tctgtctctt acttccaatc 180

00196PCT1.ST25

caataaaacc tgtatggatt tgttttgtat ttcataat gtcattattc attctaagag 240
 tcaactgcctg accatttccc tgcctatagc ataattagct attaaaaagc tacactggca 300
 tggttttcaa acttgcattc tctttttctg aggtgggattg attctaaact gattaaaata 360
 tctcagaatt tccaatacaa tttttaaaat gcaacagatt ttcaagactg cctcatgact 420
 ctgccaagcc aaggaggtta gctgccaact ctctctgact gccaaggaag ccaataaat 480
 aatcctgatg gtggttttaa aatgagaggc aagtgcccat ttcttaggtt gacagtgcc 540
 ccctacacat tgacttctcc aggttttgta agacaccaag ggtgatgttt cagattttcc 600
 c 601

<210> 29
 <211> 597
 <212> DNA
 <213> Homo sapiens

<400> 29
 tccacattct ttctaaagtt ctgagctttt ccatgggctt ccatggtagg gaaagcacat 60
 ggcctgggtg tgggtagagc aggtgcccgc atttatatgt atggttcttt gcaagtctgg 120
 catttgtaaa atgggtgatg cttgtattgt gttattttat tcaatcatgt aatagaagat 180
 gcacataaga ttattttgaa aagtatgcct tccattttca tgctgagaat aatgcaggaa 240
 gttcagtgtg atgcagttat aataaaatag tagcaaaaca atattttgct ttaaatcatg 300
 gaattagcaa gtaaagacta attggaagcc aatcttttgc aaatttttta aatgtaagtt 360
 tatttgagg atatgacttg ttggcccaga gtacatataa agaacaaaag agtataatta 420
 acaacagttt caaatatgga cttaccaggc atcttgataa aatcagtatt gacatgtatg 480
 tgaatgcaa cattgtgttt ttccaattca atactatgtt atgccataaa actggttagca 540
 gttatgaaaa ttagaattgg ttaaaaactg ttgaaatctt taaatttttc ctgttta 597

<210> 30
 <211> 618
 <212> DNA
 <213> Homo sapiens

<400> 30
 tacgtgggtt tccctatcgt cctcttcatg agttctttgt gaaaacagaa agactgagtc 60
 tgccaataac cagcaagaga acaagataaa ataaataaaa ttaaccataa gactttaaca 120
 tatgacaaac aactggttaag gattttcaaa atcttttggc caactttgat ggtatttttc 180
 catacaatga actctaaaat atgaaaaacg tacatccata ttttagatat aaaagtctct 240
 tgcacaggcc agaaaatgaa actttaattt aagcaataaa attccccctt gtagactgca 300
 aatggagaac atgctatcta gtttcatttt tcttcaactt acataaaaat gaaacaatgg 360
 ttaatgttct ggcggcatct ctaaacatat tcagtgaac aaaatttcct tacaatgtc 420
 aacagcttac aacaaataac attttatcct gttaattat ttagaaacaa aatcagttat 480

00196PCT1.ST25

gctgagatat gtttgcatgg gatttatata ctctgatcat agaaacaaat tattgacatc 540
 tgaatctgaa agctgcaaaa catgataaaa gacataataa aatcacagat ttgttattct 600
 ctcaggaact ttttctag 618

<210> 31
 <211> 655
 <212> DNA
 <213> Homo sapiens

<400> 31
 ggtgcccattg cttggtggta ggatgtatga agctccttgc tcctccagct gggcatcctg 60
 ccacttgctg agccaaataa ggaatgtggg gaagcagcag gccaccagcc agatgagggc 120
 cactgccttc caggcagccc catgggacat gaaggagagg tagcgagctg gatggatgac 180
 tgccaggtag gtgtgcagca caatggcggg gaaggacagg atgggtgctg tgcaggcggc 240
 gaagacagca tcagtggaaa tgccacaggc catgcggccc agctcccagc caccaggct 300
 gctggaggag atgagcatgt ggaggagaat gtaggccagg tctgagagca ggatgttagc 360
 cgggagcagg tagtggggct cctgtcgag cgggtggttc cgcaggatgg tcaccagcag 420
 cagggggctg acagccagtg tggctgcagc cagcaggctt gagggaaagga aaagccagta 480
 cagcatggag ctgggcaccc tgaggtcccc caggcccaag gaagtgttgc tggctgcttg 540
 gggcatgcag ggtgtcttgc tgatgagctg gatcagggcc ggccaagctg tagtgccac 600
 agggcaaggt gccagctcat ccccatgct tcctggcagg gatggctggc tttgt 655

<210> 32
 <211> 697
 <212> DNA
 <213> Homo sapiens

<400> 32
 aacactgact tctctgaagc agttgtctaa aagaacctac accattttta tttagcaaaa 60
 aggcttttgt taaaagcagg ggatagcaga aagagctttg taaaaaatat gtcattggatt 120
 ttaggagttt ctaagagcaa gaaaacgttt cttaaataga ggaatgaagc aattagagtt 180
 ccataaaaat cacctaattg gccttccaaa aggcaaatgc taaagcccca gaaatcatca 240
 ctgaggaagt ctgaagtagg aagagacctt gttctagaaa gccgacaagg tagaaattaa 300
 aatggaacag gcccaacttg aaattccgag accaaaagag gagctgatga cattggtggg 360
 agacagggtg gggaaataag aatgttggta gattctagag acattccagc gataacacag 420
 acaggacttt gtgactgact gtatggggca gctgcagggg taggagagga ggaacgatta 480
 agacatgatg aactgggcta tgagttggca gctccattta ctccagagaa cacaggaggt 540
 gaaaatcatg ggagacttga tgaaaacact ttgagaggca ccatggggat aaaagccaga 600
 aataaggtgg gaaatggtgg aagctattca ttctagaaaa gaggggtggga ggatgagcat 660

00196PCT1.ST25

aagttaacag gaaacaagtt aattttttaaa aagtgct

697

<210> 33
<211> 611
<212> DNA
<213> Homo sapiens

<400> 33
tttcccagat aaattgtatg cacagtaact ggtgttgacg tataccatag catatataca 60
tccatttggc aactgcagg tgccagtggg acaacatacc agagtgtgag tcttcctgat 120
cattttcatg atgtcctcag ttattttacct tgtaataagc ttgtaaacgt ctatgattgt 180
ttttgagtca tcccaatgca gtcagtgaat aacaacatgt attttaaatg aaacttgggg 240
attttcctcc atacctgaat ctctagtatt cacataaatg aaaaatcaaa attaggataa 300
gttagtgtca aacattaatg gattttttaca atgctaattg gtgttccttt ttaaattatt 360
gtgcctaca gacacatagc tatagtttcca tgcactttca accaccaatg ctgccaggct 420
agtaaagcag ttaatgtata tttgggggta attatcagaa tcaccagaaa caattttttt 480
aatttttaaa atattttatt tttccacaga ttattggggg acagatgctg tttgattaca 540
taagttcttt actggtgatt tgagagattt ggggtgcacc aacatccgag cagtatacat 600
tattccctat g 611

<210> 34
<211> 602
<212> DNA
<213> Homo sapiens

<400> 34
cgggttatit agagaaccac ttgaaatacc acctccttgg taacaccagc tccctccacc 60
ccctgagctc acggtctctt ccctgtgaga tgcagcacca ggtaagggtca ttaacaacca 120
ggtttagagt aaacagtgtc ggggtgtatt tctgatcctg cctttcccta actgggtgct 180
ctttggcaag ttattaagtt acttcatctg tacaatgggt tacacttatg ccttttacat 240
atgggtgttg cgaagattga gtgatatgca taccaaaaat gctgagcaga acaccttgct 300
catatccttc ctctctgtta ttaaatggag gcctttaagg ttaagtaatt tgttattgtt 360
gtgggttaatt ttagtcctct gaattttaat ctagtacaaa ttgtgctgca tttggcacat 420
ggtacatgtt catgaatatt gagtgttgta taaaggaatg aaaaatcaat tacatgaaaa 480
gaaattccaa atcttacatt ttacaaacac agacacaaaag aataactaaga ttttaactcag 540
gggcaaaaagt taagatttgg ccaccagcac gtggtgagct tccttgaaag tttgtttctg 600
gc 602

<210> 35
<211> 428
<212> DNA
<213> Homo sapiens

00196PCT1.ST25

<400> 35
 attctaattgg cattaatcca tagcattatc tccatctctg ttttaaataat catgcatcct 60
 cattctatga ttcaatttaa aggaatcctt caaaaggatc aatctaaata aataacaagt 120
 tagctttcag gcaaacaaat aaatttgctt tgttttatat tcaccataaa tatttcactt 180
 aattactgag gtaccttggt caggaaacac aaaacaacat tataaattaa ttagcactgt 240
 ccctgctgac gtttttagtcc tgtggaatgc aaaagctaaa agtaaaaaca ggccatgaag 300
 cccaaccaga gcacacatcg tatgcaaag ataaagccca caaacatcat gggatcattc 360
 ctgggacatt ctgaatcacc aaaattttgt cttaaatcaa gtattgccct atttattttc 420
 aaattcaa 428

<210> 36
 <211> 643
 <212> DNA
 <213> Homo sapiens

<400> 36
 tttgggtcta gaatccccctt ggtttgggaa gcatccagaa ggagcttcca tccccatcca 60
 tttcttctc acttctcct ctctagcttt gtttacatgt ctctcgatac ctagacagag 120
 gcagaggcat ggacttctgg ttctaactcat tcaagcctta cacgtccttc aaggctccat 180
 tcagaattat ctttctcgg ggagtctgct tctcctattt caggatttac tgactattct 240
 cttatctctt gtaacattta atatccact ccttagcatt aactttttaa cttgctttct 300
 aatcctgagg tttgtgttcc ttgccttgtt aaaattcttt gtaaatggcc agcccagtac 360
 gtagccagct cccaagcacc acgtaggcaa ttgaaggagc tgggacaaaa agagttcttt 420
 gtttgaattt cttttactgc tctgagttaa ctctgtattt gcacatgagt tttaatgttt 480
 tggggccatt gaactatttg agaactctaga agataatata cctcttttca gaaaaacaca 540
 tatgaatata cacacacaca tgccacctac acacacaatt ttgcatgtaa ttttaaggat 600
 tcattaacct tagcttacca gactgtaagt tcctttgcat att 643

<210> 37
 <211> 567
 <212> DNA
 <213> Homo sapiens

<400> 37
 gcaagttatc tgtatttata cccctacaaa cacacactcc taacatacag tggtagaga 60
 ggaacaacat aactgcagag gaagtaagt agagacaaa agcagtcatt ggttcattgc 120
 tataatgaaa ttctcttaga caaatgctgc caggatctct tccctgggga taaggcttag 180
 ttatcttctt ggaagtgggt tccagctcac tattctctac tgtataatta cagtactcc 240
 ctcatccatc ctcttgctt ctcagatctt aactttatcc tctagactcc aggtcctcc 300
 tctgagatgt tctcactttt ctgcaacaaa agctgagtct atttctcaat ctgtttgctg 360

00196PCT1.ST25

tccatagaaa atggaagggtt cagaggcttt tattcaattt tctcagtctc tttattgcaa 420
 gctgggtccc atttacttat ataactcttt taaaaagttt ttgtgggctt tctatgtatc 480
 agataataga ccacttcatt tgataaaaag ccacattctt tgttttccag acaagctttc 540
 tatatttttg acaagtaagg ccactta 567

<210> 38
 <211> 594
 <212> DNA
 <213> Homo sapiens

<400> 38
 ttatggtgtt tgtaagatct tattgccc aa agagtctgtt ctgtccatct tatgatctct 60
 gttttaacat taatgatgct cagttgtgtc tagaccctaa aagaagaagt ttgtatgact 120
 ttccatgctg ttatggtcag gaatttagtt ttaagctttt ttggggcctc taagccacaa 180
 ggggatctgt tcagtcagtt cagtagaggg cttaggattt atcatcttta attcacattc 240
 cccatttttg gtcaaaatat gccaaaagta gcatcaatag ccaagctctt atttcattcc 300
 atattattac cagggtggtg ggctatctat ctcagatata ttctgttctt caatgggacc 360
 catatagcca agggacttat agccaaaaga cttacagcca attaacatt ctaggacaaa 420
 agggaaatgga ggtgggaagg cattcattat tccttaaaaa ccttttgagc aatataagag 480
 ccacaaacca aaagccaaaa agtaagctta caaaaccgat ttatctataa gttctatgtg 540
 ttgggccatc ggctcttagg catctgtgag cccatctttt ttggaggatc tgaa 594

<210> 39
 <211> 282
 <212> DNA
 <213> Homo sapiens

<400> 39
 aaggcagagg gggccagcag ggcggtttac agaaccatga tgtgttttta actggactca 60
 cttctgccag tatctgctg actcttcagc ccatgtctct ttctctgtt gtaataactaa 120
 tgggggcatt aaggagccag agaagggggc tccgacgcca ctgcttgtag ctctggagtt 180
 acatttagcg gcatttatat tttgtcatgt gaaattcgaa atcctcatcc aaaatgcaac 240
 tgtgggggaa ctctcatagg aatttcagcc aattctggct cc 282

<210> 40
 <211> 626
 <212> DNA
 <213> Homo sapiens

<400> 40
 cccttctctc ccagccatac cgtgaccac ccataagctg gcccccttag ctctggctca 60
 cctggctcag acttagaggt ggcaggattc ctgctgctca ggaaataagg actgctcttg 120
 agctcctcac agggccccag aatcccaaca aaagccaacc aaggctacct tcaggccttc 180

00196PCT1.ST25

cagaaggggg tggtagtgtc ctcacaggt tcccaagtt tagggagagg gcagctgggc 240
 ccagggccct tctccttggt gctcaggatt tagcccccact taccatggtg cagccccagc 300
 cttccagcca acccagcatt agaggcagtg gctcctctta atgccaggcc ctagttggct 360
 caggcataat ccagccagga aacctctcac ctttccacag caatggccac cagtgtgaaa 420
 acggaagccg acacagacat gccctgcacc aagccgctca tcttgcatgt ggcattgtcg 480
 aagggccacc ctgcaatgac agaggccccc acagagttag agatgccac gcacaaagag 540
 ccagagactg aaagccctcc aagccaggtc ccctctgagc ttggatcttt cctccatgac 600
 ctgctagggtg ttatctggtc tctgct 626

<210> 41
 <211> 685
 <212> DNA
 <213> Homo sapiens

<400> 41
 aaactcccaa acgatagtaa cttaaataac ttaggtcttt aatactctct tcagtaaaaag 60
 aattctagta gttggagagt ccaccatccc taggaatgta gttcttgtcc tcatggttca 120
 atatagctgc tgatgctcca gccattacag ccacattcca gacagcaaaa tatggaaaga 180
 gaatgaagag aagaagagcg tgcctaggag tcccatgtat tatttccata tatatttggg 240
 cagaacctag tcacagggcc actccatacg tatctgttag ctattgctac atagcaacca 300
 caaaatttcc atgtcataca acacatatct gcaggttggc tagggttcag ttcctccatg 360
 ctgggtctcag acaggcagtt ctgcttcggg ttacagtggc tgagctgatt ccatttctca 420
 ctgcagggtct gtgtttcagt tgagtgactg tcccatgtgc ctttcatctt ccttgggttg 480
 atgaaaggaa gccacatctt tcaacagggc tagccacatc tggttcctcat ggcccaaaga 540
 gacaccaaag agcagataga aatagggtgag acctcttaag gtctagactc aaaactggca 600
 cactgccacg tctgttcaca agctattagc caaagcatag atgcattacc aagccccaag 660
 tcaagggcaa gaagtacaat ccacc 685

<210> 42
 <211> 566
 <212> DNA
 <213> Homo sapiens

<400> 42
 acaataatgt gttgtatatt ccaaaatagc tagtagtgta atgtttccaa tacaagaaa 60
 agataaatgt ttgtggtgat gcataattca agtaccctga tctgataatt gcacattgta 120
 tacatctatc aaaaatcag cagtacctcc aaaatatgct caattattgt ataagtacaa 180
 aaaaatttaa acaattataa tgtattattt atttctaaat gggttattag attttaaatt 240
 ttcttggtgt ttaatttttt catatattac cttataccct ttaacttcct aaaatatatt 300

00196PCT1.ST25

aggtcttcat atttttagagt aaaattctga aaatcctttg agtatctgat tttacaatct 360
 tttcttccac tgattttccc ttagcaatgg cctgtttaaa gtgttgatgat gatgttactg 420
 agaaaatgggc tggctacctg atgcacatag aagccaatac tatggcagtg gttttctaga 480
 aaagaaaagg ctttactgtg agtctactgg caaggagaca ggtggcaaca ctcaaatctg 540
 tctccctgaa ctgaggatgg tgggggt 566

<210> 43
 <211> 578
 <212> DNA
 <213> Homo sapiens

<400> 43
 cctcttactt gggccccgtt cactagtcct tcagccaaac tgcctcacat gctattccca 60
 gtatgaaaat cttgccattc cctttatctt tttctcttc tctcatttac agccctgtgc 120
 tagtttcttc attcccttca agttctggcc aaactttatt tacctcttga ctgaccactc 180
 catctaaaat agtactcatc actgtgtatc cctcaacac actttatagg tcatggccat 240
 cacctgataa tgtgttatgt attttttgggt ttacttgttg tgttagttca ttcttgcat 300
 gctgtaaaga aattcctgag actgggtaat ttataaagaa aagaggttta attgactcac 360
 agttctgcag gctgtatggg aagcatgttg ctggcatctg cttggcttct ggggaggact 420
 caggaaactt acaatcatgg ggaagggtgac gggggagcag gcacatctga catagcagga 480
 gcagcaagtg agcaaagggg gacgtgccac acacttctaa gtaaccagac ctcatgagaa 540
 ctcactatca tgagaacagt accaggggat ggtgctag 578

<210> 44
 <211> 684
 <212> DNA
 <213> Homo sapiens

<400> 44
 agtataacaa ttactgctt tacatctcta tattttgctt atctcaagta tccactttgt 60
 ctggtatagt gtgtcattc cacagttttt ggctgtcctg ggaacaacaa tctagtcaa 120
 ctccagcaat gtgagttata gtgcaaagt caaaccagag cagcatcacc atctagaggt 180
 caaatgata actgcaaact ttctcacctt tatgagcctt ccgtattctg tatacatagc 240
 agtttatgtg aatgtacaga aaataatgtt tgctattgtt ttctctccag ttgggtttcc 300
 agaaagagat catggcataa agcaggaacc acctgtattt acagatggca tagggaagca 360
 tacatgcag agccatatat cagcagcact acagcatgtt tcaaccaaag atgagcctcc 420
 cacatgtcag acaaaccacc tacattggga ccacagcagt gacagtgttt tttagcacat 480
 tcctgataat gaaatctatg ttgaactcaa catgaatggc ttttccttc tcttggcagt 540
 caacagccta caccattctg catttgactg tttagtttat tctccctct ggaaaggcat 600
 gactatggaa acagagtaga ggatattttg gggatttatg aaactattaa tataatttac 660

00196PCT1.ST25

tctcattgct gtgctttcta caaa

684

<210> 45
 <211> 693
 <212> DNA
 <213> Homo sapiens

<400> 45
 tcagcttatac tggtaaatag cttttcgctc tgttgcatatc cttgagcata tgcatacagct 60
 acaatgttta taggtagctg tatggtgttt gacacagcac atggcgtacc tttaaaacaa 120
 ttatagcact gggatttgga tctgaattta tgttgctctg tcaaagtttc ctctttgtaa 180
 catggttagcc ttttaaataat taggcagcta cctgcaacac tgggcattca gactaaccca 240
 tcaggcttat ggcattctgc tcttctcgtt cctctctgtg gtgttggtac atcatgttag 300
 gtttatgcag tagacgtaga taggaagcaa gccaatggc tacagggtat tgaaagtcaa 360
 ttgctgagaa tgataaaaga caaggatagc cttctctgca aagaagtgtc aagaagattc 420
 taaacgtata caaggatctc aagagaaaca gtcccagata gcaacactat tcagtcttag 480
 actatggctg atactatata cttctccagc tcctctgctc ctcagagcag aaaacagaag 540
 attttgaaat gagcaccacc ccagctcctg aatacaatgg tacctttcat ctatttctgg 600
 tgacttttat tttcttttgt tgctggatcc cctacataat tgtaagcata tcgcaggcaa 660
 gcacaatggt aaacagtggg tggacgcttc ctc 693

<210> 46
 <211> 677
 <212> DNA
 <213> Homo sapiens

<400> 46
 atagaacttg attatattgg tatttttatt tcaaattttc aatttttga atggcagaat 60
 gttgctattg aaaagtgtct taaagggtcac cactgtaacc ccttcattgt gcttgagacc 120
 tgctcagctc cttaaattta caggggacgg atctgagaaa ctgactcaa gttgtaacct 180
 cttgcttagt tttctttcta gggagataac cgtctctcca aacctgtcga aatctaaatt 240
 tattacctct tacctaatac ttgggtccct gtggacttca cttcactgtt tgtgctaata 300
 gccctttcat caccatcttg actttggatt ctagagcatc acctacttcc ccattttctg 360
 tgacccttac attcctcctg tcagtcacta tgtctgattt attgttctcc cctatctttt 420
 gccctttgca aatcctcaag ccttcattct ggttcagacc tttaaaaggc tgagttactg 480
 gagtatagtg ttacccaaag tgagttgttc cataaaaaat tagtaagttg gaaaaaaaaa 540
 caaaaaacaa aaaaataccc taccataaaa gttggtaaat gttcctgtaa aaagggttcc 600
 ttggccagggt acatgttaga atagctgggt aagtttcttt gcagaaagac ttctcctggc 660
 cttcatttgt gactgtg 677

00196PCT1.ST25

<210> 47
 <211> 729
 <212> DNA
 <213> Homo sapiens

<400> 47
 gcaattaagt tttgtactgt atggacagtg tgaaaaacat tatggaaaaa caacttgaaa 60
 gaaaatgtga cagaatttct cctaacaatg tcattgcttc aaccagctac aaatttccaa 120
 cctagtttct ttcttttgcg gtttcttctt ttgtctttga tacaatcata cagcctctct 180
 tccttgaaga gataataaaa gactaacagt taaaagatct ggaagactca tattcttttc 240
 tttttcactg gctacgggtt tgaaaagagg ctgttggtct ttgatttttt cttttggggt 300
 ctttacatcg cccaattcaa acaggtctgc tctcaaagaa aacaaatcaa aattgtcaag 360
 acctgtgaag catgaaaaat aaattgcttt ttccaactcc aaaaagcacc agaaaagcat 420
 taattttgat cttttataaa cctctatccc ctatcctcta atctatagat ttcacagaat 480
 gtttatatat tcttctgtat aatacaggag atcaaacctt attatgaata aattgaattg 540
 aacctgtaat acaactaata tttaaactag tgttattttg gagttcaact agacacatat 600
 aaaacatttc aagtgagatg acacaaattc ctggggctgc cagtataaaa taaacagtcc 660
 agtaagctgc atctaccatg ccgttaaggg actctgtcct ttagctggtt gggagcacag 720
 gcttcataa 729

<210> 48
 <211> 595
 <212> DNA
 <213> Homo sapiens

<400> 48
 tcctgagaag acctgcagca cagggtaaaa tatgcaaggg agggccatat aacttttatac 60
 ttacttaat ttattttaat ttactaattt ttaagtatta acctattttg tttttattaa 120
 atctctgtgg ttgcacagaa ttcaaattgc agcaaaaatc attcagggct aaacactgga 180
 aaaatctctt aattctaagg tacatgacac aatggactca aaaacagtgt ctgagtcctt 240
 ttactgtgag aaattttaag aaagggtata gaaaagtttt gaccaattcc acccaatcct 300
 gcatcccaa ttccaatctc aaggaccagt ttccatctga tctctctcca cctacagatg 360
 gtggtcctga atctccaaat caacaaacca aaaactgaat ccatcatctt ctcacacctg 420
 gtttttctt ccaactccct catttctgtg acctgcccc taaccttacc aggaatccag 480
 ccccaaagc aggttggtact cctccctctg caatggacac cagggtattca ggtcctgttg 540
 ctggtcctca aatgcccaca atgccctgtt ctcccaaatc agcacattca acagt 595

<210> 49
 <211> 710
 <212> DNA
 <213> Homo sapiens

00196PCT1.ST25

<400> 49
 ttgacttaa ccctttgtag cccaggtaat aaatccaaac tcagcaagta tgggctggac 60
 cccagtagct ctgtggttgc cacttttttg cccatattga accgacgtcc ccttggcac 120
 taccagggaac tcctcaggga gagtgtggga atgatggggg aagactcgtc actcttttgt 180
 agagcgtggg gcagatgata gcagagacct tccaggggcc agggctgggg tcttgtcttc 240
 cttggatgtg gtctagcgtt gctccagatg gtgggtttgt ggcagggtgg gcagaagcag 300
 atgatgcagt tgaggcgggt ctctggtaga gaggatgtc aaagatgagc actcctttta 360
 tcccctgact cttctgagga tggctgcctc cttggtgagc cacttgaggg tctcaggccg 420
 atcatgcggg atggtggccc agatgaggaa ggggatccaa ggcggtggcc tcccagatg 480
 cactgggccc cagcccttct tcctagcttc ggctgattac tgtgggcttc agcaaccagg 540
 gcctacctgt aggtctccac atttgtaagc accacagaac ccagtgcac tttgtgcaact 600
 caaatcgccc ttgatccagg ggtattttct cattcagaac acactttgaa aggcggccat 660
 tccccttctg gagaaagcct ggagaatcta cagtgcctt aattacagt 710

<210> 50
 <211> 550
 <212> DNA
 <213> Homo sapiens

<400> 50
 agatgccag acaccttcac ttcagcagac aaggggcaga gtcctggaaa atctaggcag 60
 ggaagacttg cgcctctaag agtaaaaggc ctcccagaga ggacatggat gaaaggagga 120
 ccaccttcca atgccactct ccaaagcagg aaacatccaa ataaaggatg ttgattttca 180
 ggaccccatc ccttcagtag tgcttacaca actggtatat cctctcccg tcttctctct 240
 ggtagccaag accttatacc agtttgagta tcctttatcc aaaatgcttg gggtcagaag 300
 tgttttgaat ttcagatatt tttaaatttt ggaatattta tatcatacct cttggttgaa 360
 ccttcagat acaaaaatct ggagtccagt gaggatttcc tttgagtgtc atgtcagtgc 420
 tcaaaaagtt ttagattttg gagcgtttca gatttcagg ttttgaaatt ggaataactca 480
 acctgtactc tctgtccttg ttctacctct accagaccct ccccccacagg aatgaattta 540
 gatctgaaaa 550

<210> 51
 <211> 747
 <212> DNA
 <213> Homo sapiens

<400> 51
 tcacccctcg ctgtctatct tgagctgtga gtttatccac aaaggaacag agctgaaatg 60
 aaacaatttc accacagtaa cttgttaatc gggcatcctt taagtatgct ggatttaaca 120
 ctggaagttc ttttgaagac tctgaaagtt ttctttaatc gtcagtgat ttttccaaac 180

00196PCT1.ST25

```

taagttcatg atatggattt ttttcaactgt atctagctta agtcacattt caattcaaatt 240
ctaaacctaa actgatggag ctggagctag tgacttcagg caattggcat cttttcgctg 300
aatacaaaaca tcctatttaa aagaccaaac acatgactcc attcaaaaat taaaacagtc 360
atgtgtagtg aaacagcaag aacacggtct gagaaacgtg tccttgacaca cacagcgtga 420
atgcactcac gcaagcctag acggtgcggc tgccgcacac caggccctgt ggtacagcct 480
gtcaattcca ggccccaagc ctgcatacca tgttgctgtg cgggacgctg ccggcggctg 540
tagcacaatg ctaagtatct gtgtatctca acacagaaga ggtagagtaa agtacagtat 600
tatgatcgta cgggacgctg gttgtacaca cagtctatca ttgatggaag catcggtata 660
tggcacatta ctgcactgta aaaagacacc aaacttcggc cggcgagtg gctcatgcct 720
gtaatccag cactttggga ggctgag 747

```

```

<210> 52
<211> 695
<212> DNA
<213> Homo sapiens

```

```

<400> 52
ttcttccttt tcctttcatt atcattttct tttgtctca aaataatgaa aaatgcataa 60
gggtctgtag agagaagaaa atgtccttgc ccatgaactt cttgcaggta tttatcttgc 120
ttctttatct tactaaaaat agaattgaaa gtttttcatt tttgttttt caattttaga 180
ggatacaatg gagattcagg aacgaataga aaatagtttt aagtctttac tagaccagt 240
aaaaggtaag ttttctact gtagattcct gtatttgtat ctggttgat ggcaatagct 300
tcgaagttct ttccctatt cccaagccca atcaccaga gataagtaag tagttttaac 360
actttggagt caatactcct agatgccacc taaacacata tgtgtgtgaa tgaaaataca 420
gataaaaagt aatctttaa cataggaaat ggtgtaatcc atgctttttt gactttaatt 480
ttttgttat tttggatacc ttccatgac agttatata accccattta tttcaagac 540
tgcgtaatat tctatagtat tgtattaaca tttttatgt tatcgcaatt ggtgacatat 600
tatgtatatg agttatttct tctactgatg ctgaaatgaa tatcttgga caaattgtta 660
ggggtattat ttgagtcctt cttgggatt aaatt 695

```

```

<210> 53
<211> 735
<212> DNA
<213> Homo sapiens

```

```

<400> 53
cttttgagga ttaaaaattc ctgcttactg tcgttataac acggggatta ataagcacct 60
tactggaatc tctcacctac cataatttta gtatgctatg tgagggaatg aacagtctca 120
cacatttaat aatgactact catataatgc ttttaattgg taatgaccta tatgaaacat 180

```

00196PCT1.ST25

gatatagaaa acacattaca gcttctcaaa tgacccctat aagttaacca attgcttagg 240
tttctgacaa atttgaatct ggcccatgc acctttgctg ggcccacaa aacaaggagg 300
tagattatit atgaaggcca accactctgg caatatcacc attaaatata aagctcatct 360
gcccatagc tcctccatct tcaggccag gactctggat tggaatgacc tacctccaca 420
ttcagttctg taagtcatta ggcatcatcc aagatggtag atgatgaata aatggacaat 480
gacttaagct ttttttactc tctcatccat tccaatgctt tcttcctgg tctttgctca 540
ttatttccat gttatttaata atatatitgg aagaattcat ggcatgtata acaataatgg 600
ctacaatttt ttattaccta tgtatgccag gcattgtgct aagtgttca ggtataagat 660
cttgaaggga attggttaca ttttacagat ggtaagactg ggattcagat gttagttgcc 720
tgtttaagtc aataa 735

<210> 54
<211> 427
<212> DNA
<213> Homo sapiens

<400> 54
ctcttctccc taggtggttt gctggcaatc tttggcattc cttagcttgt ggaagtatca 60
ctccatctct gtcctgattt ctacatggtg ttcttcctgt gtgcatgtct gtctccaaat 120
ttccccattt tataaggaca cagtcatact ggattcgggc tcattctaaa gacctcattt 180
aatttaattc cataaagacc ctatctccaa ataatgtcac attctgtggt actggggggt 240
atgacttaaa catataaatt ttagggagac aaatttgaac ctctaacagt actgaacatc 300
caggatggaa gaacatggta ttaggttgag ccaaacacag ttgcttacgt tttggttttc 360
ctcaccagga caagaaaccc ccagtgcagg aaaattggag acatggaaaa cagggcttaa 420
gtaaaca 427

<210> 55
<211> 713
<212> DNA
<213> Homo sapiens

<400> 55
ttattatact caacactgct aggaagaat cagtgtgtt gaagatatat atatatatat 60
ttgcttgtgt atttgtgtgt gagagacaca catagaaaaa aagagagaga gaaatatatt 120
ggttgacact ggcttctttg aaaaaaggca gtttagtaac aatggccttt actagacaga 180
catgttagaa ggcagcagga gaaagggaat gtggtatcag atattttctg taaaagggtt 240
gttattaata ttcattgtggc aaattgtagc tgatgtcaaa gtagttataa agcaaggga 300
acacaattct ttacagcaa tgttgaggtc taagaacat aaaacaaata cctggtaagt 360
accatgcata tatacatata taaacaatca ataactcaca aaacattcac atatttgcaa 420
cactgctttt cagtttatgc agtttatttt ttgttctttt taagcttttt attatagtga 480

00196PCT1.ST25

atgtcttata tttcattaaa agttttgata ttatatgtga aacaacagtt ctgataaagc 540
aatatctaga taaaggctat tacttacctt tctcaaattg atagattttc tccttgtaac 600
aagctctgat ataaaatatg ataatttggt gaaaactttt acacattcaa aactaaatta 660
tcatatatattt aatgagactt tgggtgtgta tgtgtgagtg tgtgtctgtg tgt 713

<210> 56
<211> 607
<212> DNA
<213> Homo sapiens

<400> 56
aaaacttggt tttttaaagc aaacacagaa acaatgtaat ataggtctta ttacatatgt 60
aggaaataaa aataatatgt atgacgacaa cagtagtcta aaattcagga gacagagaat 120
ggaagtacat tgttgcaagg ttttctaata cacatgtaca aagtggata atgttacttg 180
aaagataact gtgataagtt aaagacgtaa tcaatgacac tatatcaacc actaaaataa 240
tacaacaaag gatatacgaa atatttttaa aagtataatt aaccctaaag aaagcataga 300
ggaaaaaggg aacaaagaat aatagatgga ataaacagaa aaaactagcc agctggtaaa 360
tttaaaaccg atcatataca tattcacatt aaatacaaaa agtttaaaaca cttcaaagtc 420
aagtcagagg tgtcatattg gataaaaaga aagactcaac tatatgttac ctataaggaa 480
tgcactttaa atatacaaac atattaaaat aaaaagatga aaagttatat acaatgttaa 540
tactcatcaa aataaagcta atgaggctat attcatatta aaaagtaggt tttaaagcaa 600
agattac 607

<210> 57
<211> 746
<212> DNA
<213> Homo sapiens

<400> 57
tcaagtccat gcttttacgg aaagacccca gttcctgcct cttctatata tttatctacc 60
ttgtggtgaa gagcatgtgt gtgcaacacc ttgcctgaa atggtatggt ttggcattaa 120
tgaattgtgg gtccattgaa aagaaatctc ctcttgtttc tcgtgttatg gacagttcaa 180
ggtttgctt agaactaact tcaaggaaaa gtagcagaat cgtaggaagg gacaatcttg 240
ccttcagtcc caccctctgt tccgggcagg tctgggtggc tatcttcttt cgggggcttt 300
tccttgcaaga agaacttctt cagcatgtcc tggatttctt tcttaatggt cttgtgcatg 360
tagccataga cataggggtg gatgcagcac tgcaggaaga aaagccagat gattatggtg 420
atcacccact ggggtacctg ggtttcgaca tccaccaca cggccaggac tgctaaaaag 480
cagtagggcc ccagggatag cacataggag aaaatgatga tgaagatcac tttagcagct 540
ttgcactggt agcacctggg cagaggaggg ttgctgttgc tgttacgacg actgggtggg 600

00196PCT1.ST25

aggctctccg ggatgttcac tgcctcgacg tcatcctcac tgaaattgat gtcgtcttca 660
 ccaaaactcca tgtcatcttc acccaagtca atgctgcact ggttgacctc tgtgcgaccc 720
 ttgtctgcct tcatgctgtt ctccctc 746

<210> 58
 <211> 638
 <212> DNA
 <213> Homo sapiens

<400> 58
 agtggaaaga ccacacctag gaaccgactc tagctcttac caccctgtaa gcctgaggct 60
 cagttgctgt ccctggagaa cagaaaacat aatcatggct attctgaggg tcaggggcaa 120
 gtgctttgca agtgggattg tgggtggcag tgggagggat tctgggggttc actgtcatgc 180
 tagttgtgta actgggcaat gcaaccgtgt aagtgtcagg aaaccctcaa taagactgag 240
 ccagaggcca ataagaagcc agcatttaca tgatgttctt ttcctttttg taactaggaa 300
 atttcgattt gcacactgat ttggcccacc attcctggag agatctcgtg ggatgtctct 360
 tttgttactt tgaacttctt ggtgccagga ctggtcattg tgatcagtta ctccaaaatt 420
 ttacaggtat gttttctgca agtgctgccca ctgaacttca cccaggcttg gggttatttc 480
 tgctagaatc ttagaatttg gggtcggaga acacctaaga gttcacgcca gctcaatctt 540
 gattcactgc ccaggctctac aacactgagg aaggagagga tttttttaga agttatatct 600
 ttgtgattat gttttttgct catcactaaa gtaatact 638

<210> 59
 <211> 216
 <212> PRT
 <213> Homo sapiens

<400> 59
 Asp Ala Phe Leu Phe Pro Cys Pro Glu His Gly Ser Val Met Thr Ser
 1 5 10 15
 Gly Ser Cys Lys Glu Ala Gly Leu Arg Phe Phe Gln Ala Trp Gly Glu
 20 25 30
 Val Gly Glu Glu Cys Val Leu Met Arg Arg Ala Gly Cys Ala Gly Ala
 35 40 45
 Glu Ser Ser Thr Ser Leu Gly Ser Arg Cys Pro Thr Ser Pro Ser Leu
 50 55 60
 Gln Pro Ala Leu Pro Lys Gly Ala Arg Ala Trp Pro Pro Leu Asp Met
 65 70 75 80
 Ala Ser Gln Pro Phe Gly Lys Cys Gly Arg Pro Cys Cys Arg Ala Pro
 85 90 95
 Val Thr Val Ser Val Trp Val Trp His Gly Trp Cys Ser Pro Ala Gln
 100 105 110
 Asn Pro Ala Cys Asn Ser Thr Gln Ser His Ile Pro Gly Gly Gln Ala

00196PCT1.ST25

115 120 125
 Leu Leu Leu Cys Ser Gln Met Pro Pro Ala Gln Lys Glu Asp Thr Pro
 130 135 140
 Ser Ser Ser Ala Glu Ala Ser Leu Thr Glu Gly Gly Cys Val Lys Ala
 145 150 155 160
 Ser Glu Ala Glu Leu Pro Ala Ala His His Gln Asp Ala Leu Glu Ala
 165 170 175
 Arg Ser Trp Ile Gly Ser Gly Cys Thr Glu Pro Ser Leu Pro Arg Asn
 180 185 190
 Thr Gly Asn Ala Lys Cys Ala Gly Gln Ala Val Gly Glu Gly Gly Met
 195 200 205
 Ser Leu His Val Cys Ala His Cys
 210 215

<210> 60
 <211> 204
 <212> PRT
 <213> Homo sapiens

<400> 60

Leu Glu Lys Gly Thr Lys Ser Gly Ser Val Phe Ser Ala Phe Phe Phe
 1 5 10 15
 Phe Phe Gln Ile Leu Val Val Ile Ile Gln Leu Phe Phe Leu Cys Met
 20 25 30
 Asp Phe Val Val Leu Arg Ala Ile Tyr Arg Ser Arg Val Gln Leu Leu
 35 40 45
 Lys Val Ile Tyr Ser Gln Phe Cys Ile Lys Pro Ile Ile Tyr Lys Cys
 50 55 60
 Ile Ser Ile Gln Tyr Arg Pro Gln Arg His Lys Ile Phe Phe Ser Leu
 65 70 75 80
 Leu Ser Cys Cys Pro Thr Asn Val Cys Arg Ile Tyr Gln Asn Ser Ile
 85 90 95
 Arg Lys Leu Leu Val Tyr Ala Leu Leu Ala Val Leu Leu Leu Ala Phe
 100 105 110
 Leu Phe Arg Val Val Glu Ile His Ser Phe Ile Asp Ile Lys Gly Thr
 115 120 125
 Val Lys Met Ser Leu Pro Val Asn Ile Asn Arg Leu Val Ile Leu Gly
 130 135 140
 Leu Gln Leu Asp Leu Leu Ile Cys Cys Ser Cys His Met Ser Thr Asn
 145 150 155 160
 Leu Ile Cys Ser Pro Phe Gln Lys Leu Asn Tyr Leu His Phe Phe Gly
 165 170 175
 Gly Ala Leu Val Trp Lys Val Arg Glu Ile Phe Thr Phe Thr Leu Phe
 180 185 190
 Phe His Phe Phe Leu Lys Thr Ser Ile Pro Pro Leu
 195 200

00196PCT1.ST25

<210> 61
 <211> 96
 <212> PRT
 <213> Homo sapiens

<400> 61

Val Glu His Ser Trp Pro Cys Ile Gln Tyr Ile Ser Trp Val Arg Pro
 1 5 10 15
 Gly Val Pro Val Ser Ile Ser Val Asp Leu Leu Ser Met Leu Pro Val
 20 25 30
 Ser Thr Trp Val Val Pro Trp Gln Glu Arg Cys Ile Cys Val Leu Thr
 35 40 45
 Glu Val Pro Tyr Arg Cys His Phe His Cys Gly Ser Ser Asp Pro Gly
 50 55 60
 Lys Asp Ser Phe Gln Gly Pro Gln Val Gly Ser Gly Gly Gly Gly Ser
 65 70 75 80
 Gln Thr Pro Asp Pro Val Thr Pro Ser Arg Pro Val Leu Glu Gly Pro
 85 90 95

<210> 62
 <211> 213
 <212> PRT
 <213> Homo sapiens

<400> 62

His Gln Ile Leu Leu Cys Cys Leu Arg Leu Gln His Ile Ser Met Ala
 1 5 10 15
 Ser Ser Leu Gly Met Val Thr Val Ala Glu Leu Gly Gly Phe Val Leu
 20 25 30
 Pro Ile Ile Ile Ile Thr Tyr Phe Thr Trp Lys Thr Arg Lys Ser Leu
 35 40 45
 Trp Glu Phe Gln Val Pro Pro Arg Asn Thr Lys Glu Arg Lys Lys Ala
 50 55 60
 Leu Arg Met Val Leu Met Cys Glu Val Val Phe Ile Val Cys Phe Thr
 65 70 75 80
 Pro Tyr His Leu Asn Phe Pro Phe Phe Met Met Val Lys Glu His Val
 85 90 95
 Phe Leu Asn Cys Ser Phe Ile Lys Ile Ile Leu Cys Phe His Ile Ile
 100 105 110
 Ser Leu Cys Leu Ala Asn Leu Asn Cys Cys Leu Asp Pro Val Val Tyr
 115 120 125
 Tyr Phe Met Thr Ser Lys Phe His Asp Gln Phe Ser Asp His Gly Ser
 130 135 140
 Leu Val Leu Gln Ser Cys Met Arg Cys Asn Asn Ser Thr Leu Glu Ile
 145 150 155 160
 His Gln Arg Lys Gly Gly Ser Ser Asn Tyr Leu Ser Met Phe Glu Arg
 165 170 175

00196PCT1.ST25

Phe Gln Asp Asn Ile Ile Lys Leu Thr Arg Lys Ile Asp Met Leu Tyr
 180 185 190

Cys Ile Tyr Val Thr Leu Lys Ile Phe Leu Phe Phe Phe Ser Phe Phe
 195 200 205

Leu Leu Tyr Phe Lys
 210

<210> 63
 <211> 197
 <212> PRT
 <213> Homo sapiens

<400> 63

Cys Tyr Cys Ser Cys Ile Leu Leu Ser Val Cys Leu Leu Cys Pro Lys
 1 5 10 15

His Arg Leu Phe Gln Lys His Phe Leu Leu Ser Pro Phe Ser Leu Ala
 20 25 30

Glu Ser His Phe Ser Val Ser Ser His Ile Ser Tyr Leu Phe Leu Leu
 35 40 45

Lys Thr Arg His Phe Arg Cys Val Val Ala Val Gln Ile Leu Ile Leu
 50 55 60

Ser Pro Arg Ser Cys Cys Leu Ser Tyr Leu Tyr Met Cys Leu Val Thr
 65 70 75 80

Trp Leu Asp Tyr Phe Asn Asn Val Tyr Phe Pro Val Val Tyr Thr Ile
 85 90 95

Phe Tyr Thr Asn Val Thr Phe Pro Ile Val Gln Pro Trp Ala Trp Thr
 100 105 110

Glu Leu Ser Trp Asp Asp Ser Asn Phe Gly Ser Leu Leu Ser Leu Ser
 115 120 125

Leu Met Ser Leu Leu Ser Tyr Leu His Leu Leu Val Ser His Leu Ala
 130 135 140

Phe Asp Phe His Leu Phe Asp His Cys Leu Thr Val Phe Gly Ser Ala
 145 150 155 160

Leu Arg His Lys Val Phe His Ser Leu Ile Leu Asn Ser Asp Ser Tyr
 165 170 175

Lys Ser Gly Leu Gly Gln Ser Leu Arg Phe Val Leu Thr Leu Gly Gly
 180 185 190

Leu Lys Cys Phe Pro
 195

<210> 64
 <211> 132
 <212> PRT
 <213> Homo sapiens

<400> 64

Pro His Ile Pro Phe Pro Ser Asn Pro Gly Asn Pro Lys Leu Phe Leu
 1 5 10 15

00196PCT1.ST25

Thr Ala Ser Phe Gly Ile Ser Ser Phe Trp Cys Gln Ile Ser Gln Gln
 20 25 30
 Asn Phe Leu Pro Ile Ile Tyr Gln Cys Leu Ser Val Lys Phe Arg Phe
 35 40 45
 Asn Phe Leu Leu Pro Arg Ala His Tyr Leu Ala Pro Ile Ile Pro Ser
 50 55 60
 Pro Asn Ser Gln Thr His Lys His Ser Leu Leu Gln Leu Trp Ala Ser
 65 70 75 80
 Tyr Leu Ser Pro Ser Gly Lys Lys Cys Cys Val Thr Pro Leu Ala Val
 85 90 95
 Ser Val Asp Leu Val Gln Gly Arg Ala Pro Val Arg Ala Ala Gly Pro
 100 105 110
 Ser Ser Leu Pro Gly His Gln Gln Ile Ser Thr Ala His Arg Cys Pro
 115 120 125
 Gly Asn Gly Ser
 130

<210> 65
 <211> 202
 <212> PRT
 <213> Homo sapiens

<400> 65

Ile Thr Ile Phe Gln Pro Leu Leu Leu Gln Gly Leu Leu Cys Thr Leu
 1 5 10 15
 Ser Leu Asn Ser Pro Ser Ile Cys Ser His Asn Pro His Asp Pro Gln
 20 25 30
 Phe Tyr Asn Thr Thr Val Arg Ser Pro Lys Leu Pro Phe Ile His Phe
 35 40 45
 His Ile Thr Ile Phe Gln Pro Leu Leu Leu Gln Gly Leu Leu Cys Thr
 50 55 60
 Leu Ser Leu Asn Ser His Asp Ser Ser Cys Thr Leu Gly Ser Ser Val
 65 70 75 80
 Ser Pro Leu Leu Leu Ile Ser Arg Val Pro Phe Cys Phe Cys Trp Leu
 85 90 95
 Pro Tyr Lys Ala Cys Asn Ile Ile Ser His Phe Arg Lys Glu Leu Asp
 100 105 110
 His Leu Leu Met Asn Pro Ala Phe Met Thr His Cys Leu Thr Cys Leu
 115 120 125
 Trp Leu Cys Met Ser Pro Ser Phe Arg Phe Phe Leu Trp Lys Glu Arg
 130 135 140
 Leu Pro Lys Ser Pro Ala His Gln His Tyr Lys Cys Met Gln Thr Ser
 145 150 155 160
 Phe Ser Cys Leu Pro Thr Leu Lys Met Ser Lys Gln Phe Ser Lys Gly
 165 170 175

00196PCT1.ST25

Glu Lys Ile Ser Ser Pro Pro His Thr Asn Tyr Leu His Asn Ser Val
 180 185 190

Thr Phe Tyr Lys Pro Cys His Cys Ile Ser
 195 200

<210> 66
 <211> 221
 <212> PRT
 <213> Homo sapiens

<400> 66

Thr Val Leu Ile Met Ile Val Phe Val Ile Cys Cys Trp Gly Pro Tyr
 1 5 10 15

Cys Phe Leu Val Leu Leu Ala Ala Ala Arg Gln Ala Gln Thr Met Gln
 20 25 30

Ala Pro Ser Leu Leu Ser Val Val Ala Val Trp Leu Thr Trp Ala Asn
 35 40 45

Gly Ala Ile Asn Pro Val Ile Tyr Ala Ile Arg Asn Pro Asn Ile Ser
 50 55 60

Met Leu Leu Gly Arg Asn Arg Glu Glu Gly Tyr Arg Thr Arg Asn Val
 65 70 75 80

Asp Ala Phe Leu Pro Ser Gln Gly Pro Gly Leu Gln Ala Arg Ser Arg
 85 90 95

Ser Arg Leu Arg Asn Arg Tyr Ala Asn Arg Leu Gly Ala Cys Asn Arg
 100 105 110

Met Ser Ser Ser Asn Pro Ala Ser Gly Val Ala Gly Asp Val Ala Met
 115 120 125

Trp Ala Arg Lys Asn Pro Val Val Leu Phe Cys Arg Glu Gly Pro Pro
 130 135 140

Glu Pro Val Thr Ala Val Thr Lys Gln Pro Lys Ser Glu Ala Gly Asp
 145 150 155 160

Thr Ser Leu Asp Gly Trp Asn Gly Gln Leu Met Lys Ala Asn Phe His
 165 170 175

Ser His Tyr Leu Met Met Glu Asp Ser Gly Gly Glu Leu Trp Ile Ser
 180 185 190

Ser Gln Thr Phe Lys Ala Arg Asp Gly Gly Gly Leu Pro Leu Ser Pro
 195 200 205

Asn Asn Ile Lys Asp Asn Val Pro Ser Phe Lys Lys Cys
 210 215 220

<210> 67
 <211> 595
 <212> PRT
 <213> Homo sapiens

<400> 67

Leu Glu Pro Thr Ser Lys Ala Pro Pro Gly Pro Gln Arg Pro Pro Pro
 1 5 10 15

00196PCT1.ST25

Leu Arg Pro Ser Pro Ala Pro Arg Gly Gly Arg Pro Pro Ala Pro Ser
 20 25 30
 His His Ser Asp Leu Ala Ala Ala Ala Pro Gly Ala Gly Gly Asp Pro
 35 40 45
 Arg Pro Pro Leu Gly Pro Met Glu Glu Pro Gln Pro Pro Arg Pro Pro
 50 55 60
 Ala Ser Met Ala Leu Leu Gly Ser Gln His Ser Gly Ala Pro Ser Ala
 65 70 75 80
 Ala Gly Pro Pro Gly Gly Thr Ser Ser Ala Ala Thr Ala Ala Val Leu
 85 90 95
 Ser Phe Ser Thr Val Ala Thr Ala Ala Leu Gly Asn Leu Ser Asp Ala
 100 105 110
 Ser Gly Gly Gly Thr Ala Ala Ala Pro Gly Gly Gly Gly Leu Gly Gly
 115 120 125
 Ser Gly Ala Ala Arg Glu Ala Gly Ala Ala Val Arg Arg Pro Leu Ala
 130 135 140
 Thr Glu Ala Ala Pro Leu Leu Ser His Gly Ala Ala Val Ala Ala Gln
 145 150 155 160
 Ala Leu Val Leu Leu Leu Ile Phe Leu Leu Ser Ser Leu Gly Asn Cys
 165 170 175
 Ala Val Met Gly Val Ile Val Lys His Arg Gln Leu Arg Thr Val Thr
 180 185 190
 Asn Ala Phe Ile Leu Ser Leu Ser Leu Ser Asp Leu Leu Thr Ala Leu
 195 200 205
 Leu Cys Leu Pro Ala Ala Phe Leu Asp Leu Phe Thr Pro Pro Gly Gly
 210 215 220
 Ser Ala Pro Ala Ala Ala Ala Gly Pro Trp Arg Gly Phe Cys Ala Ala
 225 230 235 240
 Ser Arg Phe Phe Ser Ser Cys Gly Ile Val Ser Thr Leu Ser Val Ala
 245 250 255
 Leu Ile Ser Leu Asp Arg Tyr Cys Ala Ile Val Arg Pro Pro Arg Glu
 260 265 270
 Lys Ile Gly Arg Arg Arg Ala Leu Gln Leu Leu Ala Gly Ala Trp Leu
 275 280 285
 Thr Ala Leu Gly Phe Ser Leu Pro Trp Glu Leu Leu Gly Ala Pro Arg
 290 295 300
 Glu Leu Ala Ala Ala Gln Ser Phe His Gly Cys Leu Tyr Arg Thr Ser
 305 310 315 320
 Pro Asp Pro Ala Gln Leu Gly Ala Ala Phe Ser Val Gly Leu Val Val
 325 330 335
 Ala Cys Tyr Leu Leu Pro Phe Leu Leu Met Cys Phe Cys His Tyr His
 340 345 350
 Ile Cys Lys Thr Val Arg Leu Ser Asp Val Arg Val Arg Pro Val Asn
 355 360 365

00196PCT1.ST25

Thr Tyr Ala Arg Val Leu Arg Phe Phe Ser Glu Val Arg Thr Ala Thr
370 375 380

Thr Val Leu Ile Met Ile Val Phe Val Ile Cys Cys Trp Gly Pro Tyr
385 390 395 400

Cys Phe Leu Val Leu Leu Ala Ala Ala Arg Gln Ala Gln Thr Met Gln
405 410 415

Ala Pro Ser Leu Leu Ser Val Val Ala Val Trp Leu Thr Trp Ala Asn
420 425 430

Gly Ala Ile Asn Pro Val Ile Tyr Ala Ile Arg Asn Pro Asn Ile Ser
435 440 445

Met Leu Leu Gly Arg Asn Arg Glu Glu Gly Tyr Arg Thr Arg Asn Val
450 455 460

Asp Ala Phe Leu Pro Ser Gln Gly Pro Gly Leu Gln Ala Arg Ser Arg
465 470 475 480

Ser Arg Leu Arg Asn Arg Tyr Ala Asn Arg Leu Gly Ala Cys Asn Arg
485 490 495

Met Ser Ser Ser Asn Pro Ala Ser Gly Val Ala Gly Asp Val Ala Met
500 505 510

Trp Ala Arg Lys Asn Pro Val Val Leu Phe Cys Arg Glu Gly Pro Pro
515 520 525

Glu Pro Val Thr Ala Val Thr Lys Gln Pro Lys Ser Glu Ala Gly Asp
530 535 540

Thr Ser Leu Asp Gly Trp Asn Gly Gln Leu Met Lys Ala Asn Phe His
545 550 555 560

Ser His Tyr Leu Met Met Glu Asp Ser Gly Gly Glu Leu Trp Ile Ser
565 570 575

Ser Gln Thr Phe Lys Ala Arg Asp Gly Gly Gly Leu Pro Leu Ser Pro
580 585 590

Asn Asn Ile
595

<210> 68
<211> 201
<212> PRT
<213> Homo sapiens

<400> 68

Ala Ser Ala Ser Gln Ala Gln Phe Lys Lys Lys Met Phe Asn Leu Leu
1 5 10 15

Leu Thr Tyr Phe Cys Lys Ile Leu Lys Ile Tyr Thr Ile Tyr Trp Leu
20 25 30

His Asn Ile Val Lys Ala Leu Thr Ala Thr Lys Leu Tyr Ala Gln Lys
35 40 45

Trp Leu Lys Trp Tyr Ile Tyr Ile Thr Tyr Ile Leu Leu Gln Phe Val
50 55 60

00196PCT1.ST25

Ile Glu Lys Asn Glu Met Lys Lys Val Lys Phe Gln Pro Gln Leu Cys
 65 70 75 80
 Phe Asn Asn Ile Gln Asp Leu Val Lys Leu Leu Lys Phe Leu Asn Ala
 85 90 95
 Tyr Phe Gln Phe Leu Tyr Leu Ser Arg Cys Arg Pro Val Lys Val Cys
 100 105 110
 Met Leu Ala Ala Ile Pro Glu Leu Tyr Phe Asp Ser Thr Asp Leu Ser
 115 120 125
 Cys Glu Gly Leu Trp Leu Cys Arg Ala Ser Gln Glu Thr Phe Glu His
 130 135 140
 Lys Val Ser Cys Thr Thr Thr Pro Ser Ser Arg His Phe Trp Thr Pro
 145 150 155 160
 Gly Trp Ser Thr Pro Ser Ser Ser Gly Gln Ala His Cys Ser Asp Val
 165 170 175
 Trp Leu Thr Pro Thr Tyr Ala Pro Ala Val Pro Gln Gly Pro Cys Cys
 180 185 190
 Thr Val Val Phe Ile Tyr Phe Leu Arg
 195 200

<210> 69
 <211> 217
 <212> PRT
 <213> Homo sapiens

<400> 69

Arg Leu Lys His Ile Leu Pro Ser Ser Leu Arg Leu Ala Ser Lys Asn
 1 5 10 15
 Ala Phe Asn Trp Leu Asn Leu Arg Ile Ile Val Tyr Cys Cys Leu Gly
 20 25 30
 Ile Ile Glu Cys Cys Leu Leu Ile Lys Val Glu Phe Asp Pro Pro Arg
 35 40 45
 Leu Pro Leu Val Trp Val Gly Glu Gly Leu Gly Phe Cys Ser Phe Phe
 50 55 60
 Phe Leu Leu Leu Ile Arg Ser Thr Asn Ile Tyr Cys Met Pro Met Gly
 65 70 75 80
 Gly Lys His Arg Phe Cys Gly Ala Ser Leu Tyr Tyr Leu Gly Asp Pro
 85 90 95
 Leu Ile Lys Leu Ile Lys Leu Gln Ile Gln Asn Ala Lys Leu Phe Leu
 100 105 110
 Arg Met Gln Ile Glu Gly Thr Leu Gln Leu Lys Asp Tyr Ser Leu Tyr
 115 120 125
 Asn Lys Tyr Ala Ser Gly Ala Tyr Cys Met Ser Gly Thr Leu Gly Pro
 130 135 140
 Val Asp Lys Val Met Asn Ala Ile Val Thr Leu Thr Trp Ile Leu Gln
 145 150 155 160
 Ser Ser His Phe Gln Lys Met Val Ser Leu Phe Val Pro Pro Gln Arg

00196PCT1.ST25

165 170 175
 Ala Thr Trp Tyr Thr Ala Leu Leu Val Ala Glu Gly Pro Ser Thr Pro
 180 185 190
 Ala Leu Phe Pro Val Ser Ser Leu Leu Trp Thr Arg Lys Asn Pro Asp
 195 200 205
 Leu Thr Tyr Thr Gly Gln Ser Ala Leu
 210 215

<210> 70
 <211> 156
 <212> PRT
 <213> Homo sapiens

<400> 70

Glu Gly Leu Ile Thr Ala Gly Val Ser Asp Ala Pro Leu Pro Gln Met
 1 5 10 15
 Gln Ile Pro Gly Thr Leu Trp Met Tyr Tyr Tyr Leu Ile Ser Leu Tyr
 20 25 30
 Ile Pro Phe Ser Met Asn Gly Glu Pro Met Pro Phe Trp Arg Gly Glu
 35 40 45
 Arg Trp Ser Ser Ser Ala Ile Leu Pro Lys Leu Phe Ala Phe Leu Glu
 50 55 60
 Asp Phe Pro Phe Phe Ser Glu Leu Asp Ile Trp Met Ser Glu Ala Gly
 65 70 75 80
 Arg Gly Phe Cys Phe Leu Ile Ala Ala Leu Arg His Thr Ser Pro Ile
 85 90 95
 Pro Ala Gln Met Arg Arg Pro Leu Glu Asn Lys His Gln Phe Arg Phe
 100 105 110
 Leu Asn Ile Ile Pro Thr Leu Ala Ile Met Pro Ala Leu Glu Thr Lys
 115 120 125
 Glu Leu Cys Ser Arg Lys Val Ser Val Gln Gly Tyr Thr Val Phe Ala
 130 135 140
 Val Cys Arg Gly Lys Phe Leu Thr Asp Ile Cys Leu
 145 150 155

<210> 71
 <211> 221
 <212> PRT
 <213> Homo sapiens

<400> 71

Leu Ala Gln Val Phe Pro Val Pro Gln Gly Asn Glu Tyr Phe Lys Gln
 1 5 10 15
 Lys Arg Val Arg Asp Met Ser Asn Val Tyr Cys His Thr Leu Thr Leu
 20 25 30
 Trp Ala Leu Ile Phe Leu Val Val Leu Ser Tyr Tyr Tyr Arg Phe Leu
 35 40 45
 Pro Cys Ser Tyr Leu Phe Gly Asn Gly Thr Glu Ile Trp Leu Leu Leu

. 00196PCT1.ST25

50	55	60
Gly Thr Asn His Ser Ser Pro Leu Trp Lys Trp Leu Leu Ser His Lys		
65	70	75 80
Tyr Ser Pro Ser Cys Ser Arg Leu Leu Ile Leu Asn Leu Trp Val Asn		
	85	90 95
Lys Val Thr His Leu Tyr Lys Glu Ile Gly Asp Gln Ser Asn Ser Pro		
	100	105 110
Ile Arg Lys Pro Gln Arg Val Gly Thr Asn Ser Val Met His Leu Glu		
	115	120 125
Leu Glu His Thr Cys Ser Asn Leu Gln Ser Gly Lys Leu Ile Val Leu		
	130	135 140
Trp Trp Leu Lys Lys Gln Arg Gly Thr Gly Ser Ala Glu Lys Pro Met		
	145	150 155 160
Asn Lys Pro Pro Val Pro Tyr Gly Phe Phe Leu Lys Ser Glu Phe Arg		
	165	170 175
Ala Gln Asn Glu Ser Ile Tyr Leu Val Leu Thr His Ser Ile Lys Asn		
	180	185 190
Glu Glu Thr Gly Ala Glu Leu Leu Lys Asn Ile Pro Val Ser Cys Lys		
	195	200 205
Ala Arg Thr Gly His Pro Tyr Val Leu Thr Leu Pro Cys		
	210	215 220

<210> 72
 <211> 237
 <212> PRT
 <213> Homo sapiens

<400> 72

Leu Pro Ser Gln Gly Glu Gly Arg Ala Pro Lys Gly Leu Met Arg Gly	
1	5 10 15
Leu Thr Asp Gln Gly Arg Glu Gln Asn Thr Phe Leu Ser Ile Gly Asp	
	20 25 30
Ser Val Thr Trp Leu Ser Leu Ile Ile Ser Glu Ala Trp Arg Ile His	
	35 40 45
Leu Phe Val Ser Pro Gly Arg Glu Asn Lys Leu Trp Thr Phe Ser	
	50 55 60
Ser Leu Tyr Asp Asn Ser Leu Tyr Val Asp Cys Lys Gly Gly Thr Lys	
65	70 75 80
Pro Ser Leu Leu Ser Asn Thr Ile Trp Gln Ser Pro Trp Val Ile Ile	
	85 90 95
Leu Asn Ile Asp Ala Tyr Cys Ser Arg Val Lys Lys Ile Ser Met Thr	
	100 105 110
Ala Phe Gln Phe Tyr Lys Phe Asn Leu Tyr Ser Ala Tyr Cys His Pro	
	115 120 125
His Val Leu Lys Asn Lys Ile Lys Asn Lys Lys Pro Ser Asn Tyr Val	
	130 135 140

00196PCT1.ST25

Leu Tyr Ser Lys Glu His Ser Tyr Ile Ser Leu His Cys Ile Leu Thr
 145 150 155 160
 Thr Ile Leu Cys Ser Ile Cys Phe Thr Pro Phe Leu Leu Cys Phe Val
 165 170 175
 Tyr Lys Glu Met Ser Pro Arg Glu Leu Asn Gly Leu Pro Gln Leu Val
 180 185 190
 Lys Leu Lys Leu Gln Ser Arg Ser Phe Tyr Phe Gln Ile His Asn Leu
 195 200 205
 Gln Pro Ser Val Glu Ser Tyr Asn Glu Ile Met Val Arg Gly Leu Ser
 210 215 220
 Ile Ser Val Gln Val Cys Pro Ala Pro Thr Thr Ser Ile
 225 230 235

<210> 73
 <211> 224
 <212> PRT
 <213> Homo sapiens

<400> 73

Ser Val His Cys Tyr Gln Glu Asn Asn Ala Phe Ser Gly Ser Leu Ile
 1 5 10 15
 Leu Asn Thr Leu Ala Gly Asn Leu Leu Ala Arg Thr Gly Asp Leu Ile
 20 25 30
 Ile Ser Ser Trp Met Arg Leu Trp Gly Gly Arg Ile Leu Thr Gly Tyr
 35 40 45
 Thr Ala Ala Gln Thr Arg Val Ala Leu Gly Arg Arg Glu Gly Glu Asn
 50 55 60
 Trp Val Asn Pro Met Met Pro Val Met Thr Asp Val Gly Leu Leu Asn
 65 70 75 80
 Lys Phe Ser Ser Gln Lys Leu Met Ile Phe Thr Ile Pro Ile Trp Ile
 85 90 95
 Ser Tyr Gly Glu Ile Gln Val Trp Leu His Ser Phe Ser Leu Ser Ile
 100 105 110
 His Thr Leu Ile His Tyr Leu Leu Glu Ala Asn Phe Val Pro Gly Leu
 115 120 125
 Val Arg Tyr Gly Val Thr Ser Cys Thr Lys Gln Pro Gly Ser Leu Gly
 130 135 140
 Pro Thr Val Gly Lys Gln Gly Lys Cys Gly Arg Ile Ile Lys Ile Thr
 145 150 155 160
 His Thr Ala Pro Arg Trp Gln Gly Lys Cys His Phe Phe Tyr Phe Leu
 165 170 175
 Leu Met Asp Leu Arg Leu Phe Trp Phe Gln Trp Ser His Phe Ser Leu
 180 185 190
 Ser Ile Gln Phe Ile Gln Asn Ser Phe Ala Ser Asp Lys Ile Ala Asn
 195 200 205

00196PCT1.ST25

Trp Leu Pro Ala Asn Ser Phe Ser Pro Gln Ser Met Gly Asn Ala Gly
 210 215 220

<210> 74
 <211> 216
 <212> PRT
 <213> Homo sapiens

<400> 74

Leu Leu Leu Leu Lys Val Ile Ala Phe Arg Leu Phe Gln Leu Gln Ser
 1 5 10 15

Lys Glu Val Tyr Val Tyr Ile Val Ile Cys Glu Tyr Thr His Thr Tyr
 20 25 30

Thr Tyr Phe Tyr Met Ser Ser Ile Phe Lys Leu Ser Arg Ile Val His
 35 40 45

Thr Asp Ile Ser Asn Pro Asn Gln Leu Pro Gln Gly Leu Phe Arg Pro
 50 55 60

Phe Ser Leu Gly Ser Leu Gln Leu Leu Leu Gln Gln Leu Glu Ile Trp
 65 70 75 80

Leu Pro Tyr Ser Phe Ala Leu Phe Asn Ser Ser Thr His Lys Trp Trp
 85 90 95

Leu Gln Asn Leu Ile Pro Pro Trp Glu Ile Thr Leu Leu Thr Lys Val
 100 105 110

Gln His Leu Cys Ile Val Leu Phe Glu Phe Leu Asp Leu Glu Ile Pro
 115 120 125

Leu Leu Phe Gln Ser Tyr Leu Gly Gln Asn His Phe Pro Phe Phe Ser
 130 135 140

Glu Val Val Leu Cys Ile Cys Asn Thr Val Arg Leu Phe Cys His Met
 145 150 155 160

Val His Ser Ile Leu Gly Phe Pro Ile Ser Phe Phe Asn Ile Cys Ile
 165 170 175

Tyr Val Ser Phe Phe Cys Ala Val Ser Phe Tyr Gly Phe Gln Leu Met
 180 185 190

His Ser Val Met Asn Leu Pro Pro Glu His His Thr Glu Phe His Gln
 195 200 205

Leu Lys Lys Phe Pro Met Phe Tyr
 210 215

<210> 75
 <211> 204
 <212> PRT
 <213> Homo sapiens

<400> 75

Phe Leu Pro Leu Cys His Asn Gly His Asp Asp Ser Trp Leu Thr Gln
 1 5 10 15

Thr Phe Cys Val Trp Lys Asp Leu Ile Cys Pro Phe Leu Glu Ala Thr
 20 25 30

00196PCT1.ST25

Ile Leu Arg Phe Glu Lys Ser Phe Leu Lys Asn Lys Ile Phe Leu Ile
 35 40 45
 Lys Asn Asn Ala Ser Ser Leu Glu Lys Asn Lys Ile Asn Lys Ser Thr
 50 55 60
 Ile Phe Leu Asn His Leu Lys Met Thr Ile Val Ser Phe Phe Phe Phe
 65 70 75 80
 Leu Val Leu Phe Ser Val Ser Asn Leu Phe Ser Ile Lys Thr Ser Glu
 85 90 95
 Met Leu Gln Arg Ile Arg Gly Pro His Ile Glu Lys Phe Ile Asn Thr
 100 105 110
 Leu Ala Ser Cys Leu Ala Phe Val Pro Ser Leu Thr Gly Asn Ser Phe
 115 120 125
 Ser Ile Ser Leu Lys Leu Gln Ile Leu Asp Asn Ser Ser Arg Ser Ser
 130 135 140
 Ser Asn Val Leu Leu Asp Ser Ser Gln Gln Glu Leu Ile Tyr Phe Leu
 145 150 155 160
 Cys Ile Phe Val Pro Gln Asp Leu Leu Ser Tyr Gly Asn Tyr His Leu
 165 170 175
 Leu Pro Tyr Ile Thr Ile Phe Glu Ser Ser Asn Lys Val Phe Phe Phe
 180 185 190
 Phe Gln Met Lys Ser Arg Tyr Ile Ala Gln Ala Gly
 195 200

<210> 76
 <211> 228
 <212> PRT
 <213> Homo sapiens

<400> 76

Val Met Gly Asn Ala Arg Ile Cys Val Gln His Gly Arg Glu Ser Val
 1 5 10 15
 Trp Lys Ser Phe Asp Lys Leu Trp His Leu Ser Leu Thr Leu Pro Gln
 20 25 30
 Asn Phe Arg Leu Pro Ala Ile Tyr Lys Leu Glu Val Lys Ile Thr Ser
 35 40 45
 Met Tyr Thr Ser Gln His Lys Glu Ser Tyr Pro Ser Phe Leu Asp Gly
 50 55 60
 Ala Arg Ile Trp Val Arg Phe Ile Val Gln Ser Ser Ser Leu Phe Tyr
 65 70 75 80
 Arg Pro Gly Phe Lys Phe Thr Ser Lys Met Glu Asn Phe Gly Trp Glu
 85 90 95
 Asn Tyr Met Trp Glu Asp Ile Phe Ser Gly Asp Phe Ser Asn Tyr Ser
 100 105 110
 Phe Ser Tyr Asp Pro Thr Pro Phe Leu Leu Asp Ser Ala Pro Cys Trp
 115 120 125
 Pro Glu Ser Leu Glu Ile Asn Tyr Val Leu Ile Ile Ile Tyr Ala Leu

00196PCT1.ST25

130 135 140
 Met Phe Leu Leu Asn Val Met Asn Ser Leu Pro Met Leu Val Ile Leu
 145 150 155 160
 Phe Ser Val Ser His Cys His Arg Cys Leu Pro Ala Asp Pro Gly Leu
 165 170 175
 Gly Arg Pro Val Leu Phe Pro Asp Ile Ala His Leu Gly Cys Leu Gln
 180 185 190
 Glu Met Ala Gly Ile Phe Gly Thr Ile Cys Ala Arg Trp Ser Ser Ser
 195 200 205
 Arg Lys Ser Thr Ser Thr Gly Gly Ile Leu Leu Leu Ala Cys Arg Ser
 210 215 220

Met Gly Leu Leu
 225

<210> 77
 <211> 220
 <212> PRT
 <213> Homo sapiens

<400> 77

Val Leu Thr Thr Ser Thr Val Phe Leu Lys Gln Asn Cys His Leu Leu
 1 5 10 15
 Glu Arg Lys Ile Tyr Gly Glu Ser Pro Ser Ser Ser Leu Thr Pro Glu
 20 25 30
 Lys Ala Trp Ile Lys Asn Ser Arg Gln Pro Trp Arg Leu Ser Leu Leu
 35 40 45
 His Gly Thr Met His Pro Trp Gly Arg Gln Lys Met Glu Lys Cys Ile
 50 55 60
 Ile Ile Lys Cys Leu Leu Cys Thr Arg Ser Gln His Phe His Met Tyr
 65 70 75 80
 Ser His Pro Ala Pro Phe His Ile Cys Ser His Phe Pro Asp Glu Gly
 85 90 95
 Thr Glu Ile Pro Arg Arg Glu Val Thr Ser Gly Gln Ser Trp Asp Leu
 100 105 110
 His Thr Ala Arg Lys Ser Thr Ala Asp Ile Asp Cys Val Leu Pro Leu
 115 120 125
 Cys Gln Leu Leu Phe Glu Gly Val Ser Arg Phe Gln Leu Ile Phe Ser
 130 135 140
 Gln Lys Cys His Gly Asp Asp Glu Glu Thr Glu Ala Lys Tyr Leu Ala
 145 150 155 160
 Val Ala Gln Leu Pro Asp Asp Gly Val Arg Ile Gln Tyr Trp Gln Cys
 165 170 175
 Trp Val Gln Ser Gln Val Leu Leu Thr Leu His Pro Val Cys Tyr Pro
 180 185 190
 Leu Ser Thr Ala Ser Gln Arg Lys Thr Tyr Thr His Gly Ala Phe Met
 195 200 205

00196PCT1.ST25

Leu Phe Gly Asn Val Gln His His Gly Asn Ile Ile
 210 215 220

<210> 78
 <211> 157
 <212> PRT
 <213> Homo sapiens

<400> 78

Lys Ile His Ser Ala Ala Gly Arg His Arg Ala Phe Ser Thr Cys Ser
 1 5 10 15
 Ser His Leu Thr Val Val Leu Leu Gln Tyr Gly Cys Cys Ala Phe Met
 20 25 30
 Tyr Leu Cys Pro Ser Ser Ser Tyr Asn Pro Lys Gln Asp Gln Phe Ile
 35 40 45
 Ser Leu Val Tyr Thr Leu Gly Thr Pro Leu Leu Asn Pro Leu Ile Tyr
 50 55 60
 Ala Leu Arg Asn Ser Glu Met Lys Gly Ala Val Gly Arg Val Leu Thr
 65 70 75 80
 Arg Asn Cys Leu Ser Gln Asn Ser Glu Arg Arg Gly Asp Ser Leu Ser
 85 90 95
 Gly Lys Tyr Leu Val Pro Ala His Gln Ile Cys Met Lys Leu Arg Phe
 100 105 110
 Leu Ser Phe Gly Val Lys Thr His Leu Lys Asp Gly Ile Asn Tyr Met
 115 120 125
 Asp Thr Val Tyr Val Cys Gln Arg Phe Leu Asn Ile Ser Thr Ile Leu
 130 135 140
 Cys Asn Phe Ser Ser Trp Lys Glu Leu His Glu His Lys
 145 150 155

<210> 79
 <211> 227
 <212> PRT
 <213> Homo sapiens

<400> 79

Ile Lys Ile Arg Leu Gly Leu Lys Leu Ser Leu Pro Leu Ser Arg Glu
 1 5 10 15
 Met Lys Cys Thr Leu Ser Thr Ile Leu Ile Leu Lys Leu Phe Lys Lys
 20 25 30
 Cys Phe Arg Asp Ser Leu Pro Asp Lys Leu Ala Met Asn Phe Gln Pro
 35 40 45
 Thr Arg Ala Phe Ile Tyr Ile Arg Gly Val Gln Glu Phe Arg Gln Leu
 50 55 60
 Phe Thr Leu Lys Lys Ile Leu Ile Val Lys Thr Thr Lys Val Asp Gln
 65 70 75 80
 Leu Ile Leu Phe Leu Trp Leu Leu Val Phe Ser Lys Val Leu Ile Leu
 85 90 95

00196PCT1.ST25

Leu Tyr Leu Ala Val Ser Lys Phe Gln Lys Cys Phe Cys Thr Asp Trp
 100 105 110
 Pro His Phe Lys Phe Ser Ile Gly Asn Phe Lys Trp Val Leu Met Leu
 115 120 125
 Pro Gly Val Leu Gly Leu Ile Leu Asp Phe Ser Val Phe Ser Leu Ser
 130 135 140
 Cys Phe Phe Met Thr Ile Leu Cys Leu Pro Ser Leu Leu Lys Phe Pro
 145 150 155 160
 Lys Asp Val Phe Tyr His Pro His Ala Gln Leu Met Asn Leu Ser Ser
 165 170 175
 Tyr Phe Ala Glu Ile Met Arg Ala Ile Arg Ser Ser His His Cys Ser
 180 185 190
 Trp Gly Ile Ile Cys Leu His Phe Gln Gln Arg Pro Cys Ser Ser Pro
 195 200 205
 Arg Pro Thr Leu Leu Ala Trp Ala Ala Ile Thr Glu His His Arg Leu
 210 215 220

Gly Gly Leu
225

<210> 80
 <211> 164
 <212> PRT
 <213> Homo sapiens

<400> 80

Ser Leu Ser Ser Arg Gly Ser Glu Ala Gln Asn Cys Leu Glu Ile Cys
 1 5 10 15
 Pro Ser Ser Asp Thr Glu Leu Met Leu Glu Arg Glu Pro Asn Leu Phe
 20 25 30
 His Leu Asn Ser Cys Gly Lys Met Asn Thr Asn Cys Phe Leu Tyr Tyr
 35 40 45
 Asp Asn Lys Lys Leu Ser Ser Ile Phe Leu Tyr Lys Lys Ala Ile His
 50 55 60
 Met His Gln Ser Gly His Leu Leu Val Thr Phe Phe Pro His His Phe
 65 70 75 80
 Thr Thr Phe His Phe Thr Thr Cys Cys Leu Asn Pro Leu Ile His Phe
 85 90 95
 Phe Lys Lys Glu Asn Glu Phe His Tyr Tyr Gln Thr Pro Gly Ser Ser
 100 105 110
 Cys Asp Gln Leu Phe Leu Val Val Lys Cys Cys Pro Glu Thr Lys Val
 115 120 125
 Asn Leu Ser Val Leu Leu Cys His Asn Arg Thr Phe Pro Val Arg Arg
 130 135 140
 Glu Cys Gly Arg Phe Gly Val Asn Pro Gly Met Gly Gln Gly Arg His
 145 150 155 160

00196PCT1.ST25

Lys Ser Arg Asn

<210> 81
 <211> 221
 <212> PRT
 <213> Homo sapiens

<400> 81

Leu Glu Phe Tyr Ser Lys His Gln Ser Arg Gly Ile Val Arg Glu Arg
 1 5 10 15

Asn Met Leu Ile Gln Asp Ser Gly Ser Leu Phe Phe Ser Ser Phe Phe
 20 25 30

Ser Gln Asn Asp Leu Asp Ser Cys Lys Val Leu Val Tyr Leu Val Ser
 35 40 45

Lys Ser Leu Phe Leu Leu Asn Phe Ile Cys Ile Asn Gln Leu Tyr Met
 50 55 60

Thr Lys Met Ser Pro Lys Phe Lys Ser Leu His Ser Lys Ala Leu Tyr
 65 70 75 80

Val His Leu Ala Ser Phe Gln Lys Thr Lys Ala Val Val Leu Lys Phe
 85 90 95

Ser Cys Thr Leu Ile Thr Gly Lys Leu Phe Lys Leu Leu Met Thr Lys
 100 105 110

Pro His Val Arg Leu Ile Tyr Ala Glu Ser Leu Gly Gln Gly Pro Arg
 115 120 125

Tyr Gln His Phe Leu Lys Leu Arg Asn Asn Gln Gly Glu Pro Leu His
 130 135 140

Lys Met Val Asn Ala Thr Phe Ile Val Ile Phe Phe Lys Ile Met Val
 145 150 155 160

Glu Leu Ile Leu Ile Leu Val Pro Ser His Gly Asn Phe Phe Arg Leu
 165 170 175

Arg Glu Phe Ile Leu Ala Leu Arg Leu Leu Lys Asn Leu Glu Ile Gln
 180 185 190

Val Phe Leu Phe Ile Phe Phe Leu Ile Leu Glu Tyr Ala Ser Ala His
 195 200 205

Pro Tyr Leu Ile Ile Leu Glu Lys Tyr Ile Lys Thr Phe
 210 215 220

<210> 82
 <211> 216
 <212> PRT
 <213> Homo sapiens

<400> 82

Ile Ile Ile Met Leu Ile Leu His His Ile Gln Ile Asp Cys Asn Ile
 1 5 10 15

Val Ile Cys Asn Ile Leu Phe Lys Ile Asn Leu Ser Glu Ser Tyr Ile
 20 25 30

00196PCT1.ST25

Ala Thr Val Val Ser Leu Ile His Arg Phe Ile Phe Tyr Gly Phe Ser
 35 40 45

Tyr Leu Leu Ser Thr Arg Ile Gln Gln Tyr Tyr Met Gly Lys Ser Gln
 50 55 60

Lys Thr Val Cys Lys Phe Phe Val Arg Cys Ser Gly Gln Arg Asp Lys
 65 70 75 80

Ile Ser Cys Cys Ser Ser Leu Ser Cys Leu Asn Met Asn Tyr Pro Leu
 85 90 95

Ser Ser Ile Ser Thr Leu His Met Leu Pro Ser His Ser Ser Phe Ser
 100 105 110

Ser Cys Phe Asp Tyr Leu Ile Glu Lys Thr His Ser Ile Tyr Arg Val
 115 120 125

Phe Tyr Gly Ala Arg Glu Asn Phe Leu Phe Val Leu Arg Phe Thr Glu
 130 135 140

Asn Ser Thr His Lys Gly Arg Leu Ile Gly Met Lys Val Lys Lys Lys
 145 150 155 160

Ile Tyr His Gln Trp Arg Leu Gln Ser Asp Tyr Ser Ile Ala Ile Asn
 165 170 175

Gly Leu Gln Trp Leu Lys Tyr Arg Phe Glu Val Thr Lys Arg Val Glu
 180 185 190

Val Leu Gly Ser Trp Gln Asn Arg Leu Trp Glu Glu Glu Lys Arg Asn
 195 200 205

Pro Gly Gln Arg Ser Ser Cys Asp
 210 215

<210> 83
 <211> 118
 <212> PRT
 <213> Homo sapiens

<400> 83

Phe Phe Pro Leu Ser Val Ser Leu Met Leu Ser Ser Lys Trp Arg Trp
 1 5 10 15

Arg Gly Phe Thr Ser Leu Phe Ser Asn Ser Pro Phe Phe Gly Phe Phe
 20 25 30

Ser Ser Thr Ser Lys Ser Val Gln Asn Val Pro Leu Ala His Arg Lys
 35 40 45

Ser Phe Leu Asp Pro Ala Thr Tyr Leu Thr Lys Ile Pro His Phe Ser
 50 55 60

Ser Ser Phe Lys Ile Ser Phe Ile Met Val Cys Val Asn Gly His Ile
 65 70 75 80

His Leu Ile His Ser Phe Leu Lys Phe Gln Lys Asn Gly Phe Val Ser
 85 90 95

Cys Tyr Phe Asn Gly Ile Ile Phe Pro Lys Ile Asn Arg Thr Phe Pro
 100 105 110

Gln Ala Gln Ser Ser Arg

00196PCT1.ST25

115

<210> 84
 <211> 147
 <212> PRT
 <213> Homo sapiens

<400> 84

```

Ile Glu Ile Ile Cys Thr Leu Leu Pro Leu Glu Asn Asn Glu Lys Leu
1      5      10      15
Gly Ile Ser Gln Cys Tyr Leu Leu Val Ala Ser Gly Ile Lys His Asn
20      25      30
Gln Asn Gly Ser Gly Gln Cys Thr Pro His Phe Lys Ala Cys Asn Ser
35      40      45
Glu Val Glu Pro Arg His Leu Pro Leu Val Val Tyr Ser Val Tyr Leu
50      55      60
Ile Asp Ser Pro Lys Cys Lys Leu Leu Ile Asn Arg Ala Tyr Val Arg
65      70      75      80
Ser Pro Val Met Cys Leu Ile Leu Ser Asp Val Cys Ser His His Thr
85      90      95
Ser Phe Gly Val Cys Asn Ser Phe Val Cys Gly Phe Phe Cys Leu Val
100     105     110
Ile Leu Val Cys Pro Val Cys Phe Tyr Gly Arg Val Trp Arg Asn Ser
115     120     125
Lys Ala Ile Pro His Cys Pro Ser Ser Phe Pro Trp Ile His Val Pro
130     135     140
Tyr His Val
145

```

<210> 85
 <211> 202
 <212> PRT
 <213> Homo sapiens

<400> 85

```

Thr Ser Leu Cys Ala Ser Val Ala Lys Ser Met Arg Ala Gly Lys Thr
1      5      10      15
Cys Ile Leu Ser Cys Ile Cys Ile Gln Met Leu Asp Pro His Leu Cys
20      25      30
Pro Val Gln Tyr Leu Ser Leu Leu Leu Gln Trp Val Thr Asn Glu
35      40      45
Pro Cys Leu Pro Ala Trp Gly Arg Arg Gly Leu Arg Asp Ile Ser Thr
50      55      60
Gly Ile Phe Gly Val Ser Arg Leu Glu Arg Asn Leu Leu Ile Ser Thr
65      70      75      80
Leu Tyr Asn Tyr His Asn Ile Leu Phe Leu Met Lys Gln Gln Phe Thr
85      90      95
Phe Leu Cys Trp Leu Tyr Phe Ala Ser Phe Thr Trp Gln Tyr Leu Met

```

00196PCT1.ST25

100 105 110
 Pro Ser Leu Gly Ile Arg Arg Lys Thr Arg Pro Gln Ile Pro Gly Pro
 115 120 125
 Ser Thr Leu Phe Leu Leu Gly Thr Ser Phe Thr Ser Ser Ser Ala Asp
 130 135 140
 Ala Pro Leu Leu Pro Thr Pro Pro Arg Lys Val Ser Ser Gln Gln Ala
 145 150 155 160
 Leu Thr Lys Gly Ser His Phe Leu Pro Lys Gly Glu Ser Ser Gln Ala
 165 170 175
 Val Asn Phe Ser Asn Phe Cys His Cys Ser Ser Val Ala Asp Leu Pro
 180 185 190
 Ser Ser Leu Ser Trp Arg Ile Leu Pro Gly
 195 200

<210> 86
 <211> 189
 <212> PRT
 <213> Homo sapiens

<400> 86

Leu Asn Ala Thr Pro Phe Ser Ser Glu Thr Leu Trp Cys Ile Leu Gly
 1 5 10 15
 His Tyr Leu Ser Lys Gly Pro Lys Leu Asn Ser Ser His His Pro Ser
 20 25 30
 Phe Phe Cys Leu Arg Phe Tyr Phe Pro Asn Gln Ile Trp Val Asn Phe
 35 40 45
 Gln Pro Leu Ser Val Ser Tyr Phe Gln Ser Asn Lys Thr Cys Met Asp
 50 55 60
 Leu Phe Cys Ile Ser Ser Asn Val Ile Ile His Ser Lys Ser His Cys
 65 70 75 80
 Leu Thr Ile Ser Leu Pro Ile Ala Leu Ala Ile Lys Lys Leu His Trp
 85 90 95
 His Gly Phe Gln Thr Cys Ile Leu Phe Phe Gly Gly Leu Ile Leu Asn
 100 105 110
 Leu Lys Tyr Leu Arg Ile Ser Asn Thr Ile Phe Lys Met Gln Gln Ile
 115 120 125
 Phe Lys Thr Ala Ser Leu Cys Gln Ala Lys Gly Val Ser Cys Gln Leu
 130 135 140
 Ser Leu Thr Ala Lys Glu Ala Lys Ile Ile Leu Met Val Val Leu Lys
 145 150 155 160
 Glu Ala Ser Ala His Phe Leu Gly Gln Cys His Pro Thr His Leu Leu
 165 170 175
 Gln Gly Leu Asp Thr Lys Gly Asp Val Ser Asp Phe Pro
 180 185

<210> 87
 <211> 191

00196PCT1.ST25

<212> PRT

<213> Homo sapiens

<400> 87

Asn Arg Lys Asn Leu Lys Ile Ser Thr Val Phe Asn Gln Phe Phe Ser
 1 5 10 15
 Leu Leu Pro Val Leu Trp His Asn Ile Val Leu Asn Trp Lys Asn Thr
 20 25 30
 Met Leu Ala Phe Thr Tyr Met Ser Ile Leu Ile Leu Ser Arg Cys Leu
 35 40 45
 Val Ser Pro Tyr Leu Lys Leu Leu Leu Ile Ile Leu Phe Cys Ser Leu
 50 55 60
 Tyr Val Leu Trp Ala Asn Lys Ser Tyr Pro Pro Asn Lys Leu Thr Phe
 65 70 75 80
 Lys Lys Phe Ala Lys Asp Trp Leu Pro Ile Ser Leu Tyr Leu Leu Ile
 85 90 95
 Pro Phe Lys Ala Lys Tyr Cys Phe Ala Thr Ile Leu Leu Leu His Tyr
 100 105 110
 Thr Glu Leu Pro Ala Leu Phe Ser Ala Lys Trp Lys Ala Tyr Phe Ser
 115 120 125
 Lys Ser Tyr Val His Leu Leu Leu His Asp Ile Asn Lys His Asn Thr
 130 135 140
 Ser Ile Thr His Phe Thr Asn Ala Arg Leu Ala Lys Asn His Thr Tyr
 145 150 155 160
 Lys Trp Pro His Leu Leu Tyr Pro His Pro Gly His Val Leu Ser Leu
 165 170 175
 Pro Trp Lys Pro Met Glu Lys Leu Arg Thr Leu Glu Arg Met Trp
 180 185 190

<210> 88

<211> 194

<212> PRT

<213> Homo sapiens

<400> 88

Lys Lys Phe Leu Arg Glu Gln Ile Cys Asp Phe Ile Met Ser Phe Ile
 1 5 10 15
 Met Phe Cys Ser Phe Gln Ile Gln Met Ser Ile Ile Cys Phe Tyr Asp
 20 25 30
 Gln Ser Ile Ile Pro Cys Lys His Ile Ser Ala Leu Ile Leu Phe Leu
 35 40 45
 Asn Asn Thr Gly Asn Val Ile Cys Cys Lys Leu Leu Thr Phe Val Arg
 50 55 60
 Lys Phe Cys Phe Thr Glu Tyr Val Arg Cys Arg Gln Asn Ile Asn His
 65 70 75 80
 Cys Phe Ile Phe Met Val Glu Glu Lys Ser Ile Ala Cys Ser Pro Phe
 85 90 95

00196PCT1.ST25

Ala Val Tyr Lys Gly Glu Phe Tyr Cys Leu Asn Ser Phe Ile Phe Trp
 100 105 110
 Pro Val Gln Glu Thr Phe Ile Ser Lys Ile Trp Met Tyr Val Phe His
 115 120 125
 Ile Leu Glu Phe Ile Val Trp Lys Asn Thr Ile Lys Val Asp Gln Lys
 130 135 140
 Ile Leu Lys Ile Leu Thr Ser Cys Leu Ser Tyr Val Lys Val Leu Trp
 145 150 155 160
 Leu Ile Leu Phe Ile Leu Ser Cys Ser Leu Ala Gly Tyr Trp Gln Thr
 165 170 175
 Gln Ser Phe Cys Phe His Lys Glu Leu Met Lys Arg Thr Ile Gly Lys
 180 185 190

Pro Thr

<210> 89
 <211> 218
 <212> PRT
 <213> Homo sapiens

<400> 89

Gln Ser Gln Pro Ser Leu Pro Gly Ser Met Gly Asp Glu Leu Ala Pro
 1 5 10 15
 Cys Pro Val Gly Thr Thr Ala Trp Pro Ala Leu Ile Gln Leu Ile Ser
 20 25 30
 Lys Thr Pro Cys Met Pro Gln Ala Ala Ser Asn Thr Ser Leu Gly Leu
 35 40 45
 Gly Asp Leu Arg Val Pro Ser Ser Met Leu Tyr Trp Leu Phe Leu Pro
 50 55 60
 Ser Ser Leu Leu Ala Ala Ala Thr Leu Ala Val Ser Pro Leu Leu Leu
 65 70 75 80
 Val Thr Ile Leu Arg Asn Gln Arg Leu Arg Gln Glu Pro His Tyr Leu
 85 90 95
 Leu Pro Ala Asn Ile Leu Leu Ser Asp Leu Ala Tyr Ile Leu Leu His
 100 105 110
 Met Leu Ile Ser Ser Ser Ser Leu Gly Gly Trp Glu Leu Gly Arg Met
 115 120 125
 Ala Cys Gly Ile Leu Thr Asp Ala Val Phe Ala Ala Cys Thr Ser Thr
 130 135 140
 Ile Leu Ser Phe Thr Ala Ile Val Leu His Thr Tyr Leu Ala Val Ile
 145 150 155 160
 His Pro Leu Arg Tyr Leu Ser Phe Met Ser His Gly Ala Ala Trp Lys
 165 170 175
 Ala Val Ala Leu Ile Trp Leu Val Ala Cys Cys Phe Pro Thr Phe Leu
 180 185 190

00196PCT1.ST25

Ile Trp Leu Ser Lys Trp Gln Asp Ala Gln Leu Glu Glu Gln Gly Ala
 195 200 205

Ser Tyr Ile Leu Pro Pro Ser Met Gly Thr
 210 215

<210> 90
 <211> 223
 <212> PRT
 <213> Homo sapiens

<400> 90

His Phe Lys Ile Asn Leu Phe Pro Val Asn Leu Cys Ser Ser Ser His
 1 5 10 15

Pro Leu Phe Asn Glu Leu Pro Pro Phe Pro Thr Leu Phe Leu Ala Phe
 20 25 30

Ile Pro Met Val Pro Leu Lys Val Phe Ser Ser Ser Leu Pro Phe Ser
 35 40 45

Pro Pro Val Phe Ser Gly Val Asn Gly Ala Ala Asn Ser Pro Ser Ser
 50 55 60

Ser Cys Leu Asn Arg Ser Ser Ser Pro Thr Pro Ala Ala Ala Pro Tyr
 65 70 75 80

Ser Gln Ser Gln Ser Pro Val Cys Val Ile Ala Gly Met Ser Leu Glu
 85 90 95

Ser Thr Asn Ile Leu Tyr Ser His Thr Cys Leu Pro Pro Met Ser Ser
 100 105 110

Ala Pro Leu Leu Val Ser Glu Phe Gln Val Gly Pro Val Pro Phe Phe
 115 120 125

Leu Pro Cys Arg Leu Ser Arg Thr Arg Ser Leu Pro Thr Ser Asp Phe
 130 135 140

Leu Ser Asp Asp Phe Trp Gly Phe Ser Ile Cys Leu Leu Glu Gly Pro
 145 150 155 160

Leu Gly Asp Phe Tyr Gly Thr Leu Ile Ala Ser Phe Leu Tyr Leu Arg
 165 170 175

Asn Val Phe Leu Leu Leu Glu Thr Pro Lys Ile His Asp Ile Phe Phe
 180 185 190

Thr Lys Leu Phe Leu Leu Ser Pro Ala Phe Asn Lys Ser Leu Phe Ala
 195 200 205

Lys Lys Trp Cys Arg Phe Phe Thr Thr Ala Ser Glu Lys Ser Val
 210 215 220

<210> 91
 <211> 193
 <212> PRT
 <213> Homo sapiens

<400> 91

Phe Pro Arg Ile Val Cys Thr Val Thr Gly Val Ala Val Tyr His Ser
 1 5 10 15

00196PCT1.ST25

Ile Tyr Thr Ser Ile Trp His Thr Ala Gly Ala Ser Gly Thr Thr Tyr
 20 25 30
 Gln Ser Val Ser Leu Pro Asp His Phe His Asp Val Leu Ser Tyr Leu
 35 40 45
 Pro Cys Asn Lys Leu Val Asn Val Tyr Asp Cys Phe Val Ile Pro Met
 50 55 60
 Gln Ser Cys Asn Asn Asn Met Tyr Phe Lys Asn Leu Gly Ile Phe Leu
 65 70 75 80
 His Thr Ile Ser Ser Ile His Ile Asn Glu Lys Ser Lys Leu Gly Val
 85 90 95
 Ser Val Lys His Trp Ile Phe Thr Met Leu Ile Gly Val Pro Phe Ile
 100 105 110
 Ile Ala Ala Tyr Arg His Ile Ala Ile Val Pro Cys Thr Phe Asn His
 115 120 125
 Gln Cys Cys Gln Ala Ser Lys Ala Val Asn Val Tyr Leu Gly Leu Ile
 130 135 140
 Ile Arg Ile Thr Arg Asn Asn Phe Phe Asn Phe Asn Ile Leu Phe Phe
 145 150 155 160
 His Arg Leu Leu Gly Tyr Arg Cys Cys Leu Ile Thr Val Leu Tyr Trp
 165 170 175
 Phe Glu Arg Phe Gly Cys Thr Gln His Pro Ser Ser Ile His Tyr Ser
 180 185 190

Leu

<210> 92
 <211> 191
 <212> PRT
 <213> Homo sapiens

<400> 92

Gly Leu Phe Arg Glu Pro Leu Glu Ile Pro Pro Pro Trp His Gln Leu
 1 5 10 15
 Pro Pro Pro Pro Glu Leu Thr Val Ser Ser Leu Asp Ala Ala Pro Gly
 20 25 30
 Lys Val Ile Asn Asn Gln Val Ser Lys Gln Cys Trp Ala Val Phe Leu
 35 40 45
 Ile Leu Pro Phe Pro Asn Trp Val Leu Phe Gly Lys Leu Leu Ser Tyr
 50 55 60
 Phe Ile Cys Thr Met Gly Tyr Thr Tyr Ala Phe Tyr Ile Trp Leu Leu
 65 70 75 80
 Arg Arg Leu Ser Asp Met His Thr Lys Asn Ala Glu Gln Asn Thr Leu
 85 90 95
 Ser Ile Ser Phe Leu Ser Val Ile Lys Trp Arg Pro Leu Arg Leu Ser
 100 105 110
 Asn Leu Leu Leu Leu Trp Leu Ile Leu Val Leu Ile Leu Ile Tyr Lys

00196PCT1.ST25

115 120 125
 Leu Cys Cys Ile Trp His Met Val His Val His Glu Tyr Val Leu Tyr
 130 135 140
 Lys Gly Met Lys Asn Gln Leu His Glu Lys Lys Phe Gln Ile Leu His
 145 150 155 160
 Phe Thr Asn Thr Asp Thr Lys Asn Thr Lys Ile Leu Arg Gly Lys Ser
 165 170 175
 Asp Leu Ala Thr Ser Thr Trp Ala Ser Leu Lys Val Cys Phe Trp
 180 185 190

<210> 93
 <211> 133
 <212> PRT
 <213> Homo sapiens

<400> 93

Leu Asn Leu Lys Ile Asn Arg Ala Ile Leu Asp Arg Gln Asn Phe Gly
 1 5 10 15
 Asp Ser Glu Cys Pro Arg Asn Asp Pro Met Met Phe Val Gly Phe Ile
 20 25 30
 Ile Cys Ile Arg Cys Val Leu Trp Leu Gly Phe Met Ala Cys Phe Tyr
 35 40 45
 Phe Leu Leu His Ser Thr Gly Leu Lys Arg Gln Gln Gly Gln Cys Leu
 50 55 60
 Ile Tyr Asn Val Val Leu Cys Phe Leu Asn Lys Val Pro Gln Leu Ser
 65 70 75 80
 Glu Ile Phe Met Val Asn Ile Lys Gln Ser Lys Phe Ile Cys Leu Pro
 85 90 95
 Glu Ser Leu Val Ile Tyr Leu Asp Ser Phe Arg Ile Pro Leu Asn Ile
 100 105 110
 Ile Glu Gly Cys Met Ile Phe Lys Thr Glu Met Glu Ile Met Leu Trp
 115 120 125
 Ile Asn Ala Ile Arg
 130

<210> 94
 <211> 202
 <212> PRT
 <213> Homo sapiens

<400> 94

Tyr Ala Lys Glu Leu Thr Val Trp Ala Lys Val Asn Glu Ser Leu Lys
 1 5 10 15
 Leu His Ala Lys Leu Cys Val Val Ala Cys Val Cys Val Tyr Ser Tyr
 20 25 30
 Val Phe Phe Lys Glu Val Tyr Tyr Leu Leu Asp Ser Gln Ile Val Gln
 35 40 45
 Trp Pro Gln Asn Ile Lys Thr His Val Gln Ile Gln Ser Lys Leu Arg

00196PCT1.ST25

50	55	60
Ala Val Lys Glu Ile Gln Thr Lys Asn Ser Phe Cys Pro Ser Ser Phe		
65	70	75 80
Asn Cys Leu Arg Gly Ala Trp Asp Trp Ala Thr Tyr Trp Ala Gly His		
	85	90 95
Leu Gln Arg Ile Leu Gln Gly Lys Gly Thr Gln Thr Ser Gly Leu Glu		
	100	105 110
Ser Lys Phe Lys Ser Cys Gly Val Gly Tyr Met Leu Gln Glu Ile Arg		
	115	120 125
Glu Ser Val Asn Pro Glu Ile Gly Glu Ala Asp Ser Pro Arg Lys Asp		
	130	135 140
Asn Ser Glu Trp Ser Leu Glu Gly Arg Val Arg Leu Glu Leu Glu Pro		
	145	150 155 160
Glu Val His Ala Ser Ala Ser Val Val Ser Arg Asp Met Thr Lys Leu		
	165	170 175
Glu Arg Arg Lys Ala Arg Asn Gly Trp Gly Trp Lys Leu Leu Leu Asp		
	180	185 190
Ala Ser Gln Thr Lys Gly Ile Leu Asp Pro		
	195	200

<210> 95
 <211> 178
 <212> PRT
 <213> Homo sapiens

<400> 95

Lys Leu Ser Val Phe Ile Pro Leu Gln Thr His Thr Pro Asn Ile Gln		
1	5	10 15
Trp Glu Arg Asn Asn Ile Thr Ala Glu Glu Val Ser Glu Arg His Lys		
	20	25 30
Ala Val Ile Gly Ser Leu Leu Asn Ser Pro Arg Gln Met Leu Pro Gly		
	35	40 45
Ser Leu Pro Trp Gly Gly Leu Val Ile Phe Leu Glu Val Val Ser Ser		
	50	55 60
Ser Leu Phe Ser Thr Val Leu Gln Leu Pro His Pro Ser Ser Cys Leu		
	65	70 75 80
Leu Arg Ser Leu Tyr Pro Leu Asp Ser Arg Leu Leu Leu Asp Val Leu		
	85	90 95
Thr Phe Leu Gln Gln Lys Leu Ser Leu Phe Leu Asn Leu Phe Ala Val		
	100	105 110
His Arg Lys Trp Lys Val Gln Arg Leu Leu Phe Asn Phe Leu Ser Leu		
	115	120 125
Phe Ile Ala Ser Trp Val Pro Phe Thr Tyr Ile Thr Leu Leu Lys Ser		
	130	135 140
Phe Cys Gly Leu Ser Met Tyr Gln Ile Ile Asp His Phe Ile Lys Ala		
	145	150 155 160

00196PCT1.ST25

Thr Phe Phe Val Phe Gln Thr Ser Phe Leu Tyr Phe Gly Gln Val Arg
 165 170 175

Pro Leu

<210> 96
 <211> 191
 <212> PRT
 <213> Homo sapiens

<400> 96

Met Val Phe Val Arg Ser Tyr Cys Pro Lys Ser Leu Phe Cys Pro Ser
 1 5 10 15

Tyr Asp Ile Cys Phe Asn Ile Asn Asp Ala Gln Leu Cys Leu Asp Pro
 20 25 30

Lys Arg Arg Ser Leu Tyr Asp Phe Pro Cys Cys Tyr Gly Gln Glu Phe
 35 40 45

Ser Phe Lys Leu Phe Trp Gly Leu Ala Thr Arg Gly Ser Val Gln Ser
 50 55 60

Val Gln Arg Ala Asp Leu Ser Ser Leu Ile His Ile Pro Pro Phe Trp
 65 70 75 80

Ser Lys Tyr Ala Lys Ser Ser Ile Asn Ser Gln Ala Leu Ile Ser Phe
 85 90 95

His Ile Ile Thr Arg Trp Cys Gly Tyr Leu Ser Gln Ile Tyr Ser Val
 100 105 110

Leu Gln Trp Asp Pro Tyr Ser Gln Gly Thr Tyr Ser Gln Lys Thr Tyr
 115 120 125

Ser Gln Leu Asn Ile Leu Gly Gln Lys Gly Met Glu Val Gly Arg His
 130 135 140

Ser Leu Phe Leu Lys Asn Leu Leu Ser Asn Ile Arg Ala Thr Asn Gln
 145 150 155 160

Lys Pro Lys Ser Lys Leu Thr Lys Pro Ile Tyr Leu Val Leu Cys Val
 165 170 175

Gly Pro Ser Ala Leu Arg His Leu Ala His Leu Phe Trp Arg Ile
 180 185 190

<210> 97
 <211> 91
 <212> PRT
 <213> Homo sapiens

<400> 97

Gly Arg Gly Gly Gln Gln Gly Gly Leu Gln Asn His Asp Val Phe Leu
 1 5 10 15

Thr Gly Leu Thr Ser Ala Ser Ile Cys Leu Thr Leu Gln Pro Met Ser
 20 25 30

Leu Phe Leu Val Val Ile Leu Met Gly Ala Leu Arg Ser Gln Arg Arg
 35 40 45

00196PCT1.ST25

Gly Leu Arg Arg His Cys Leu Tyr Leu Trp Ser Tyr Ile Arg His Leu
 50 55 60

Tyr Phe Val Met Asn Ser Lys Ser Ser Ser Lys Met Gln Leu Trp Gly
 65 70 75 80

Asn Ser His Arg Asn Phe Ser Gln Phe Trp Leu
 85 90

<210> 98
 <211> 201
 <212> PRT
 <213> Homo sapiens

<400> 98

Ser Arg Asp Gln Ile Thr Pro Ser Arg Ser Trp Arg Lys Asp Pro Ser
 1 5 10 15

Ser Glu Gly Thr Trp Leu Gly Gly Leu Ser Val Ser Gly Ser Cys Val
 20 25 30

Gly Ile Ser His Ser Val Gly Ala Ser Val Ile Ala Gly Trp Pro Phe
 35 40 45

Asp Asn Ala Thr Cys Lys Met Ser Gly Leu Val Gln Gly Met Ser Val
 50 55 60

Ser Ala Ser Val Phe Thr Leu Val Ala Ile Ala Val Glu Arg Glu Val
 65 70 75 80

Ser Trp Leu Asp Tyr Ala Ala Asn Gly Leu Ala Leu Arg Gly Ala Thr
 85 90 95

Ala Ser Asn Ala Gly Leu Ala Gly Arg Leu Gly Leu His His Gly Lys
 100 105 110

Trp Gly Ile Leu Ser His Lys Glu Lys Gly Pro Gly Pro Ser Cys Pro
 115 120 125

Leu Pro Lys Leu Gly Glu Pro Asp Glu Asp Thr Thr Pro Phe Trp
 130 135 140

Lys Ala Arg Pro Trp Leu Ala Phe Val Gly Ile Pro Gly Ala Cys Glu
 145 150 155 160

Glu Leu Lys Ser Ser Pro Tyr Phe Leu Ser Ser Arg Asn Pro Ala Thr
 165 170 175

Ser Lys Ser Glu Pro Gly Glu Pro Glu Leu Arg Gly Pro Ala Tyr Gly
 180 185 190

Trp Val Thr Val Trp Leu Gly Arg Lys
 195 200

<210> 99
 <211> 218
 <212> PRT
 <213> Homo sapiens

<400> 99

Thr Pro Lys Arg Leu Lys Leu Arg Ser Leu Ile Leu Ser Ser Val Lys
 1 5 10 15

00196PCT1.ST25

Glu Phe Leu Glu Ser Pro Pro Ser Leu Gly Met Phe Leu Ser Ser Trp
 20 25 30
 Phe Asn Ile Ala Ala Asp Ala Pro Ala Ile Thr Ala Thr Phe Gln Thr
 35 40 45
 Ala Lys Tyr Gly Lys Arg Met Lys Arg Arg Arg Ala Cys Leu Gly Val
 50 55 60
 Pro Cys Ile Ile Ser Ile Tyr Ile Trp Ala Glu Pro Ser His Arg Ala
 65 70 75 80
 Thr Pro Tyr Val Ser Val Ser Tyr Cys Tyr Ile Ala Thr Thr Lys Phe
 85 90 95
 Pro Cys His Thr Thr His Ile Cys Arg Leu Ala Arg Val Gln Phe Leu
 100 105 110
 His Ala Gly Leu Arg Gln Ala Val Leu Leu Arg Val Thr Val Ala Glu
 115 120 125
 Leu Ile Pro Phe Leu Thr Ala Gly Leu Cys Phe Ser Val Thr Val Pro
 130 135 140
 Cys Ala Phe His Leu Pro Trp Val Asp Glu Arg Lys Pro His Leu Ser
 145 150 155 160
 Thr Gly Leu Ala Thr Ser Val Pro His Gly Pro Lys Arg His Gln Arg
 165 170 175
 Ala Asp Arg Asn Arg Asp Leu Leu Arg Ser Arg Leu Lys Thr Gly Thr
 180 185 190
 Leu Pro Arg Leu Phe Thr Ser Tyr Pro Lys His Arg Cys Ile Thr Lys
 195 200 205
 Pro Gln Val Lys Gly Lys Lys Tyr Asn Pro
 210 215

<210> 100
 <211> 175
 <212> PRT
 <213> Homo sapiens

<400> 100

Thr Ile Ile Cys Cys Ile Phe Gln Asn Ser Cys Asn Val Ser Asn Thr
 1 5 10 15
 Lys Lys Arg Met Phe Val Val Met His Ile Ser Ser Thr Leu Ile Leu
 20 25 30
 His Ile Val Tyr Ile Tyr Gln Asn Ile Ser Ser Thr Ser Lys Ile Cys
 35 40 45
 Ser Ile Ile Val Val Gln Lys Asn Leu Asn Asn Tyr Asn Val Leu Phe
 50 55 60
 Ile Ser Lys Trp Phe Ile Arg Phe Lys Ile Phe Leu Val Phe Asn Phe
 65 70 75 80
 Phe Ile Tyr Tyr Leu Ile Pro Phe Asn Phe Leu Lys Tyr Ile Arg Ser
 85 90 95

00196PCT1.ST25

Ser Tyr Phe Arg Val Lys Phe Lys Ser Phe Glu Tyr Leu Ile Leu Gln
 100 105 110

Ser Phe Leu Pro Leu Ile Phe Pro Gln Trp Pro Val Ser Val Val Met
 115 120 125

Met Leu Leu Arg Asn Gly Leu Ala Thr Cys Thr Lys Pro Ile Leu Trp
 130 135 140

Gln Trp Phe Ser Arg Lys Glu Lys Ala Leu Leu Val Tyr Trp Gln Gly
 145 150 155 160

Asp Arg Trp Gln His Ser Asn Leu Ser Pro Thr Glu Asp Gly Gly
 165 170 175

<210> 101

<211> 184

<212> PRT

<213> Homo sapiens

<400> 101

Ser Tyr Leu Gly Pro Val His Ser Phe Ser Gln Thr Ala Ser His Ala
 1 5 10 15

Ile Pro Ser Met Lys Ile Leu Pro Phe Pro Leu Ser Phe Phe Ser Ser
 20 25 30

Leu Ile Tyr Ser Pro Val Leu Val Ser Ser Phe Pro Ser Ser Ser Gly
 35 40 45

Gln Thr Leu Phe Thr Ser Leu Thr Thr Pro Ser Lys Ile Val Leu Ile
 50 55 60

Thr Val Tyr Pro Leu Asn Thr Leu Tyr Arg Ser Trp Pro Ser Pro Asp
 65 70 75 80

Asn Val Leu Cys Ile Phe Trp Phe Thr Cys Cys Val Ser Ser Phe Leu
 85 90 95

His Cys Cys Lys Glu Ile Pro Glu Thr Gly Phe Ile Lys Lys Arg Gly
 100 105 110

Leu Ile Asp Ser Gln Phe Cys Arg Leu Tyr Gly Lys His Val Ala Gly
 115 120 125

Ile Cys Leu Ala Ser Gly Glu Asp Ser Gly Asn Leu Gln Ser Trp Gly
 130 135 140

Arg Arg Gly Ser Arg His Ile His Ser Arg Ser Ser Lys Ala Lys Gly
 145 150 155 160

Asp Val Pro His Thr Ser Lys Pro Asp Leu Met Arg Thr His Tyr His
 165 170 175

Glu Asn Ser Thr Arg Gly Trp Cys
 180

<210> 102

<211> 212

<212> PRT

<213> Homo sapiens

<400> 102

00196PCT1.ST25

Tyr Asn Asn Ser Leu Leu Tyr Ile Ser Ile Phe Cys Leu Ser Gln Val
 1 5 10 15
 Ser Thr Leu Ser Gly Ile Val Cys Ser Phe His Ser Phe Trp Leu Ser
 20 25 30
 Trp Glu Gln Gln Ser Ser Ala Thr Pro Ala Met Val Ile Val Gln Met
 35 40 45
 Ser Asn Gln Ser Ser Ile Thr Ile Arg Ser Lys Leu Gln Thr Phe Ser
 50 55 60
 Pro Leu Ala Phe Arg Ile Leu Tyr Thr Gln Phe Met Met Tyr Arg Lys
 65 70 75 80
 Cys Leu Leu Leu Phe Ser Leu Gln Leu Gly Phe Gln Lys Glu Ile Met
 85 90 95
 Ala Ser Arg Asn His Leu Tyr Leu Gln Met Ala Gly Ser Ile His Arg
 100 105 110
 Arg Ala Ile Tyr Gln Gln His Tyr Ser Met Phe Gln Pro Lys Met Ser
 115 120 125
 Leu Pro His Val Arg Gln Thr Thr Tyr Ile Gly Thr Thr Ala Val Thr
 130 135 140
 Val Phe Phe Ser Thr Phe Leu Ile Met Lys Ser Met Leu Asn Ser Thr
 145 150 155 160
 Met Ala Phe Pro Phe Ser Trp Gln Ser Thr Ala Tyr Thr Ile Leu His
 165 170 175
 Leu Thr Val Phe Ile Leu Pro Ser Gly Lys Ala Leu Trp Lys Gln Ser
 180 185 190
 Arg Gly Tyr Phe Gly Asp Leu Asn Tyr Tyr Asn Leu Leu Ser Leu Leu
 195 200 205
 Cys Phe Leu Gln
 210

<210> 103
 <211> 219
 <212> PRT
 <213> Homo sapiens

<400> 103

Ser Leu Ser Gly Gln Leu Phe Ala Leu Leu His Thr Leu Ser Ile Cys
 1 5 10 15
 Ile Ser Tyr Asn Val Tyr Arg Leu Tyr Gly Val His Ser Thr Trp Arg
 20 25 30
 Thr Phe Lys Thr Ile Ile Ala Leu Gly Phe Gly Ser Glu Phe Met Leu
 35 40 45
 Pro Cys Gln Ser Phe Leu Phe Val Thr Trp Pro Phe Lys Tyr Ala Ala
 50 55 60
 Thr Cys Asn Thr Gly His Ser Asp Pro Ile Arg Leu Met Ala Ser Cys
 65 70 75 80
 Ser Ser Arg Ser Leu Ser Val Cys Trp Tyr Ile Met Leu Gly Leu Cys

00196PCT1.ST25

85 90 95
 Ser Arg Arg Arg Glu Ala Ser Gln Leu Ala Thr Gly Tyr Lys Ser Ile
 100 105 110
 Ala Glu Asn Asp Lys Arg Gln Gly Pro Ser Leu Gln Arg Ser Ala Lys
 115 120 125
 Lys Ile Leu Asn Val Tyr Lys Asp Leu Lys Arg Asn Ser Pro Arg Gln
 130 135 140
 His Tyr Ser Val Leu Asp Tyr Gly Tyr Tyr Thr Leu Leu Gln Leu Leu
 145 150 155 160
 Cys Ser Ser Glu Gln Lys Thr Glu Asp Phe Glu Met Ser Thr Thr Pro
 165 170 175
 Ala Pro Glu Tyr Asn Gly Thr Phe His Leu Phe Leu Val Thr Phe Ile
 180 185 190
 Phe Phe Cys Cys Trp Ile Pro Tyr Ile Ile Val Ser Ile Ser Gln Ala
 195 200 205
 Ser Thr Met Val Asn Ser Gly Trp Thr Leu Pro
 210 215
 <210> 104
 <211> 208
 <212> PRT
 <213> Homo sapiens
 <400> 104
 Arg Thr Leu Tyr Trp Tyr Phe Tyr Phe Lys Phe Ser Ile Phe Gly Met
 1 5 10 15
 Ala Glu Cys Cys Tyr Lys Val Ser Arg Ser Pro Leu Pro Leu His Cys
 20 25 30
 Ala Asp Leu Leu Ser Ser Ile Gln Gly Thr Asp Leu Arg Asn Leu Gln
 35 40 45
 Val Val Thr Ser Cys Leu Val Phe Phe Leu Gly Arg Tyr Pro Ser Leu
 50 55 60
 Gln Thr Cys Arg Asn Leu Asn Leu Leu Pro Leu Thr Tyr Leu Val Pro
 65 70 75 80
 Cys Gly Leu His Phe Thr Val Cys Ala Asn Ser Leu Phe Ile Thr Ile
 85 90 95
 Leu Thr Leu Asp Ser Arg Ala Ser Pro Thr Ser Pro Phe Ser Val Thr
 100 105 110
 Leu Thr Phe Leu Leu Ser Val Thr Met Ser Asp Leu Leu Phe Ser Pro
 115 120 125
 Ile Phe Cys Pro Leu Gln Ile Leu Lys Pro Ser Phe Trp Phe Arg Pro
 130 135 140
 Leu Lys Gly Val Thr Gly Val Cys Tyr Pro Lys Val Val Pro Lys Ile
 145 150 155 160
 Ser Lys Leu Glu Lys Lys Thr Lys Asn Lys Lys Ile Pro Tyr Pro Ser
 165 170 175

00196PCT1.ST25

Trp Met Phe Leu Lys Gly Phe Leu Gly Gln Val His Val Arg Ile Ala
180 185 190

Gly Val Ser Leu Gln Lys Asp Phe Ser Trp Pro Ser Phe Val Thr Val
195 200 205

<210> 105

<211> 231

<212> PRT

<213> Homo sapiens

<400> 105

Met Lys Pro Val Leu Pro Pro Ala Lys Arg Thr Glu Ser Leu Asn Gly
1 5 10 15

Met Val Asp Ala Ala Tyr Trp Thr Val Tyr Phe Ile Leu Ala Ala Pro
20 25 30

Gly Ile Cys Val Ile Ser Leu Glu Met Phe Tyr Met Cys Leu Val Glu
35 40 45

Leu Gln Asn Asn Thr Ser Leu Asn Ile Ser Cys Ile Thr Gly Ser Ile
50 55 60

Gln Phe Ile His Asn Lys Val Ser Pro Val Leu Tyr Arg Arg Ile Tyr
65 70 75 80

Lys His Ser Val Lys Ser Ile Asp Arg Ile Gly Asp Arg Gly Leu Lys
85 90 95

Ile Lys Ile Asn Ala Phe Leu Val Leu Phe Gly Val Gly Lys Ser Asn
100 105 110

Leu Phe Phe Met Leu His Arg Ser Gln Phe Phe Val Phe Phe Glu Ser
115 120 125

Arg Pro Val Ile Gly Arg Cys Lys Glu Pro Lys Arg Lys Asn Gln Lys
130 135 140

Pro Thr Ala Ser Phe Gln Asn Arg Ser Gln Lys Arg Lys Glu Tyr Glu
145 150 155 160

Ser Ser Arg Ser Phe Asn Cys Ser Phe Ile Ile Ser Ser Arg Lys Arg
165 170 175

Gly Cys Met Ile Val Ser Lys Thr Lys Glu Glu Thr Ala Lys Glu Arg
180 185 190

Asn Val Gly Asn Leu Leu Val Glu Ala Met Thr Leu Leu Gly Glu Ile
195 200 205

Leu Ser His Phe Leu Ser Ser Cys Phe Ser Ile Met Phe Phe Thr Leu
210 215 220

Ser Ile Gln Tyr Lys Thr Leu
225 230

<210> 106

<211> 188

<212> PRT

<213> Homo sapiens

<400> 106

00196PCT1.ST25

Ser Glu Asp Leu Gln His Arg Val Lys Tyr Ala Arg Glu Gly His Ile
 1 5 10 15
 Thr Phe Ile Phe Thr Phe Ile Leu Ile Tyr Phe Leu Ser Ile Asn Leu
 20 25 30
 Phe Cys Phe Tyr Ile Ser Val Val Ala Gln Asn Ser Asn Cys Ser Lys
 35 40 45
 Asn His Ser Gly Leu Asn Thr Gly Lys Ile Ser Phe Gly Thr His Asn
 50 55 60
 Gly Leu Lys Asn Ser Cys Val Pro Phe Thr Gly Glu Ile Arg Lys Gly
 65 70 75 80
 Ile Glu Lys Phe Pro Ile Pro Pro Asn Pro Ala Ser Pro Ile Pro Ile
 85 90 95
 Ser Arg Thr Ser Phe His Leu Ile Ser Leu His Leu Gln Met Val Val
 100 105 110
 Leu Asn Leu Gln Ile Asn Lys Pro Lys Thr Glu Ser Ile Ile Phe Ser
 115 120 125
 His Leu Val Phe Pro Ser Asn Ser Leu Ile Ser Val Thr Cys Pro Ile
 130 135 140
 Thr Leu Pro Gly Ile Gln Pro Pro Lys Gln Gly Gly Leu Leu Pro Leu
 145 150 155 160
 Gln Trp Thr Pro Gly Ile Gln Val Leu Leu Leu Ala Pro Lys Cys Pro
 165 170 175
 Gln Cys Pro Val Leu Pro Asn Gln His Ile Gln Gln
 180 185

<210> 107
 <211> 230
 <212> PRT
 <213> Homo sapiens

<400> 107

His Cys Asn Gly His Cys Arg Phe Ser Arg Leu Ser Pro Glu Gly Glu
 1 5 10 15
 Trp Pro Pro Phe Lys Val Cys Ser Glu Glu Asn Thr Pro Gly Ser Arg
 20 25 30
 Ala Ile Val His Lys Asp Ala Leu Gly Ser Val Val Leu Thr Asn Val
 35 40 45
 Glu Thr Tyr Arg Ala Leu Val Ala Glu Ala His Ser Asn Gln Pro Lys
 50 55 60
 Leu Gly Arg Arg Ala Gly Ala Gln Cys Ile Trp Glu Gly His Arg Leu
 65 70 75 80
 Gly Ser Pro Ser Ser Ser Gly Pro Pro Ser Arg Met Ile Gly Leu Arg
 85 90 95
 Pro Pro Ser Gly Ser Pro Arg Arg Gln Pro Ser Ser Glu Glu Ser Gly
 100 105 110

00196PCT1.ST25

Asp Lys Arg Ser Ala His Leu His His Ser Leu Pro Glu Thr Arg Leu
 115 120 125

Asn Cys Ile Ile Cys Phe Cys Pro Thr Cys His Lys Pro Thr Ile Trp
 130 135 140

Ser Asn Ala Arg Pro His Pro Arg Lys Thr Arg Pro Gln Pro Trp Ala
 145 150 155 160

Leu Glu Gly Leu Cys Tyr His Leu Pro His Ala Leu Gln Lys Ser Asp
 165 170 175

Glu Ser Ser Pro Ile Ile Pro Thr Leu Ser Leu Arg Ser Pro Trp Met
 180 185 190

Pro Arg Gly Arg Arg Phe Asn Met Gly Gln Lys Val Ala Thr Thr Glu
 195 200 205

Leu Leu Gly Ser Ser Pro Tyr Leu Leu Ser Leu Asp Leu Leu Pro Gly
 210 215 220

Leu Gln Arg Val Lys Ser
 225 230

<210> 108
 <211> 178
 <212> PRT
 <213> Homo sapiens

<400> 108

Phe Arg Ser Lys Phe Ile Pro Val Gly Glu Gly Leu Val Glu Val Glu
 1 5 10 15

Gln Gly Gln Arg Val Gln Val Glu Tyr Ser Asn Phe Lys Asn Leu Lys
 20 25 30

Ser Glu Thr Leu Gln Asn Leu Lys Leu Phe Glu His His Asp Thr Gln
 35 40 45

Arg Lys Tyr Ser Leu Asp Ser Arg Phe Leu Tyr Leu Glu Gly Ser Thr
 50 55 60

Lys Arg Tyr Asp Ile Asn Ile Pro Lys Phe Lys Asn Ile Asn Ser Lys
 65 70 75 80

His Phe Pro Gln Ala Phe Trp Ile Lys Asp Thr Gln Thr Gly Ile Arg
 85 90 95

Ser Trp Leu Pro Glu Glu Glu Thr Gly Glu Asp Ile Pro Val Val Ala
 100 105 110

Leu Met Lys Gly Trp Gly Pro Glu Asn Gln His Pro Leu Phe Gly Cys
 115 120 125

Phe Leu Leu Trp Arg Val Ala Leu Glu Gly Gly Pro Pro Phe Ile His
 130 135 140

Val Leu Ser Gly Arg Pro Phe Thr Leu Arg Gly Ala Ser Leu Pro Cys
 145 150 155 160

Leu Asp Phe Pro Gly Leu Cys Pro Leu Ser Ala Glu Val Lys Val Ser
 165 170 175

Gly His

00196PCT1.ST25

<210> 109
 <211> 237
 <212> PRT
 <213> Homo sapiens

<400> 109

Ser Ala Ser Gln Ser Ala Gly Ile Thr Gly Met Ser His Cys Ala Gly
 1 5 10 15
 Arg Ser Leu Val Ser Phe Tyr Ser Ala Val Met Cys His Ile Thr Met
 20 25 30
 Leu Pro Ser Met Ile Asp Cys Val Tyr Asn Thr Arg Pro Val Arg Ser
 35 40 45
 Tyr Cys Thr Leu Leu Tyr Leu Phe Cys Val Glu Ile His Arg Tyr Leu
 50 55 60
 Ala Leu Cys Tyr Ser Arg Arg Gln Arg Pro Ala Gln Gln His Gly Met
 65 70 75 80
 Gln Ala Trp Gly Leu Glu Leu Thr Gly Cys Thr Thr Gly Pro Gly Val
 85 90 95
 Arg Gln Pro His Arg Leu Gly Leu Arg Glu Cys Ile His Ala Val Cys
 100 105 110
 Ala Arg Thr Arg Phe Ser Asp Arg Val Leu Ala Val Ser Leu His Met
 115 120 125
 Thr Val Leu Ile Phe Glu Trp Ser His Val Phe Gly Leu Leu Asn Arg
 130 135 140
 Met Phe Val Phe Ser Glu Lys Met Pro Ile Ala Ser His Leu Gln Leu
 145 150 155 160
 His Gln Phe Arg Phe Arg Phe Glu Leu Lys Cys Asp Leu Ser Ile Gln
 165 170 175
 Lys Lys Ser Ile Ser Thr Phe Gly Lys Ile Ser Arg Leu Lys Lys Thr
 180 185 190
 Phe Arg Val Phe Lys Arg Thr Ser Ser Val Lys Ser Ser Ile Leu Lys
 195 200 205
 Gly Cys Pro Ile Asn Lys Leu Leu Trp Asn Cys Phe Ile Ser Ala Leu
 210 215 220
 Phe Leu Cys Gly Thr His Ser Ser Lys Thr Ala Glu Asp
 225 230 235

<210> 110
 <211> 221
 <212> PRT
 <213> Homo sapiens

<400> 110

Phe Phe Leu Phe Leu Ser Leu Ser Phe Ser Phe Cys Leu Lys Ile Met
 1 5 10 15
 Lys Asn Ala Gly Ser Val Glu Arg Arg Lys Cys Pro Cys Pro Thr Ser

00196PCT1.ST25

20	25	30
Cys Arg Tyr Leu Ser Cys Phe Phe Ile Leu Leu Lys Ile Glu Leu Lys		
35	40	45
Val Phe His Phe Leu Phe Phe Asn Phe Arg Gly Tyr Asn Gly Asp Ser		
50	55	60
Gly Thr Asn Arg Lys Phe Val Phe Thr Arg Pro Val Lys Arg Val Phe		
65	70	75
Leu Leu Ile Pro Val Phe Val Ser Gly Cys Met Ala Ile Ala Ser Lys		
85	90	95
Phe Phe Pro Leu Phe Pro Ser Pro Ile Thr Gln Arg Val Ser Ser Phe		
100	105	110
Asn Thr Leu Glu Ser Ile Leu Leu Asp Ala Thr Thr His Met Cys Val		
115	120	125
Asn Glu Asn Thr Asp Lys Lys Ser Leu Asn Ile Gly Asn Gly Val Ile		
130	135	140
His Ala Phe Leu Thr Leu Ile Phe Leu Leu Phe Trp Ile Pro Phe His		
145	150	155
Val Ser Tyr Ile Tyr Pro Ile Tyr Phe Gln Asp Cys Val Ile Phe Tyr		
165	170	175
Ser Ile Val Leu Thr Phe Phe Met Leu Ser Gln Leu Val Thr Tyr Tyr		
180	185	190
Val Tyr Glu Leu Phe Leu Leu Leu Met Leu Lys Ile Ser Trp Asp Lys		
195	200	205
Leu Leu Gly Val Leu Phe Glu Ser Phe Leu Gly Ile Lys		
210	215	220

<210> 111
 <211> 235
 <212> PRT
 <213> Homo sapiens

<400> 111

Phe Glu Asp Lys Phe Leu Leu Thr Val Val Ile Thr Arg Gly Leu Ile		
1	5	10
Ser Thr Leu Leu Glu Ser Leu Thr Tyr His Asn Phe Ser Met Leu Cys		
20	25	30
Glu Gly Met Asn Ser Leu Thr His Leu Ile Met Thr Thr His Ile Met		
35	40	45
Leu Leu Ile Gly Asn Asp Leu Tyr Glu Thr Tyr Arg Lys His Ile Thr		
50	55	60
Ala Ser Gln Met Thr Pro Ile Ser Pro Ile Ala Val Ser Asp Lys Phe		
65	70	75
Glu Ser Gly Pro Met His Leu Cys Trp Ala Pro Gln Asn Lys Glu Val		
85	90	95
Asp Tyr Leu Arg Ser Thr Thr Leu Ala Ile Ser Pro Leu Asn Ile Lys		
100	105	110

00196PCT1.ST25

Leu Ile Cys Pro Ile Ala Pro Pro Ser Ser Gly Pro Gly Leu Trp Ile
 115 120 125
 Gly Met Thr Tyr Leu His Ile Gln Phe Cys Lys Ser Leu Gly Ile Ile
 130 135 140
 Gln Asp Gly Arg Ile Asn Gly Gln Leu Lys Leu Phe Leu Leu Ser His
 145 150 155 160
 Pro Phe Gln Cys Phe Leu Pro Trp Ser Leu Leu Ile Ile Ser Met Leu
 165 170 175
 Phe Asn Ile Tyr Leu Glu Glu Phe Met Ala Val Ile Thr Ile Met Ala
 180 185 190
 Thr Ile Phe Tyr Tyr Leu Cys Met Pro Gly Ile Val Leu Ser Ala Ser
 195 200 205
 Gly Ile Arg Ser Cys Lys Gly Leu Val Thr Phe Tyr Arg Trp Asp Trp
 210 215 220
 Asp Ser Asp Val Ser Cys Leu Phe Lys Ser Ile
 225 230 235

<210> 112
 <211> 134
 <212> PRT
 <213> Homo sapiens

<400> 112

Ser Ser Pro Val Val Cys Trp Gln Ser Leu Ala Phe Leu Ser Leu Trp
 1 5 10 15
 Lys Tyr His Ser Ile Ser Val Leu Ile Ser Thr Trp Cys Ser Ser Cys
 20 25 30
 Val His Val Cys Leu Gln Ile Ser Pro Phe Tyr Lys Asp Thr Val Ile
 35 40 45
 Leu Asp Ser Gly Ser Phe Arg Pro His Leu Ile Phe His Lys Asp Pro
 50 55 60
 Ile Ser Lys Cys His Ile Leu Trp Tyr Trp Gly Leu Leu Lys His Ile
 65 70 75 80
 Asn Phe Arg Glu Thr Asn Leu Asn Leu Gln Tyr Thr Ser Arg Met Glu
 85 90 95
 Glu His Gly Ile Arg Leu Ser Gln Thr Gln Leu Leu Thr Phe Trp Phe
 100 105 110
 Ser Ser Pro Gly Gln Glu Thr Pro Ser Ala Gly Lys Leu Glu Thr Trp
 115 120 125
 Lys Thr Gly Leu Lys Thr
 130

<210> 113
 <211> 229
 <212> PRT
 <213> Homo sapiens

<400> 113

00196PCT1.ST25

His Thr Asp Thr His Ser His Ile His Thr Gln Ser Leu Ile Lys Tyr
 1 5 10 15
 Met Ile Ile Phe Met Cys Lys Ser Phe Gln Gln Ile Ile Ile Phe Tyr
 20 25 30
 Ile Arg Ala Cys Tyr Lys Glu Lys Ile Tyr Gln Phe Glu Lys Gly Lys
 35 40 45
 Pro Leu Ser Arg Tyr Cys Phe Ile Arg Thr Val Val Ser His Ile Ile
 50 55 60
 Ser Lys Leu Leu Met Lys Tyr Lys Thr Phe Thr Ile Ile Lys Ser Leu
 65 70 75 80
 Lys Arg Thr Lys Asn Lys Leu His Lys Leu Lys Ser Ser Val Ala Asn
 85 90 95
 Met Met Phe Cys Glu Leu Leu Ile Val Tyr Val Cys Ile Tyr Ala Trp
 100 105 110
 Tyr Leu Pro Gly Ile Cys Phe Met Phe Leu Arg Pro Gln His Cys Cys
 115 120 125
 Lys Arg Ile Val Phe Pro Leu Leu Tyr Asn Tyr Phe Asp Ile Ser Tyr
 130 135 140
 Asn Leu Pro His Glu Tyr Gln Thr Phe Tyr Arg Lys Tyr Leu Ile Pro
 145 150 155 160
 His Ser Leu Ser Pro Ala Ala Phe His Val Cys Leu Val Lys Ala Ile
 165 170 175
 Val Thr Lys Leu Pro Phe Phe Lys Glu Ala Ser Val Asn Gln Tyr Ile
 180 185 190
 Ser Leu Ser Leu Phe Phe Tyr Val Cys Leu Ser His Thr Asn Thr Gln
 195 200 205
 Ala Asn Ile Tyr Ile Tyr Ile Phe Asn Ile Thr Asp Ser Phe Leu Ala
 210 215 220
 Val Leu Ser Ile Ile
 225

<210> 114
 <211> 189
 <212> PRT
 <213> Homo sapiens

<400> 114

Ser Leu Leu Asn Leu Leu Phe Asn Met Asn Ile Ala Ser Leu Ala Leu
 1 5 10 15
 Phe Val Leu Thr Leu Tyr Ile Thr Phe His Leu Phe Ile Leu Ile Cys
 20 25 30
 Leu Tyr Ile Ser Ala Phe Leu Ile Gly Asn Ile Leu Ser Leu Ser Phe
 35 40 45
 Tyr Pro Ile His Leu Leu Asp Phe Glu Val Phe Lys Leu Phe Val Phe
 50 55 60

00196PCT1.ST25

Asn Val Asn Met Tyr Met Ile Gly Phe Lys Phe Thr Ser Trp Leu Val
 65 70 75 80
 Phe Ser Val Tyr Ser Ile Tyr Tyr Ser Leu Phe Pro Phe Ser Ser Met
 85 90 95
 Leu Ser Phe Gly Leu Ile Ile Leu Leu Lys Ile Phe Arg Ile Ser Phe
 100 105 110
 Val Val Leu Phe Trp Leu Ile Cys His Leu Arg Leu Leu Ile Thr Val
 115 120 125
 Ile Phe Gln Val Thr Leu Tyr His Phe Val His Val Tyr Lys Thr Leu
 130 135 140
 Gln Gln Cys Thr Ser Ile Leu Cys Leu Leu Asn Phe Arg Leu Leu Leu
 145 150 155 160
 Ser Ser Tyr Ile Leu Phe Leu Phe Pro Thr Tyr Val Ile Arg Pro Ile
 165 170 175
 Leu His Cys Phe Cys Val Cys Phe Lys Lys Pro Ser Phe
 180 185

<210> 115
 <211> 242
 <212> PRT
 <213> Homo sapiens

<400> 115

Glu Glu Asn Ser Met Lys Ala Asp Lys Gly Arg Thr Glu Val Asn Gln
 1 5 10 15
 Cys Ser Ile Asp Leu Gly Glu Asp Asp Met Glu Phe Gly Glu Asp Asp
 20 25 30
 Ile Asn Phe Ser Glu Asp Asp Val Glu Ala Val Asn Ile Pro Glu Ser
 35 40 45
 Leu Pro Pro Ser Arg Arg Asn Ser Asn Ser Asn Pro Pro Leu Pro Arg
 50 55 60
 Cys Tyr Gln Cys Lys Ala Ala Lys Val Ile Phe Ile Ile Ile Phe Ser
 65 70 75 80
 Tyr Val Leu Ser Leu Gly Pro Tyr Cys Phe Leu Ala Val Leu Ala Val
 85 90 95
 Trp Val Asp Val Glu Thr Gln Val Pro Gln Trp Val Ile Thr Ile Ile
 100 105 110
 Ile Trp Leu Phe Phe Leu Gln Cys Cys Ile His Pro Tyr Val Tyr Gly
 115 120 125
 Tyr Met His Lys Thr Ile Lys Lys Glu Ile Gln Asp Met Leu Lys Lys
 130 135 140
 Phe Phe Cys Lys Glu Lys Pro Pro Lys Glu Asp Ser His Pro Asp Leu
 145 150 155 160
 Pro Gly Thr Glu Gly Gly Thr Glu Gly Lys Ile Val Pro Ser Tyr Asp
 165 170 175
 Ser Ala Thr Phe Pro Ser Phe Gly Lys Pro Thr Val His Asn Thr Arg

00196PCT1.ST25

180 185 190
 Asn Lys Arg Arg Phe Leu Phe Asn Gly Pro Thr Ile His Cys Gln Thr
 195 200 205
 Ile Pro Phe Gln Ala Lys Val Leu His Thr His Ala Leu His His Lys
 210 215 220
 Val Asp Lys Tyr Ile Glu Glu Ala Gly Thr Gly Val Phe Pro Lys His
 225 230 235 240
 Gly Leu

<210> 116
 <211> 206
 <212> PRT
 <213> Homo sapiens

<400> 116

Ser Gly Lys Thr Thr Pro Arg Asn Arg Leu Leu Leu Pro Pro Cys Lys
 1 5 10 15
 Pro Glu Ala Gln Leu Leu Ser Leu Glu Asn Arg Lys His Asn His Gly
 20 25 30
 Tyr Ser Glu Gly Gln Gly Gln Val Leu Cys Lys Trp Asp Cys Gly Gly
 35 40 45
 Gln Trp Glu Gly Phe Trp Gly Ser Leu Ser Cys Leu Cys Asn Trp Ala
 50 55 60
 Met Gln Pro Cys Lys Cys Gln Glu Thr Leu Asn Lys Thr Glu Pro Glu
 65 70 75 80
 Ala Asn Lys Lys Pro Ala Phe Thr Cys Ser Phe Pro Phe Cys Asn Glu
 85 90 95
 Ile Ser Ile Cys Thr Leu Ile Trp Pro Thr Ile Pro Gly Glu Ile Ser
 100 105 110
 Trp Asp Val Ser Phe Val Thr Leu Asn Phe Leu Val Pro Gly Leu Val
 115 120 125
 Ile Val Ile Ser Tyr Ser Lys Ile Leu Gln Val Cys Phe Leu Gln Val
 130 135 140
 Leu Pro Leu Asn Phe Thr Gln Ala Trp Gly Tyr Phe Cys Asn Leu Arg
 145 150 155 160
 Ile Trp Gly Arg Arg Thr Pro Lys Ser Ser Arg Gln Leu Asn Leu Asp
 165 170 175
 Ser Leu Pro Arg Ser Thr Thr Leu Arg Lys Glu Arg Ile Phe Leu Glu
 180 185 190
 Val Ile Ser Leu Leu Cys Phe Leu Leu Ile Thr Lys Val Ile
 195 200 205

<210> 117
 <211> 9
 <212> PRT
 <213> Artificial Sequence

00196PCT1.ST25

<220>

<221> misc_feature

<223> Peptide substrate

<400> 117

Ala Pro Arg Thr Pro Gly Gly Arg Arg
1 5

Continuation of Transmittal Letter:

- 4(c) The present application claims priority of Application Serial No. 60/195,150 filed 6 April 2000; Application Serial No. 60/195,099 filed 6 April 2000; Application Serial No. 60/195,151 filed 6 April 2000; Application Serial No. 60/195,148 filed 6 April 2000; Application Serial No. 60/195,093 filed 6 April 2000; Application Serial No. 60/195,098 filed 6 April 2000; and Application Serial No. 60/230,149 filed 5 September 2000.